



0...

SE F









ACOBI PETIVERI

HISTORIAM NATURALEM spectantia:

CONTAINING

SEVERAL THOUSAND FIGURES

Birds, Beasts, Fish, Reptiles, Insects, Shells, Corals, and Fossils;

ALSO OF

Trees, Shrubs, Herbs, Fruits, Fungus's, Mosses, Sea-Weeds, &c. from all Parts, adapted to Ray's HISTORY OF PLANTS,

On above Three Hundred COPPER-PLATES, with English and Latin Names.

The Shells have English, Latin, and Native Names.

N. B. Above One Hundred of these Plates were never published before.

To which are now added

SEVENTEEN CURIOUS TRACTS,

Most of them so scarce as not to be purchased,

Which completes all he ever wrote upon NATURAL HISTORY; viz.

Gazophylacium Naturæ et Artis. Musei Rariora Naturæ, &c. 238 Land and Water Beetles. 26 Humble-Bees.

100 \ Sea-Shells, Eggs, Stars, &c. 26 Kentish 25 Sheppey

59 Gloucestershire

112 Merian's Surinam Insects. On Snails and Worms being Hermaphrodites.

Ray's Method of Plants Illustrated.

Herbs of the same Class have similar Virtues. Virginian Cicada, Wasps, Bees, &c.

The Two Volumes containing above Ten Thousand Articles, Engraved in the most Accurate Manner, from Originals, the Gifts of the most eminent Persons in all Nations.

The ADDITIONS Corrected by the late Mr. JAMES EMPSON, of the British Museum.

V O L U II.

CONTAINING

Rudiments of Botany. Ray's English Herbal. South-Sea Italian and Egyptian American Plants. Montpellier

Silesian and Etrurian Plants. 164 British Butterflies, &c. Collectiana Petiveriana. English Grasses, Fungus's, Mosses, &c. Hortus Siccus.

N D O L O N.

Printed for JOHN MILLAN, Bookfeller, near White-Hall.



E

•			
VОL. I.	Pag.	-	Pag.
Irections for Travellers	I	Herbs of the same Class have similar Virtues	29
156. Gazophylacium Naturæ & Artis		10 Virginian Wasps, Bees, &c	30
Catalogus Classicus & Topicus		g and springers	30
22. Above 400 Amboina and East-India		VOL. II.	
Shells, &c	2	, o E. II.	
Musei Rariora Naturæ		2. Rudiments of Botany	
238 English Land and Water Beetles, in		82. Ray's English Herbal and Catalogue;	
	12-18	864 Plants, &c	
26 Humble-Bees, &c. in five Classes -	18	7. Hortus Peruv. Chili, &c. or South-	
28 De Animalibus Crustaceis, Echinatis,	10	Sea Herbal; 133 Plants -	
Stellatisque Britannicis	ib 1	7. Italian and Egyptian Plants, &c. 120	5
19 Fresh-Water Fishes	10.	23. Pteriagraphia Americana; 506 Ferns,	
42 DeCochleis Terrestribus ac Fluviat. Brit.	20		
39 De Pectinibus, Pectunculis, & Ostreis Br.		6. Papilion. Britan. about 164 English	
26 Fossilia Cantiana; Fossils found in the	10.	Butterflies	
Chalk-Pits of Kent	0.1	Monspelienses Plantæ; 460 Plants -	
25 Sheppey Fossils		Plantæ Silesiæ rariores, &c	
59 Fossils at Pyrton-Passage in Gloucestersh.		Plantæ Etruriæ rariores, &c	
112 Merian's Surinam Infects		Cat. Collectania Petiver. of 1080 Articles	
Snails and Worms Hermaphrodites in a	- 1	Gramin. Brit. Concordia; above 400 Eng-	
double Sense	26		
Ray's Method of English Trees and Shrubs	- 1	Hortus Siccus, Chirurgicus, & Pharma-	
Illustrated	26	ceuticus	12

Printed for J. MILLAN, near WHITE-HALL.

- I. Catefby's Hortus Americanus; containing Eighty-Five of the most curious Trees, Shrubs, &c. with their Blossoms, Fruits, and Seeds, on Seventeen large Copper-Plates coloured after Nature; with Observations on their Constitution, Growth, Culture, Uses, and Virtues, adapted to the Climates of Great-Britain, Ireland, and most Parts of Europe, &c. Likewise Directions to collect them in America, and to pack up and secure the Trees and Seeds in their Passage to Europe. 21.2s.
- II. Dillenius's General History of Land and Water Mosses, &c. containing all the known Species, exhibited by about a Thousand Figures on Eighty-Five large Royal Quarto Copper-Plates, their Names, Places of Growth, and Seasons, in English and Latin, referring to each Collected, drawn, and engraved, by the Author. 11.5s.

III. Sheldrake's Herbal, on a Hundred and Eighteen large Folio Copper-Plates, drawn in the most masterly Manner from the Originals when in the highest Perfection.

- IV. Sheldrake on Heat and Cold, for Green-Houses, 1s.
- V. Columella on Agriculture, by Gibson, Quarto, 14s. VI. Letters from a Persian in England to his Friend at Ispahan, 3s.
- VII. Hill's History of the Royal Society, 10s. 6d.
- VIII. List of his Majesty's Land-Forces and Marines at Home and Abroad, &c. by Permission, for 1767, 5s.
- IX. The New Manual Exercise, with Explanations, by his Majesty's Order, 1s.
- X. Manœuvres for a Battalion of Infantry upon Fixed Principles, Command of small Detachments, &c. with Fifty-seven Plates, 10s. 6d.
- XI. Recruiting-Book, proper for all Officers on that Service, 2s. 6d.
- XII. Regimental Book, with proper Heads beautifully engraved, 4l. 4s.
- XIII. General Returns for Reviewing Horse, Dragoons, and Foot, as now given in to his Majesty.
- XIV. Ditto, Monthly and Weekly Recruiting, &c. &c.
- XV. The Majors or Adjutants Book; containing the different Forms of Returns for each Corps,
- by the Adjutant General. XVÍ. A System of Camp-Discipline, Adjutants and Garrison Duty, Regulations for Land-Forces, Kane's Campaigns and Military History from 1660. with many Copper-Plates; now reprinting, 7s. 6d. with great Additions.

A Catalogue of MRAY'S ENGLISH HERBAL Illustrated with FIGURES on Folio Copper PLATES. On which are Engraven Their NAMES to each, as also the Places where Growing. Colour of their Flowers and Time of Flourishing, if Annual or Perennial, Common or Rare. The Flower, Seed, Vessell &c. Figured from y Life & added to each Different Kind, By JAMES PETIVER Fellow of y Royall Society London.

SECT. I. ENGLISH HERBS with Sta = 9 Curl'd Pondweed O. L. Rivulets. mineous, Threddy or Blink FLOWERS. 10 Broad Grass Pondweed O. Slow waters. TAB. I. Fig. 1. Male Hops. O. Flowers in July. KENT &c.

2. Female Hops. O. * Ripe beginning of Sep!

3. Male Hemp. O. Sow'n in Welds. Ft. July.

4. Female Hemp. O. * Seeds in Aug!

Mela Dan's Herrory. O. Ft. March. 5 Male Dog's Mercury. O. Fl. March. woods & hedges. 6 Female Dog's Mercury. O. Seeds Apr. LOND. 7 Male French Mercury. . Gardens Wild. & Temale French Mercury. O. x . Iuno &c. g Great Nettle. O. * L. July Ditches & Dunghills Com. 10 Small Nettle. . L. Com. Rubbish Iun. &c. 11Roman Nettle. . Rare. June. 12 Small Burdock. L. Rare. by Roads. 12 Small Burdock. C.L. Rare. by Road.

TAB. II. 1. Great Water Dock. O. * I. by Sow Rivulets Iul.

2 Curld Sharp Dock. O. I.Wet ground.

3 Smooth Sharp Dock. O. L. Com. Wast & Wet g. Iul.

4 Small sharp Dock. O. L. Wet Ditches. Y.

5 Blood vein d Dock. O. L. Paths Plow d g. Y. 6 Bloodles Dock. O. L. Hedges . Y. 7 Gold Dock. . L. Slow Waters. Y. 8 Bur Gold Dock. . L. Slow Waters. Y. 9 Broad Dock O. L. Com . Wast g. dout Houses . Y. 10 Fiddle Dock O. L. War Fields. Y. "Buck-wheat, O.r.L. Sown in Fields Y. &c White.
12 Trailing Buck-wheat O.L. Corn & Sowed g. Y pale. TAB. I. Common Sorrel O. * I. Meadows. Iune. AB. I. Common Jorrel. S. L. Meadows. Tune.

2 Sheep's Jorrel. O. L. Jandy soil. May.

3 Barren Sheep's Sorrel. O. L. Jand, May.

4 Welsh Jorrel. O. In Rills on Rocks. May. &c.

5 Common Arfmart. S.* L. Wet Ditches. Aug.

6 Creeping Arfmart. S. L. Moist Meads & Aug.

7 Spotted Arfmart. S. L. Corn & Wet g. Aug. Flesh. 8 Spotted hoary Arfmart. O. L. Rare Aug. 9 Silver Arfmart. O.L.Corn Iul.
10 White branched Arfmart. O.L.Rare. " Bobarts Arfmart. O. L. Sand. Corn &c Iul. Pale. 12 Amphibious Arfmart. O. L. Fig. A. Land B. Water. TAB. N. 1 Common Plantain. O. L. In Paths Iun to Aug. 2Dented Plantain O.L. In Paths. 3 Hoary Plantain. O. L. Sandy Paths May &c. 4 Rose and spiked *Plantain*. O. Rare.
5 Besom *Plantain*. O. Rare.
6 Common *Ribwort*. O. *. Pastures. Com. May & c. 7 Mountain Ribwort. O. *. Pastures. Com. May &c.
7 Mountain Ribwort. O. Wales &c.
6 Childing Ribwort. O. Rare.
9 Sea Plantain. O. L. Salt Marshes
10 Bucks-horn Plantain. O. *. Sand & Sea Joore. Int.
11 Great Burnet. O. * Ch. 11. 11.
11 Small Rurnet. O. * Ch. 11. 11. 12 Small Burnet. O. * Chalk-hill Im.
TAB.V. 1 Common Biftort. O. *. L. Moist Meads. Y. pale. 2 Small Bytort O. North. pale 3 Wellh Bytort. O. Rare 4Broad Pondweed. O.L. Boggs & Ponds. 5 Long Pondweed. O. L. Thames. 6 Round Pondweed. O. L. Slow Rivers.

7 Heart Pondiveed. O. Slow Waters.

Plain Pondweed. O. L. Rivulets.

12 Fine Grass Pondweed O. Rivers. 3 Fennel Pondweed O. Brooks TAB. VI. 1 Tassel Pondweed. O. Salt Rivulets. 2 Horned Pondweed. O. L. Shallow Ponds. Iul. 3 Water Starwort. O.L. Wet Ditches. 4 Long Water Starwort. O.L. Ponds & Ditches.
5 Feather Pondweed. O.L. Rivulets.
6 Fine Feather Pondweed. O. Slow Waters. 7 Millfoil Pondweed. O. Low waters. 8 Knotted Pondweed Water Ditches. g Golden Saxifrage. O. L. Shady bogs. Apr. Yellowijh. 10 Long Golden Saxifrage. O. With & Com. 11 Cornwal Pennywort O. Rare. 12 Marsh Pennywort O. L. Bogs & Rills. Iul. &c. TAB.VII. 1 Delt Orrach. . L. Gardens, Dunyhills. &c. Aug. 2Delt Sea Orrach. O.L. Sea Shores. Y.&c. 3 Frosty Sea Orrach. . Sea Shores. ESSEX. &cY&c. 4 Jagged long Orrach. C.L. Sea Marshes & Aug. 5 Spear Orrach. L. Com Dunghills. Aug. 6 Grass Sea Orrach Sea Side. 7 Sea Purflain. O. Salt May hes. Y. 8 Heart Sea Purstain. O, Rave. Sea coasts.
9 Small Garden Blite. Old Gardens. Aug.
10 All feed Blite. O.L. Gardens. Hsp-ground &c. 11 Stinking Orrach. • *L. Dunghills &cY&A.

12 Wild Spinage O.*L. Wast y. Apr. &c.

TAB.VIII. 1 Rands Oak Blite • L. Rubyh. Inl. &c. 2 Frost Blite. . L. Wast Dury &c. Com. Iul. &c. 3 Buddle's Fig Blite. . L. Rubish Inl. &c. 4 Buddle's round Blite. L. Rubish. Wast. &c. Iul. &c. 5 Thick shining Blite. L. Wast Dung & Aug. 6 Sharp pointed Blite. L. Rubish Dung. Aug. 7 Maple Blite. L. Rube Broad pointed Blite. L. Com. Rubish &c. 9 Sea Beet. O. Salt Marshes. 10 Wall Pellitory. O. * L. Walls & Ruors. Iul. &c. 11 Tree Stonecrop. O. Sea Shores. Aug. 2 Shrub Stonecrop. O. Salt Marshes. Aug. TAB. X . 1 Small Glaffwort. . Salt Marshes. Com. 2 Prickly Glasswort. Sea shores. Aug. 3 Joynted Glaffwort. Sea side. Com. 4 Woody jointed Glasswort. O. Shepey Isle &c. 5 Bastard Toad-slax. O. Chalk hills. Ind. &c. 6 Upright Knawell. • L. Sandy gr. d Inl. &c. o Upright Knawell. ■ L. Jandy gr. a Iul &c.

7 Spreading Knawell. ■ L. Corn & Liul &c.

8 Pepper Grafs. L. Shallow Ponds. Iul &c.

9 Ladies Mantle. ○. * Hilly ground &c.

10 Small Ladies Mantle. ○. Rare.

11 Silver Ladies Mantle. ○. Rare. North Mount. 12 Parfly Piert: *. L. Corn & Gravell Iun &c. TAB. X. I Common Knotgrafs. . * . L. Wast gr. 4 Paths. &c Com. hig Thick fet Knotorafs. O.L. Rubish Aug '&c. 3 Short leaved Knotgrafs. . L. Wast gr. 4 Aug. 4 Narrow Knotgrafs. . L. Corn &c. Sea Knotorafs. O. Sandy Sea Shores. 6 Newton's Knotyrafs. on Seaver 7

Naked FLOWERS . THISTLES &c. TAB. XI. Fig. 1 Comon Moule - ear. O. * L. Dry Parturer. It Y. Yell 5 Narrow smooth Moule-ear. O. WALES. Yell. 6 Pluckenette Moule-ear. O. Yell. 7 Comon Dandelion O. * L. Paftures &c Com. Vell all Sumer. 8 Narrow Dandelion O.L. Brick walls &c. Yell. 9 Comon rough Dandelion. O. L. Maul &c. May. Yell. 10 Sandy rough Dandelion. O. Dry banks. Yell. 2 Hemp Agrimony. O. R. L. River sides. Y&A. uRough branched Dandelion. O. R. L. Pastures & Com. IY&A Yelz American Live-long: O. Rure. Y&A. White. 2 Smooth branched Dandelion. . Yell. TAB XIL 1 Comon Hawk-bit O. L. Patures. Y& A. Yell. 2 Jugged Hawk-bit O. L. Pafures. Iul. &c Yell. 3 Small jagged Hawk-bit. O. L. Yell. 4 Dandelion Hawk-bit. Q. L. Yell 5 Buddle's Hawk-beard. O. L.Y& A. Yell. 6 Comon Hawk-bead. D. L. Pafaras. Y& A. Yell. 7 Succory Hawk-beard. L. Parwas. Y& A. Yell. 8 Caftor Hawk-weed. . L. Chalk pits &c. I.Y& A. Yell. 9 Small Succory Hawk-weed. L. Yell. o 10 Tall Succory Hawk-weed. RENT. YEA. Yell. n Curld Hawk-weed.Q.L. Field sides. Yell. 12 Ox-tongue Hawk-weed Q. L. Ditch banks. Y& A Yell. TAB, XIII. 1 Broad Hawk-lung O. Rare Yell. 2 Round Hawk-lung. O. Rare Yell.

3 Broad dented Hawk-lung. O. Rare Yell.

4 French Hawk-lung. O. K.L. OU Walls. Woods. &c. I & Y. Yoll. 5 Narrow Hawk-lung. O.A. Val. Wall. Wood 6 Welfh hoary Hawk-lung. O. H.Y&A. Yell. 7 Broad hairy Hawk-lung. O. H.Y&A. Yell. 8 Long hairy Hawk-lung. O. H.Y&A. Yell. gBroad smooth Hawk-lung. O. Rare. WESTMERL. Yell. 10 Narrow hairy Hawk-lung. O. H. Woods. Yell. "Narrow smooth Hawk-lung. O.H. Woods. Yell. 12 Grass Hawk-lung. O. Sondhills. Yell.
TAB. W. 1 Round leaved Sowthistle. . L. Corn. Yell. hon.

3 Narrow Sowthistle. . Rare. Yell. 4 Prickle dented Sowthistle. . L. Banks & Lell. I & Y. 5 Prickle jagged Sowthistle. . * L. Banks & Yell I & Y. 6 Corn Sowthiftle. O. K.L. Banks. Corn. YKA. Yell. Tall Marsh Sowthistle. O. L. River-banks. Aug. Yell. 9 Smooth jagged Sowthiste. . *L. Garden &c. IY&c Yell. 10 Buddle's jugged Sowthiftle. O. L. Gardens. IY& A Yell. " Wild Succory. O. r.L. Jul & dug. Field paths Blue. 13 Nipple - Wort. O. *. L. Gard Tilled gr. 4 IY&A. Yell. TAB. W. 1 Broad Lettice. O. L. Field banks. Y& A. Yell. i Cutt Lettuce. . L. Ditch banks. IYka Yell. 3 Jugged Lettuce. . A. Stoney hedges Gard &c. Y&A. Vell. 4 Small jagged Lettuce. . L. 1&Y. Duch bunks. Yell. 5 Ly Lettuce. O. H. Walls. Shady banks. Y&A. Vell. 6 Narrow Goats-beard. O. L. Meads. I.Y &c. Yell. Comon Goats - beard. . *. H. Mead. I.Y& a Yell. & Purple Goats - beard. . Rare Gardens. 9 Small Swines Succory. O. L. Sand Corn. In &c. Yell. to Comon Carline . L. Inl&cDry partures. & Heath Yell.A n Butter-burr. O. * L. River fides. &c. 12 It' Flower. Apr. & May. Flesh. TAB. XVI. 1 Elecampane O. * Mout Mead & I&Y Yell. 2 Comon Fleabane O. +. L. Rill Ditha Y & A Vell. 3 Small Pleabane O.L. Dry Pond Y&A. Vell.

ABlew Fleabane O. L. Dry Patures. Y& A. Purphin. 5 Marsh Fleabane . Fone & Lun & Y. Yell.

6 Jagged Fleabane . Fen. &c. Ium &Y. Yell.

2 Broad Sowthiftle. O. Gard Dungh I & Y. Yell.

7 Bobart's Golden-rod. O. River Sides. Rare. Y&A. Yell. & March Golden-rod O. Err Fens &c. Yell. g Narrow Golden-rod. O. Rare Yell. 10 Comon Golden-rod . . H. Woods. Heaths . Y& A. Vell. 11 Welh Golden-rod O. Mount Yell. 12 White Golden-rod. Gard. Walls. Aug. Pale. TAB. INI. 1 Comon Ragwort O. * . Pafaves &c Com Y. A &c Yell. 2 May/h Ragwort O. Wet gr. Meads Y& A. Yell. 3 Grounds ell Ragwort. O. H. Hedy a. Fields Aug. Vell. 4 Cambridge Ragwort. O. Heath. &c. Inn. Yell. Comon Groundfell . * . L. Till gr. Wast &c Com. Feb. to Nov. 5 Comon Groundfell. . H. Sandy Soil. I.Y & Vell. 7 Colts-foot. O. * L. Rills. Wet gr: 8 It's Flower. March & Apr. Yell. 9 Golden Samphire. O. Jea Coa/t Yell. 10 Great Sea Starwort. O. Y. L. Salt Mar hos. Y& A. Redish. 11 Small Sea Starwort. O. L. Salt Marshes. Y&A. Redish. 12 Naked Sea Starwor t.O. Salt Marshes BRISTOLL &c. Yell TAB. XXIII. I Montpelier Fleabane. O. A. L. Dry Hills. Banks Y& A. 2 Hemp Agrimony. O. t. L. River sides Y& A. Flesh. 4 Cats-foot. O. Mount. Heaths. Wh. & Flesh 5 Jersey Live-long. . Banks. Walls. Rare. Pale. 6 English Live-long. O. H. Broom & Sand fields. Y & A. 7 Long Cudweed. O. L. Dry Ponds & c. Com. Blackish. 8 German Cudweed. . Sandy ground &c. 9 Childing Cudweed . L. L. Paths. dry ponds. Y& A. Pale. 10 Low Childing Cudweed. Barren gr. Y& A.Palc. 11 Least Cudweed. M. Sand Heaths. Corn. Y& A. Pale. 12 Grafs Cudweed . Corn Sandy gr. Pale TAB. XX. 1 Great Dalie O. x. L. Field sides. &c I & Y. Wh. 2 Comon Dalie O *.L. All Meads &c Apr. to Aug. Wh. Sneezewort. O. r. H. Wet ground Y& A. Wh. 4 Yarrow. O. *. L. All Parturas. Com. I.Y&c. Wh. & Bluth. 5 Feverfew. O. *. L. Nat Hedga. Y& A. Wh. 6 Corn Marygold. O.L. Corn. Y&A. Yell 7 Yellow Ox-eye. O. r. River Sides. Rare. Y & A. Yell. 8 White Ox-eye.Q.L. Meads &c. May &c. Wh. g Bitter Camomile. L. Tilld gr. Wh. 10 Sweet Camomile O. * L. Wet Comons. Aug. &c Wh. 11 Stirking Camomile. . * L. Wast & Tilld. gr. 3 Y& A. Wh. 12 May-weed. . x.L. Wast Fields. Y& A.Wh. TAB. X. 1 Comon Wormwood O. * L. Wast gr. A &c. Yell! 2 English Sea Wormwood. O. *. Salt Musher. Aug &c Yell. A. 3 French Sea Wormwood Q r. Salt Marshes Aug &c Yellsh 4 Wild Southernwood. O. Way sides. Rare. Y & A. 5 Mugwort. O. *. L. Field sides. Hedges. LY&A Whiteifh, 6 Bwr Marygold. . . I. By Ponds &c A& Sep. Vell. 7 Trifid Burr Marygold . L. By Ponds &c A & Sept Yell. 8 Sea Cottonweed. O. Sandy Thores. Rare. Yell. 9 Tanley. O. *. L. Field banks &c Aup &c Yell 10 Curlet Tanley. O. Gardens Aug &c Vell. 11 Agrimony. O. * L. Hedges. Field sides. Y& A. Yell. 12 Enchanters weed O.L. Shady Woods &c.Y&A. Wh. th TAB. XXI. 1 Mu/k Thiftle. L. Flinty, burren gr. Y& A.Red.
2 Welted Thiftle. L. Wast Ditcher Y& A.Red.
3 Gray Thiftle. L. Wast banks. I & Y.Blush.
4 Marsh Thiftle. H. Wet gr. Bogs. YA & c. Red. Comon Thifle O . L. Paths Fields &c. Com. Y& A.Red. 6 Dwarf Thistle. Q. L. Stoney dry hill Y& A.Red. 7 Spear Thiftle Q.L. Wast Banks & I.Y & Red. O Fryers Thiftle Q. Hills & Plains Rare I.Y & a Red. 9 Milk Thistle . * L. Banks . Field Sides . Y& A. Red. 10 Cotton Thiste . * L. Path, Banks & c. Y& A. Red. Com. 11 Star Thiste . * L. Wast Paths & Aug & Red. 12 Yellow Thiftle. D. Ry Hedges &c. Rare. Y& A. Yell. TAB. XXII. 1 Peckam Thille O. L. Wet Meads &c.Y& A. Red. 2 Northern Thiftle O. Mountains Rare. Y & A. Red. 3 Saf Flower. . *. Sowed. Y& A &c Saffron 4 Blue Bottle. . *. L. Corn Ivi & c Blue 5 Broad Saw-wort. O. H. Woods Y & A. Red. 6 Comon Saw-wort. O. H. Wood. Y &A. Red. 7 Matfellon. O. L. Corn &c Y &c Red. 8 Comon Knapweed. Ork. Pastteres &c. Com. I.Y &c. Red.

9 Jugged Knapweed. O. L. Partures &c. LY&c. Red.

10 Comon Scabious. O. * L. corn &c. I.Y & c. Flesh.



```
n Field Scabious. ⊙. Dry hilly gr. dug. Flash.1..
12 Devills -bit. ⊙. *. Woody Pafur es. Aug. Blew. H.
TAB.XXII. 1 Burdock. ⊖. *. L. Y & A. Com. Redish.
       AB.XXIII. 1 Burdock. O. * I. Y&A. Com. Redish.

2 Rose Burdock. O. L.Y. & Aug.

3 Small-head Burdock. O. L.Y & A. Com. Rubijh.

4 Small woolly Burdock. O. L.Y & A. Com. Wast gr. 4

5 Great woolly Burdock. O. L.Y & A. Com. Hedge Paths.

6 Cobweb Burdock. O. Mont. Y & A.

7 Manured Teafel. O. * Sowen. Y & Aug. Flesh.

8 Wild Teafel. O. * I. Het ditches. Y. & C. Flesh.

9 Small Teafel. O. * L. Het ditches. Y. & A. Palo.

10 Sea Eringo. O. * Sandy Shores. Y & A. Palo.

11 Comon Eringo. O. Rare. Aug. White. h.

12 Sanicle. O. * H. Woods & Hedges. May. & C. White.
               SECT. III. Um belliferous PLANTS,
               Star-foils. &c.
```

```
TAB. XXV. 1 Cow-Parinep. J. L. Noist gr. 2 Iun. &c. White.
2 Jagged Cow-Parinep. Meads. Ium. &c. White.
3 Garden Parinep. Jown Jul. &c. White. *
4 Wild Parinep. L. Jul. &c. Yell. *
5 Great Hartwort. Rare. Y&A. White.
6 Small Hartwort. L. Rare. Y&A. White.
7 Hogs Fennel. O. * Jalt Marshes. Y&C. Yell. *
8 Sampire. O. Sea Rocks. Aug. Vell. *
9 Angelica. . * Gardens. &c. Y&A. White.
10 Water Angelica. . H. Wet places. Y&A. White.
  10 Water Angelica. H. Wet places. Y&A. White.
11 Coriander. S. Sowin White.
12 Alexanders. S. * Ja Rocks. I&Y.
TAB. XXV. 1 Venus Comb. L. Corn. May. &c. White.
                                2 Cow-weed. O. L. Hedges. Orchards. May. &c. Wh.
3 Wild Chervil. O. L. Hedges. Com. ds. May. &c. Wh.
4 Spignel. O. * WALES &c. Iul & A. White.
5 Water Dropwort. O. L. Wer Moads. &c. White.
6 Small Water Dropwort. O. L. Shall. Ponds &c. White.
7 Henry Lock Dropwort. O. L. Disserberg. I. Van W.
                                  7 Hemlock Dropwort. O.L. River banks. Iul &c. Wh.
8 Narrow Dropwort. O.Rare. Iun. &c. White.
9 Earthnut. O. H. Jandy 91. Iun. &c. White.
10 Goutwood. O. K. Gardens & Hedges. White.
11 Bastard Stone Parsley. O.L. How thickers. Y&A. Wh.
go Courtweed. O. K. Earden & Hedger. White.

11 Bustard Stone Parsley. O. L. Moist Ditches. Y&A. Wh.

12 Hone wort. O. K. L. Clay & Corn. Aug. White.

TAB. XXXI. 1 Broad water Parsnep. O. L. Rivers. Ind. & Wh.

2 Navrow water Parsnep. O. L. Brooks. Inn. & Wh.

3 Creeping water Parsnep. O. L. Brooks. Inn. & Wh.

4 Leaft water Parsnep. O. L. Weedy Ponds. Ins. & Wh.

5 Great broad Saxistrage. O. & By Woods. Ins. & Wh.

6 Great Cut Saxistrage. O. By Woods. Ins. & Wh.

7 Small broad Saxistrage. O. L. & Jandy gr. d.

8 Small Cut Saxistrage. O. L. & Jandy gr. d.

9 Cornwall Saxistrage. O. Rare.

10 Carawaies. O. * Marsh Meads. Ins. & Flesh.

11 Scotch Parsley. O. Sea Rocks. Wh.

12 Smallage. O. * L. Jean Marshes. & Ditches. Aug. Wh.

TAB. XXVII. 1 Comon Fennel. O. * Sea cliffs, banks. Aug. Yellsh.

2 Rock Parsley. Rare White.

3 Mountain Stone Parsley. Rare Wh.

4 Comon Carrot. * L. Jean gr. Ins. & Wh.

6 Fines-Caved Bur-Parsley. O. Corn. Ins. & Redish.

7 Broad jagged Bur-Parsley. O. Corn. Y& A. Redish.

8 Fines-Caved Bur-Parsley. O. L. Rare. Corn. Blush.

9 Rough Parsley. O. L. Thickets, Hedges. Y& Ellush.

10 Corn Rough Parsley. O. L. Crn. Y& Wh.

11 Knotted Rough Parsley. O. L. Field banks. Wh.

12 Fine Rough Parsley. O. L. Field banks. Wh.

13 Lefter Hemlock. A. L. Gom. Ins. & Wh.

4 Comon Hemlock. & L. Gan. & Corn. Ins. & Wh.

4 Comon Hemlock. O. L. Mud ponds. Ins. & Wh.

5 Meadow Saxistrage. O. * H. Meads. Ins. & Wh.

5 Meadow Saxistrage. O. * H. Meads. Ins. & Wh.

6 Prickley Sampire. O. Sandy Sea Jh. Wh.

7 Least Hares-ear. O. Rare. Ins. & Vellsh

6 Thorow-wax. O. L. Corn. Ins. & Vellsh

6 Thorow-wax. O. L. Corn. Ins. & Vellsh

6 Thorow-wax. O. L. Corn. Ins. & Vellsh

6 Comon Borage. O. * Garden & C. Blue.

10 Corn. Thorow. & Vellsh

11 Comon Borage. O. * Corn. Sea Blue.
    6 Prickley Sampire. O. Jandy Joa Jh. Wh.
7 Least Hares-ear. R. R. v. Iul & Vell. h

8 Thorow-wax. D. L. Corn. Iun & Vell. h

9 Comon Borage. .* Sardens & Blue.
10 Small wild Buglofs. .* L. Corn & May & C. Blue.
11 Great Hounds-tongue. .* L. Nat & C. Iun. Purple.
12 Green Hounds-tongue. .* L. Jhady Road. I. & Y. Pwp.

TAB. XXX. 1 Vipers Buglofs. .* L. Barren pl. Com. Y. Pwp.
2 Small Vipers Buglofs. .* Corn & Jand Iul & c. Pwp.
3 Jea Buglofs. .* Sea Shores. Blue.

4 Long Buglofs-Cowflips O. Rare. Purplift.
5 Comfry. O.* L. River banks. May & C. Whor Purplift.
6 Blew Gromill. O. Wales & Blue.
7 Shop Gromill. O. Wales & C. Blue.
8 Bastard Gromill. .. L. Corn. Wast & C. Iun & C. Pale.
8 Bastard Gromill. .. L. Corn. Wast & C. May & C. Wh.
9 Water Scorpion-grafs. .. L. Fallow gr. May & C. Blue & Yell.
10 Field Scorpion-grafs. .. L. Fallow gr. May & C. Blue & Yell.
11 Small Scorpion-grafs. .. L. Jand banks. Walls. May & C. Blue & 12 German. Madder. .. Rare. Purplift.

TAB. XXX. 1 Crofs-wort. or Mugweed. .. * L. Nood hedges. May & C. & Garden. Madder. .. * Rare. Purplift.

TAB. XXX. 1 Crofs-wort. or Mugweed. .. * L. Nood hedges. May & C. & & Garden. Madder. .. * Rare. Purplift.

TAB. XXX. 1 Crofs-wort. or Mugweed. .. * L. Nood hedges. May & C. & & Garden. Madder. .. * Rare. Purplift.
```

3 Wild Madder. O. Rare. Jun &c Yell-h

```
4 Great Smooth Madder O. L. Holger & I. & Y. White.
5 Small Smooth Madder O. L. By Ponds. I un & Wh.
6 Rav. Small Madder. H. Heath & c. Iul & c. Wh.
7 Crokwort Madder. NORTH. White.
8 Vellow Bedfiraw. O. * L. Dry Junny pl. Yell.
9 Woodruff O. * H. Woods. May & c. Wh.
10 Blow Field Madder. L. Corn. Purplift.
11 Cleavers or Goofe-grafs. * L. Hedges. Com. Wh. 1
12 Squinancy-wort. O. L. Chalk-hills. Iul & Elesh.
```

SECT. IV. Verticillate or Whirl PLANTS. &c

```
TAB. XXI. 1 Comon Mother of Thyme. O.*.L. Dry gno Y&cPur. 2 Large flowred Mother of Thyme. O. Dry hill. Y&cPurp. 3 Upright broad Thyme. O. Rave. Aug. Purp. 4 Lemon Thyme. O. **. Hilly gno I um &c Purp. 5 Water Whirl-Mint. O. L. By Pond. Y&c Redish. 6 Comon Water Mint. O. L. Wet dither. Rills.
6 Comon Water Mint. O.I. Wet ditches. Rills.
7 Red Mint. O.**. Iul & c. Rediff.
6 Cross Whirl-Mint. O. Iul & c. Rediff.
9 Essex Pepper Mint. O.Aug. Rediff.
10 Eales Pepper Mint. O.**. Aug. Rediff.
11 Long Horse-Mint. O.Aug. Pale. By Brooks.
12 Round Horse-Mint. O. River sides. Aug. Pale.
12 Round Horse-Mint. O.**. L. Iun & c. Hedges. Flesh.
12 Penny-Royal, or Pudding grass. O.**. L. Wet Ponds. Aug.
13 White Horehound. O.**. L. By Ways. Wast grid Iul & c. Wh.
14 Stinking Horehound. O.**. L. Hedges. Com. Y& A. Purp.
15 Water Horehound. O.**. L. Brooksides & c. Y& c. Wh.
16 Wood Bitony. O.**. H. Woods & c. Iun & c. Redish.
17 Hedge Nettle. O.L. Hedges & c. May & c. Purp.
             7 Hedge Nettle . O. L. Hedges & May & Pup. O English Clary. O. r. L. Stone & Gravel, Y& A. Blue. 9 Great Wild Basil. O. L. Thickets & 28 A. Pup.
   10 Small Wild Bafil. O. L. Dry Chalk hills & L. Iun & L. Pwp.
11 Comon Self-heal. O. *. L. Pwtw & & L. I & A. Pwp.
12 Base Horehound. O. Hills. Padw. Iun & Pwp.
TAB. XXIII. 1 Ground Ly, or Ale hoof. O. *. L. C. Hedger. Blue. Apr.
           AB. XXIII. 1 Ground by, or Ale Hoot. O.* L. C. Hedger Blue. Apr. 2 Red Arch-Angel. . *L. Com. Hedger & C. Apr. May & C. Red. 3 Red cut leav'd Arch-Angel. . L. Eard & Corn. May. Red. 4 Great Henbitt. . L. Fields & Gardens. Purp. 5 White Arch-Angel. O.* L. C. Hedger & C. Apr. & C. Wh. 6 Vellow Arch-Angel. O. H. Shady Hedger. Woods. May & C. Vell. 7 Baftard Balm. O. Woods. Rare. Rediff. 8 Purple Nettle-Hemp. . H. Corn & C.Y. & Aug. Purp. 9 Clowns All Heal. . X.L. By Brooks & Liul & C. Purp.
              10 Yellow Ironwort. O. Corn.Rare. Iul. Yell.
11 Narrow Ironwort. O. L. Corn.Iuly. Red.
12 Round Ironwort. O. L. Corn.I & Y. Redich.
   TAB. XXXIV. 1 Comon Calamint. O.*. L. Paths & Hedges. I & Y. Fish.

2 Field Calamint. O. *. L. Paths. Hedges & CY& A. Flesh.

3 Comon Bugle. O.*. L. Woods. Wet Meads. May & Blue.

4 Mountain Bugle. O. Rare. WALES. Blew.
             5 Motherwort. O. *. Wast Gardens. Y&A. Pwp.
6 Scordium. O. *. Fens. Rare. Pwp.
7 Wood Sage. O. r. H. Woods. Y&A. Pale.
8 Wild Marjoram. O. *. Hedges. Thickets. Blush Y&A.
             g Ground-Pine . . *. Corn & c. I & Y.
to Hood-Mint . O. L. River sides. Iul & c. Blue .
    11 Small Hood-Mint.O.H.Bogs.Aug.Bhsh.
12 Vervain .O.*.L.Paths.Walks.Y&A.Flesh.
TAB. XXXV. 1 Small Toad-Flax. O.L. Corn &c.I &Y.Bhssh.
             AB. XXX. I Small Toad-Hax. W.L. Corn & C.I & I. Blush a Small Snap-Dragon. Q.L. Corn & C.I & Y. Redish. 3 Pointed Fluellin. Q.L. Corn & C.Yell. & Purp. I & Y. 4 Round Fluellin. Q.L. & Corn & C.Yell. & Purp. I & Y. 5 Comon Toad-Flax. Q. * L. Burren fields. Aug. Yell. 6 Blewish weet Toad-Flax. Q. Hedges. Rare Y & A.
  oBlewyn. wee Load-Flax. . Hedger. Rave. 1 & A.
7 Butterwort. O. Begs &c. 1 & Y. Blewijh.
8 Purple Fox glove. O. * H. Wurrens &c. Y & Aug.
9 Brown Fig-wort. O. * L. Thickets &c. Y & A. Red.
10 Water Fig-wort. O. X. L. Water sides. 1 & Y. Red.
11 Jage Fig-wort. O. Rave.
12 Comon Henbanc. O. * L. Wast Rubijh &c. I & Y. Pale.
TAB. XXVI. 1 Quick in hand. Touch me not. O. Rave. Y & A. Yell.
             2 Yellow Rattle. . L. Pafwer Jun &c.
3 Tall Red Rattle. L. Marshes &c Int. Red.
4 Comon Red Rattle. . H. Wet Heaths &c. Inn. Y&A.
5 Comon Eye-bright. . *L. Pafwer &c Y. to Oct. White.
6 Yellow Marsh Eye-bright. . Rare.
              7Rcd Eye-bright. . H. Corn. Wet on I&Y. 8 Or ton Red Eye-bright. Rare. 9 Comon Cow-wheat. . H. Woods &c. I&Y.
 10 Crested Cow-wheat. • Woods & Purp & Yell.

11 Great Bladder-shout. O.L. Bogs & Fens Yell.

12 Small Bladder-shout. O. Rave Moors & L. Yell.

TAB. XXXVI. 1 Purple Violet. O. * L. Gardens & March.
              2 Double Violet. O. Rare. Gardens &c. 3 White Violet. O. Hedges &c. 4 Dogs Violet. O. H. Woods & c. Apr. &c.
            5Bog Violet. O. March &c. o Yellow Rock Violet. O. WALES &c1 &Y.
7Hair Wood Violet: O. Wood sides.
8 Garden Pantie. S. K. Corn &c1 & Y. Mixt.
                                                                                                                                                                                                                                                                      9
```



o Corn Pansie. D.L. Corn & al & Y. Vell & Purple.
10 Great Yellow Pansie. O. WALES. high Bogs. May &c.
11 Base Dyers-weed. D. Chalky 91.3 Inl & Pala.
12 Comon Dyers-weed. D.L. N. Wall. & LY & Aug. Yelloh

SECT. V. Bur & Starheaded PLANTS.

TAB. XXXXII. 1 Pilewort. O. * I. Wet Neads & a M & Apr. Yell.

2 Sweet Wood Crowfoot O. H. Woods. Apr. Yell.

3 Upright Meadow Crowfoot. O. L. Meads. Com. May & a. Yell.

4 Bulbows Crowfoot. O. L. Meads & a Com. M& Ium. Yell.

5 Pale upright Crowfoot. O. L. Dry Ponds. May & a Yell.

6 Doody. byid Crowfoot or B. Banks & a May Yell. 7 Creeping Crowfoot.or Buttercups . Wet Meads. Com. May. 8Thames Crowfoot & L. River fides. Com. May & & Yell. 9 Ray. Hairy Crowfoot. L. Ditch banks. May. Yell. 10 Corn Crowfoot. L. Corn. May & Vell. 11 Manual & Crowfoot. & L. Pond & Com. M. I. Yell. TAB. XXXX. 1 Water Crowfoot, M. J. May &c. White.

TAB. XXXX. 1 Water Crowfoot, with various leaves O. L. Wh.

2 Fine Water Crowfoot, May &c. White.

3 Fine Trimed Water Crowfoot, O. L. Muddy Ponds May &c. Wh.

4 Fennel Water Crowfoot, O. Ponds &c. M. I.Y. Wh.

5 Great Spear—wort, O. L. Fens &c. I. &Y. Yell.

6 Small Spear—wort, O. L. Wet Mead &c. I.Y. A. Yell.

7 Mous for Table O. I. Constituted the A. S. W. Y. T. A. W. C. T. Mous for Table O. I. C. Wet Mead & &c. I.Y. A. Yell. 5 Small Spear-wort. O.L. Wet Mead & LYA. Vell.
7 Mouse Tail. I. Corn. Wet gr. & LA & May. Vell.
8 Adonis Flower. Red Maithes. Corn. Inn. & L.
9 Comon. Mallow. O. * L. Ditther & L. M. IX. A. Purp.
10 Dwarf Mallow. O. * Sandy Wart gr. Y& A. Flesh.
11 Marsh Mallow. O. * Salt Marshes. IX & Flesh.
12 Vervaine Mallow. O. * L. Field & L. X. A. Purplish.

TAB. XL. 1 Avens, Herb Bennet. O. * L. Woods. Hedges Yell. 2 Great flowred Avens. O.R.w.e.May &c Vell.
3 Mountain Avens. O. WALES &c. Redift. May &c.
4 Childing Avens. O. Rave. M & I. Redift.
5 Conquefoil Avens. O. SCOTLAND. Rave. Yell. 4 Childing Avens. O. Rare. M & I. Redish.
5 Cinque foil Avens. O. S'COTLAND. Rare. Yell.
6 Germander Avens. O. IRISH MOUNT. Wh.
7 Comon Strawberry. O. H. Woods. May & wh.
8 Barren Strawberry. O. H. Banks & Apr. Wh.
9 Wood Anemony. O. H. M& Apr. Blush.
10 Pasque Flower. O. H. wood Hedges. Aug to Dec. Wh.
12 Plain Travellers Joy. O. L. Wood Hedges. Aug to Dec. Wh.
12 Plain Travellers Joy. O. L. Same &c.

TAB. XII. 1 Comon Marsh Cinque foil. O. Bogs. Purp.
2 Irish Marsh Cinque foil. Bogs. Rare.
3 Comon Cinque foil. O. *L. Banks & CY& A. Yell.
4 Small rough Cinque foil. O. YORKSHIRE. Yell.
5 Tormentil Cinque foil. O. Nondy gr. I Inn & C. Yell.
6 Upright Welsh Cinque foil. O. Mount Yell.
7 Scotch Cinque foil. O. Woods. Rare. Yell.
9 Tormentil. Septfoyl. O. *H. Heaths. Woods. Y. &c. Yell.
10 Plots Tormentil. O. Corn & C. Y. A. Yell.
11 Silver weed wild Tansey. O. *L. Dry Ponds & Y. A. Yell.
12 Moscatel. O. L. Shades & Mar & Apr. Greenish.

TAB. XII. 1 Orpine, or Live-long. O. *L. Shadey Hedges. I & Y.
2 Rose-Root. O. *Mount Bare. I & Y. Redish.
3 Navel-wort. O. *. Old Walls. Rock. Y& A. Pale.
4 Houseleek. O. *L. On Houses & C. I. Y. Blush.
5 Comon Yellow Sengreen. O. I. Houses & M. H. Heaths. W. H. I. & Y. H 3 Navel-wort. O.*. old Walls. Rock. Ye. A. Pale.
4 Houseleek. O.*. L. On Hower & L. I.Y.A. Bluyh.
5 Comon Yellow Sengreen. O. L. Hower. Walls. I & Y. Yell.
6 Crooked Yellow Sengreen. O. Walls & L. I & Y. Yell.
7 Purple marsh Sengreen. O. Rock. Rare Y & A. Yell.
9 Stonecrop. or Wall Pepper. O. Hower. Walls. Imm. Yell.
10 Mild white Stonecrop. O. Rocks. Apr & Bluyh.
11 Green Bares-foot. O. CAMBR: & JUSTEX Apr. Green.
12 Setter wort. O. Hedge. Rare. Febr. & Greenish.

TAB. XIII. 1 Marsh Marygold. O. L. Wet Meads. Apr & Vell.
2 Globe Crowfoot. O. Hilly Meads. NORTH. Vell.
3 Water Gladiole. O. Muddy Rill. & CI & Y. Floth. 1AB. XIM. 1 Marsh Marygold O.L. Wet Meads. Apr & c. Vell.

2 Globe Crowfoot O. Hilly Meads NORTH. Vell.

3 Water Gladiole. O.Muddy Rill. & c.I & Y. Flesh.

4 Columbines. O. & Wood & Callue.

5 Star water Plantain. O. L. Ponds I & Y. Flesh.

6 Broad water Plantain. O. & L. Ponds & c.I & Y. Flesh.

7 Narrow water Plantain. O. L. Ponds & c.I & Y. Flesh.

9 Grass Arrowhead. O. Rivulets.

10 Broad Arrowhead. O. L. Rivers & c.I & Y. Wh.

11 Narrow Arrowhead. O.L. Rivers & c.I & Y. Wh.

12 Small Arrowhead. O.L. Rivers & c.I & Y. Wh.

13 Small Arrowhead. O.L. Rivers & c.I & Y. Wh.

14 Small Arrowhead. O.L. Rivers & c.I & Y. Wh.

15 Small Arrowhead. O.L. Rivers & c.I & Y. Wh.

16 Small Arrowhead. O.L. Rivers & c.I & Y. Wh.

17 Small Arrowhead. O.L. Hedges & C. Pale.

3 Hack Briony. O. X.L. Hedges & C. Pale.

3 Herb Christopher. O. Woods & C. Whitish.

6 Sweet Solomons-seal. O. * Woods & C. Whitish.

6 Sweet Solomons-seal. O. Rocky Woods. May & c.

7 Lilly - Convally. O. * H. Woods. Heaths. May & c. Wh.

8 Herb Paris. One Berry. O. May & c. Greenish. Woods & c.

9 Bitter sweet. O. * L. By Ponds. I& Y. Blue.

10 Conton Nightshade. O. * L. Doughill & C. Wh. Iuly.

11 Dwale Deadley Nightshade. O. Rubish & c. Brownish.

12 Wild. Sparagus. O. Sea Marshes. Pale.

SECT. VI. PLANTS with 4 Leav'd Flowers.

Flowers.

TAB. XI.V. 1 Jack by & Hedge . . * L. Bank. I & Y. White.

2 Treacle Wormfeed . . * Fen. Int. & Yell.

3 Wall-Flower, or Stock . O. * L. Wall. Apr. & Yell.

3 Wall-Flower, or Stock . O. * L. Wall. Apr. & Yell.

4 Sea Stock Gylly Flower . • Welfh coafes & Ye. A Purp.

5 White Thorow Colewort . • Rare. Jun. & Wh.

6 Sea Colewort . • Dover Cliffs Yell.

7 Round Turnep . • * L. Fields Sowen Yell.

9 Wild Navew . • X. L. Banks Com Apr. M & I Yell.

10 White Mustard . • X. L. Ditches & L. I. Y. Yell.

11 Wild Mustard . • X. L. Ditches & L. I. Y. Yell.

12 Wild Mustard . • X. L. Corn. Com. Int. & Xell.

TAB. XIM. 1 Winter Crefs . • Apr. Yell.

3 Hedge Mustard . • * L. Wast gr. & Yell.

4 Broad Hedge Mustard . • L. Banks & May & I. Yell.

5 Water Rocket . • L. Wet Ditches I& Y. Yell.

6 Sea Rocket . • Sandy Shores Y& A. Purplish.

7 Man Rocket . • Sea Shores Yell.

8 Wall Rocket . • Lond. Walls Com. YA & C. Yell.

9 Garden Radish . • L. Gard & Fields Int. & C. Blush.

10 White Charlock . • L. Corn. Com. I& Y. Pale.

11 Great White Charlock . • L. Corn. Com. I& Y. Pale.

12 Flixweed . • * L. Old Walls Ruins Y& A. Yell.

TAB. XIMI. 1 Bitter Crefs . O. L. Wet Ditches I& Y. Wh.

2 Early water Crefs . O. L. Rivelets Iune . Wh.

4 Hairy Crefs . O. L. Rivelets . Com Yay. Wh. 2 Water Crefs. O. * L. Rivulets Com Y&A. Wh.
3 Early water Crefs. O. L. Rivulets Iune. Wh.
4 Hairy Crefs. O. L. River sides &c. Apr. May. Wh.
5 Lady fmock or Crefs. O. X. L. Wet Meads Apr. &c. Blush.
6 Double Lady Crefs. O. Rare. Blush.
7 Impatient Crefs. ... Rills &c. White.
8 Trefoild Crefs. O. Rare. Blush.
9 Daisy Crefs. O. Rocks & c. Rare. Wh.
10 Tower Mustard. O. L. Walls &c. I&Y Pala 9 Daisy Cress. O. Rocks & c. Rare Wh.
10 Tower Mustard. O. L. Walls & c. I&Y. Pale.
11 Great Tower Mustard. O. Rare Pale.
12 Daisy Tower Mustard. O. Sandy dry grd Wh.

TAB. XIVIII. 1 Thale Cress. O. L. Walls. Sandy grd Apr. & c. Wh.
2 Broad Thale Cress. O. Walls. Apr. & c. Wh.
3 Wieathed Cress. O. Walls & c. I&Y. Wh.
4 Small wreath Cress. Walls & c. I&Y. Wh.
5 Speedwell Cress. O. Hill sides Apr. & M. Wh.
6 Whitlow Cress. O. * I. Walls & c. Mar. & Apr. Wh.
7 Notched Whitlow Cress. Banks & Walls. Apr. Wh.
8 Irish Rush Cress. Loughs & Meers.
9 Woad. O. * Fields. Sowed. Iun. & c. Yell.
10 Pepper -wort. Dittander. * Water sides Y&A. Wh. 10 Pepper -wort. Dittander. * Water sides Ysch. Wh. o 11 Gold of Pleasure. • Flax fields. Y & c Yell. 12 Sea grape Colewort. O. Sandy Shores. Wh. n Gold of Pleasure. • Flax fields. Y&c. Yell.

12 Jea grape Colewort. • Sandy Shores. Wh.

TAB. XIX. 1 Garden Scurvy-grass. • Sand&c. Apr. Wh.

2 Jea Scurvy-grass. • Sea Shores. Wh.

4 Sheaperds - purse. • XI. Wast grd Com. Mar. to Nov. Wh.

5 Ragged Sheaperds-purse. • I. Apr. &c. A.M. & Iun. Wh.

6 Jmall Sheaperds-purse. • I. Apr. &c. Wh. Walls &c.

7 Whole Sheaperds-purse. • I. Banks. Walls Apr. &iWh.

8 Broad Water-Radish. • I. Slow Waters. I & Y. Yell.

9 Small jagged Water Radish. • I. Wet ponds I & Y. Yell.

10 Great jagged Water Radish. • I. Wet ponds I & Y. Yell.

11 Broad Horse Radish. • X. Gard & Wet Ditches. Wh.

12 Jagged Horse Radish. • X. Gard & Wet Ditches. Wh.

2 Jheaperds - Cress. • I. Sandy grd May &c. Wh.

3 Weish Rock Cress. • Mount. Rare.

4 Fine Cress. • Strincents Rock. Wh.

5 Valencia Cress. • Salfbury Plain. Pale.

6 Wart or Swines Cress. • I. Wet banks &c. I&Y. Wh.

7 Comon Cow-Cress. • XI. Corn &c. I&Y.

8 Smooth Cow-Cress. • XI. Corn &c. I&Y.

8 Smooth Cow-Cress. • Rare.

9 Penny Cress. • Males &c. Mount. Wh.

10 Hoary Cress. • Milly Pastures North Wh.

12 Small Thorow Cress. • Stone pitts &c. Wh.

MARKS & LETTERS Explained.

Signifies an ANNUAL Plant, with dies Yearly.

ABiennial Plant, with lives 2 Years.

APerennial Plant or Tree.

APlant of Noted VERTUE. *A Plant of Less.


H. Plants growing ab Hanfted Heath & c.

L. Plants growing within 10 Mile of London.

M&I. Plants Flowring in May & Tune.

I & Y. In Iune & Iuly. Y&A. In Iuly & August.

This, or a Latine Catalogue, and y Dry Samples of & Plants themselves may be had, if desired from y humble Servant





HERBARIJ BRITANNICI Clarifs. D. RAIJ Catalogus Cum ICONIBUS ad Vivum delineatis, & A.RE incisis; in FOLIO. Quibus adjunguntur Singulorum Nomina, summa cum Arte sculpta, simul cum Locis Indigenis, Tempore Florendi, Coloribus Florum, Notis seu Annuis, seu Perennibus, Vulgaribus, vel Raris, cum Floribus, Seminibus, eorumque Vasculis, ab ipsis Plantis Figuratis & singulis Speciebus annexis

A JACOBO PETIVER, Regiæ Societatis Socio &c. LONDIN.

SECT. I Herbæ BRITANICÆ Flore Stamineo RAY Hist.

Plant. 154. Syn. 28. Edit. 2. p. 52. MORIS. 577.

IAB. I. Fig. 1. Lupulus mas O.H. Duments. Sepibus Y.A.

2 Lupulus fæmina O.* Fruct. Sept. RAY Hist Plant 196. 1

3 Cannabis mas. Agris feritur. Y.A.

4 Cannabis fæm. Agris feritur. Y.A.

4 Cannabis fæm. Agris feritur. Y.A.

4 Cannabis fæm. Aug. RAY H.P. 198 pl. 1

5 Cynocrambe mas O.H. Sylvis Sepibus MAm.

7 Potamogiton pennis mille falis O.Fofsis ag.

7 Potamogiton pennis mille falis O.Fofsis ag. 7Potamogiton pennis mille folis O. Folsis aq. R. 1912.
8 Potamogiton pennat: verticill. O. L. Aq. pigris R. 1322.1.
9 Chrysosplenium vulg O. L. Humidis umbros. Apr. R. 207.
10 Chrysosplenium pedic. longis O. Apr. luts/is. R. 207. 5. Cynocrambe mus. o.H. Sylvis. Sepibus. M.Apr. 6. Cynocrambe fiem. o.H. Sem. Apr. RAY 163.1. 7. Mercurialis mas. O.L. Hortis Olitarys. I.Y. 8. Mercurialis fam. O. + L. Hortis I.Y. RAY 163. 2. 11 Chrysosplenium Cornubiense O. Raro. Udis &cY.A.R 211. g. Urtica major 0 * I. Sepib Ruder YA &c. Com. R. 160.4. 12 Hydrocotyle vuly O.H. Palustribus Y.A.R. 1343.1.

[14.
[14.
[14.] TAB.VII.1. Atriplex foldeltoide 6.L. Hort Timetis Y.A.R. 1923. 10. Urtica minor . L. Fimetis, Horns J.Y&A. R. 161.5. 11. Urtica Pilulifera . Raro. I&Y. R. 161.6. 2 Atriplex marit fol. deltoide . Udis fallis. Com.Y. A. R. 193.10. 12 Xanthium . L. Aquosis, letts &c .I.Y. R 165.1 3 Amplex marit laciniata . Arenos falis Y.A.R. 193. 8.

4 Amplex angustif dentata O.L. Udis falis & alijs. A SR 192,

5 Amplex Polygoni folio L. Timens & Com.Y.A.R. 192. 4.

6 Amplex Grammea marit Salis. TAB.II. Fig 1 Lapathum aquat may 0.* L.Rivis & c.V.A.R.177. 4.
2.Lapathum acutum crypum 0 L.Humid Hort YA.R.173.6.
3 Lapathum acut planum 0.* L.Inculus Aq. & c.Y.A.R.173.7. 3 Lapathum acut planum © ** L. Inculus. Aq. & c.Y.A. R. 173.7.

4 Lapath. acut. minimum. © L. Folsis. Aq. Y.A. R. 174.

5 Lapathum Sanguineum. © L. Hovt. Viis. Semit. Y.A. R. 174. 10.

6 Lapathum viride © L. Sepib Umbros. Y.A.

7 Lapathum aureum. © L. Aq. pigris. Y.A.S.

8 Lapathum aureum. © L. Aq. pigris. A. S. R. 174.8.

9 Lapathum obtusum. © L. + Areis. Humid. Y.A. Com. R. 173.5.

10 Lapathum. cipustum. © L. A. spris yarder. Y.A. R. 174.6. 7 Halimus O. Agger marit Y.A.R. 195. 3. 8 Halimus capsula. cordata. O. Marit. Raro. R. Syn. 63.7. gBlitum rubrum minus . Hortis . R 200.
10 Blitum poly/permon . L. Hort. Arvis Fimet. R 196. 2.
11 Blitum foly/mem . * L. Fimetis Ruderatis Y.A. R 198.11. 12 Spinachia sylvest. 0 * L. Areis, compitis &c.M.I R. 195. 1. 10 Lapathum sinuatum O.L. Agris, ruder. Y.A. R. 174.9 TAB VIII 1 Blitum Quercus falio . I. Ruderibus &c. A. S. 11Feg opyrum .+ . Agris sering Y.A . Fl. alb.R. 1821. 2. Blinum fol candicante . L. Fimetis Com. Y.A. R. 198. 8. 12Fegopyrum scandens O.L. Hort. Cultis. &c.Y.A. R. 182. 2. 3 Blitum Ficus folio . L. Fimetis Y.A. TAB III. 1. Acetosa vulg. 0. *. L. Pratis. & c. IY. R. 178. L. 2 Acetosa lanccolata 0. L. Arenos simis. M. I. R. 180. 14 4 Blinum fel subretundo • L. Ruderibus A.S.
5 Blinum fol lucido crasso • L. Finetis & c. A.S.R. Syn. 64.5.
6 Blinum acutore folio • L. Ruder Fimens A.S.R. 197. 9.
7 Blinum Aceris folio • L. Ruderibus A.S. 3 Acetosa lanceol.mas. seu Sterilis O.L. Ar en. sicc. A & M. 4Acetosa rotundif. Cambrix O. Clivis. aq. M.IY.R. Syn. 57. 10. gPersicaria acris O.*L. Folsisaq. A. J. R. 182. 6Persicaria minor O.L. Pratis aq. Y.A. R. 183. 3. 7Persicaria maculosa O.L. Aq. leget. A.S. Carn. R. 183. 2. 8Persicaria mac. incana O.L. Ray. Y.A. 8 Blitum lattore folio . I. Fimetis . A. S. 9 Beta maritima O Sallis palultr. A.S. R. 204.3. 10 Parietaria O XI. Muris Ruderatis Y.A.R. 206. 1 oPersicaria argentea .H.Seget.Y.
oPersicaria albida ramosa .L.Raro:
11Persicaria maj. pallida .L.Aren seget.Y.R.Syn.58.
12Persicaria amphibia .L. in & extra Aquas Y.R. 148.6.
TAB.IV. 1. Plantago latif. vulg. O.A.L.Vijs ubiqe.Y.A.S.R. 876.1 11 Vermicularis arborescens O Maritimis. R. 199. 17. 12 Vermicularis frutescens O. Littor. Sallis R. 199.16. TABN 1. Kali minus album Sallis marit. Com. • R. 198.14. 2. Kali Spinosum Avenolis marit. . R. 212. 2. 3 Kali geniculatum Salûs marit. Com. •.
4 Kali geniculat lignosum O. Maritimis R. 211. 1.
5 Anonymus Lini folio O. Mont. Cretaceis Y.A. R. 399. c.7.
6 Knawel erectum • H. Arenos. sterilibus Y. 4B.W. 1. Plantago latif. vulg. 0.4L. Vijs ubiqe.Y.A.S.R. 876.1.

2Plantago dentata O.L. Advias. & c.Y.A.

3Plantago incana O.L. Glaveo fis. vijs. M.I.Y.R. 877. 5.

4Plantago Rosea & spicata O.Raro'.R. 876.

5Plantago panicula sparsa' O.Raro'.R. 877. 4.

6Plantago angustif. major. O. **L. Pascuis ubique. M.I.Y.R. 877.7.

7Plantago angustif. cambria O.Rupibus. R. Syn. 185.7.

8Plantago angustif. prolifera O.Raro'.R. 877.1.

9Plantago marina O.L. Maritimis. R. 880.5.

10Plantago Coronovi fol. O.+H. Aren & Marit. R. 870.2. 7 Knawel supinum & H. Arvis arenosis YR. 213.1. 8 Gramen Fiperinum L. Puteis Y.A. R. 13.25. c. 10. 9 Alchimilla vulg O. ** Pratis Pakuis mont R. 208.1. 10 Alchimilla minor O.Raro . R. 202.2. 11 Alchimilla argentia O Mont Rupibus R. 209. 4.
12 Percepier Anglorum * I. Solo restibili steril • R. 209. c.14. 10Plantago Coronopi fol. O. + H. Aven & Marit. R. 879.3. 11 Sanguisorba major 0. +. Prat. humidis R 40 2. 2.
12 Sanguisorba minor 0. *. Cret. prat. & Mont. Y. R. 401. 1.
TAB.V. 1. Bistorta maj. vulg. 0. * L. Pratis humid. Y. R. 186. 1.
2Bistorta minor 0. Pratis mont. R 187. 4. TAB.X.1. Polygonum latiful. • * L. Yijs. Semitis &cYAR. 184. 1.
2. Polygonum fol roundo • L. Incultis Y.
3. Polygonum fol brevi • L. Glareolis Y. 4Polygonum fol. angus O.L. Seget. steril. 5Polygonum marimum O. Aren saliis. R. 185. 4 2Bistorta minor. O. Pratis mont. R187.4.

3Bistorta min. fol. vario. O. Clivis Rupium. R.187.6.

4Potamogiton latifoil O. L. Stagnis. Pissinis. A. S. R188.1.

5Potamogiton fol. longo acuto O. L. Fluvijs. Y. R188.2.

6Potamogiton perfoliatum O. L. Fluvijs. Y. R188.3.

7Potamogiton fol. cordato O. L. Fluvijs. Y. R189.3.

9Potamogiton planum O. L. Rivuijs. Y. R189.4.

9Potamogiton crivpum O. L. Rivuijs. Y. R189.5.

10Potamogiton Gram. latum O. Aq. pagris.

11Potamogiton Gram. angust. O. Rivis.

12Potamogiton tenuifol. O. Rivis. R189.7.

13Potamogiton Laniculas cum O. Rivis. 6 Polygonum fol. circinato. Maritimis . Raro 7 Polygonum *verticillatum* Pasciris Udis Paliutr. R. 214.3. 8 Polygonum *Valentinum* Maritimis R. 215.8. 9 Herniaria glabra Promont marit R. 214.6. 20 Herniaria hirsuta O Arenosia. 11 Erica maritima. Avenos & sallis marit Y.A.R.1716.15. 12 Portulaca aq. tricoccos . H. Seget Udis Palustr. R. 1035.26. H. Plantæ circa Hamsted crescentes. L. Pl. intra X Milliaria LONDIN nascentes.

M.I.Y.A . & c. Pl. Mensib. May. Lunig. Iul. & Flor.

R. Pagina in qua Hist. RAY Planta describitur.

Notæ & Liter æ Explicatie.
• Planta Annua. • Planta Biennis.

O. Planta Perennis. * . Planta Officinalis . + . Pl. minor. Virium .

```
SECT. II. PLANTA Lactes centes & Pappe scentes. TAB XVII. Fig. 1 Baccaris Montpel O+L Mont siccis. Vijs YA R292
TAB XI. Fig. 1. Pilos ella officin. O *I. Pascuis siccis. IY. lut RAY 2 Eupatorium Cannabinum O+L. Ripis rivul stagn YA carn R293

242-1. a Carabelliam Annue con vig. O Barb V A. alb R. 284 15
                                                                                         3 Gnaphalium Americanum O Baro Y.A. alb. R. 284.15.
  2 Pilosella Cambria RAY 241. 19.
                                                                                          4 Gnaphalium Montanum O Ericetis. mont. YA.R. 283,12
  3 Pilosella folio longo L.Fl. luteo.
  4 Pilosella fol lato. Aggeribus folsar um. lut. R. 244.2.0.
                                                                                          5 Gnaphalium Terseianum OM ur is agger. albid. R. 284 17
  5 Pilosella fol longo glabro. O. Mont. summit. 1ut. R. 245. 8. 6 Pilosella Plukenety. O. I.Y.A. 1ut. R. Syn. 75. 11.
                                                                                          6 Gnaphalium Anglicum @ H. Pascuis arenos. YAR. 295.13.
                                                                                            Gnaphalium fol longo .+ L. Stagnis siccis &c. Com.Y.A.R.
  Dens Leonis vulg. O. * L. Pas cuis ubique Apr. MIY &c lut R 214.8 Gnaphalium Germanicum Avenosis &c raro.
                                                                                         o Gnaphalium proliferum . + L. Vijs siccis, steril Com Y.A.
  8 Dens Leonis angustifolius. O.L. Muris Siccis MIY. lut R 5.76. 2.
  gDens Leonis hirsutus O.L. Pascuis &c. Com. M.I.Y. lut R 245.5. to Gnaphalium prolifer minus . Sterilibus siccis YA.
  10 Dens Leonis hirsut.minus. Aggeribus I.Y. lut. R. 246.
                                                                                         11 Cnaphalium minimum . H. Seget aren siccis YAR. 196.6.
  11 Dens Leonis hirsut. ramosus O.L. Pascuis. Com. IYA. lut R 230.2.12 Gnaphalium Gramineum . Seget arenos Y.A. R.S. 85.5.
  12Dens Leon. ramos. glaber. Pascuis. lut.
TABXII. Fig 1. Hieracium præmorsum vulg OL, Pascius ComYA lut, SECT. III. HERBA Corymbifera, Nuda Capitata
                                                                                            & Nodifloræ
  2 Hieraclum pramors laciniatum. O.L. Pascuis Y.A. lut.
  3 Hieracium fol acuto minus L. Sepibus YA. lut.
                                                                                        TABXIX. Fig. 1. Bellis major O.+ L. Pratis reget IY. albR350
                                                                                         2. Bellis minor O. * I. Pratis pascuis Com. T. state tota alb R349
  4 Hieracium fol. obtuso minus . I. Sepibus Y.A. lut.
  Hieracium Lactuca folio. L. Pascuis &c.Y.A. lut. 6 Hieracium Aphacoides. L. Pratis &c. Y.A. lut. R. 234.16.
                                                                                         3Ptarmica O + H. Pratis udis Y.A. alb R.344.1.
4Millefolium yulg. O. *I. Pascuis. Com.I.Y.A. alb s purp R.345
                                                                                         5 Matricaria O. * L. Sepibus, rudetis. Y.A. alb. R. 357.1.
6 Chrysanthemum segetum. L. Seget Y.A. lut. R. 339.3.
  7 Hieracium Aphacoid acutum . L. Pascuis Y.A. lut.
  8 Hieracium Caftor odore O.L. Pascuis &c Y.A. htt. R. 232. 6.
  9 Hieracium Cichor, fol.minus Q.L. Pascuis YA hut.
                                                                                          7 Buphthalmum luteum 0 + Ripis aq raro YA Jut R.341.9.
  10 Hieracium Cichor fol majus @ Parcus &c. I.A. lut R 2,36 5
                                                                                          8 Buphthalmum album L. Pratis & c.M. I. alb.
  11 Hieracium asperum OL Agror marginibus Y.A. lut. R. 233 12
                                                                                         9 Chamemehum amarum . O.+. L. Cultis, Seget . alb . R. 355. 2.
   12Hieracium Echioides 🖭 Sepibus Agris &c. V.A. lut R. 2,33.10. 10 Chamæmelum odoratum 🔾 🖈 L. Pascuis udis Y.A.S. alb R333
TABXIII. Fig. 1. Pulmonaria maj. latifol. O. Agger folsar Y. lut R 23,97.11 Cotula fætida 0.+ L. Seg et cultis. alb. R. 355.3.

2. Pulmonaria fol. rotundo O. Rarò. I.Y. lut. R. S. 74.6.


12. Cotula inodora 0.+ L. Agris. cultis. alb. R. 359.
                                                                                          12 Cotula modora • + L. Agris . culus . alb R.355.
  3Pulmonaria fol lato dentato ORarò I.Y. lut. R. S. 74
                                                                                        TAB XX Fig 1 Absinthium vulg 0 * L. Incult vijs ComAS.
  4Pulmonaria Gallica O.H. Muris. Sylvis &c.Y.A. lut.R. 239. 9. 5Pulmonaria Gallica ang ultif. O.Rard. Y.A. lut. R. 240. 11.
                                                                                         2. Absinth marin Anglicum 0* Maritimus. Y.A.S.R. 370.16.
                                                                                         3 Absinth marin Gallicum O + Maritimis YA & R. 370. 15.
  6 Pulmonaria latif. villosa O Rupibus altis Y. lut. R. 239. 8.
                                                                                         4Abrotanum campestre O Steril vijs rard Y.A.R. 370.4.
  7 Pulmonaria frut latif hirsuta O H. Sylvis Y.A. lut. R. 238. 28 Pulmonaria frut longif hirsuta OH. Sylvis & Y.A. lut.
                                                                                         5Arthemisia O. * L. Sepes Agror Marg. IYA R. 37 2.1.
                                                                                         6 Bidens fol. integro • + I. Stagnis aq A. S. lut R.3 61. 2.
  9Pulmonaria frut longif. glabra O Raro Y.A. lut. R. 238. 1.
                                                                                          7 Bidens fol vifido + L. Stagnis aq. A. S. lut. R. 360.1.
  10 Pulmonar ia angustif hirsuta O.H. Sylvis Y.A. lut. R. 23 8.5.
11 Pulmonar ia augustif glabra O.H. Sylvis Y.A. lut.
12 Pulmonar ia Granduca O Raro Y.A. lut.
                                                                                         8 Cotonaria marina O Aren marit raro lut.R. 2941.
                                                                                         g Tanacehim vulg. O. * L. Agr. marg Vijs. A. S. lut. R. 3 65.1.
                                                                                         10 Tanacetum crispum O.+. Hortis &c. A. S. lut R. 365. 2.
TAB XIV Fig 1 Souchus rotund fol. 6. L. Seqetilus. TY: lut.
                                                                                         11Agrimonia O.*.L. Pascuis, Sepib. &c. Y.A. lut R. 400.1.
  2 Sonchus asper latifol. O.L. Hort. Fimetis . IY. lut. R. 225.7.
                                                                                         12 Circe a major O.L. Sylv Sepib umbrosis Y.A. alb R401.
                                                                                        TABXXI Fig.1. Carduus Moschatus @ L. Agris saxos YA R308
   3 Sonchus angustisolius . O. Raro'. I.Y. lut.
  4 Sonchus asper dentatus. O.L. Arvis incult. Y.A. lut.
                                                                                         2. Carduus Crispus D.L. Sepes, Vepres V.A.R. 309. 2.
  & Sonchus asper laciniatus. L. Ruderatis &c.Y.A. lut R 2 25. 6. 3 Carduus Acanthoides @L. Agger ruder I.Y.R. 309. 3. 6 Sonchus segetabis major. O. L. Agger. Cultis. &c.Y.A. lut R 226x4 Carduus palustr . H. Udis. palust. A.S.R. 309. 4.
  7 Sonchus altissimus.O.L. Ripis Fluvior. Y.A. lut. R 226.11. 5 Carduus vulgatiss O.L. Vijs arvis &c. Com. Y.A.R. 310.8.
  8 Sonchus lavis latifol. C.L. Hortis. Cultis & : Y.A. lut R. 244. 2. 6 Carduus acaulis min. O.L. Mont. sicc. Y.A. R. 310. 6.
  9 Sone hus lavis laciniatus. . * L. Horris &c IY.A. lut.R. 224.1.
                                                                                         7 Carduus lanceatus @ L. Sepib incult. Com YA R. 310.1.
  10 Sonchus lavis laceratus. L. Hortis &c. I.Y. A. lut. 8 Card Eriocephalus Camp & mont. rare Y.A.R. 311.3.

11 Cichoreum sylvestre. . L. Semitis incultis Y.A. carR 155.9 Carduus Maria X.L. Agger marg. agror Y.A.R. 313.1.

12 Lampsana. . L. Hortis, cultis. I.Y. A. lut. R. 15 6. c. 2. 10 Carduus tomentos. . L. Vijs incult. Com Y.A. R. 313.1.
TAB.XV. Fig. 1. Lattica sylv. latifol. O.L. Hort Sepibus Y.A. lut. 11 Cardius stellat. O.+ L. Vijs publicis A. S.R. 317. 1. 2 Lactica Endivia fol. O.L. Aggersaxosis I.Y.A. lut R. 221. 2. 12 Cardius Sol/litialis. O. Sepes rard. A. lut. R. 317.
   3Lactica lacerata. . +L. Saxolis lepibus Y.A. lut. R 222.3.
                                                                                       TAB XXII. Fig. 1. Cirsium palustre. OL. Pratis udis YA R306.
   4Lactuca lacerata. minor. L. Agger. folsis. IY.A. lut.R. 222.5. 2 Cirsium montanum O Mont. Eborac &c.Y.A.R. 306.8.
5Lactuca Hedera. fol. H. Muris agger. umbros. Y.A. lut.R. 223.7. 3 Carthamus Offic. * Agris seritur. Y.A. croc.R. 302.1.
                                                                                         4 Cyanus Segetum • K.L. Seget YA. carul R. 321.1.
6 Serratula fol integro. O.H. Sylvis YA. purp.
6 Serratula vulg. O. Sylvis prat. YA.R. 331.c.16.
7 Iacea major vulg. O.L. Seget I.Y. purp. R. 325.11.
8 Iacea nigra vulg. O+L. Pascuis Com. IYA. R. 325.12.
   6Tragopogon angustis: O.H. Pratis. IY. lut. R. 252. 6.
7Tragopogon vuly. O.H. Pratis. Pascuis. IY. lut. R. 252. 1.
   oTragopogon purp. Raro Hortis, I.Y. Purp R. 252.3.9 Hyoseris segetum. L. Arenosis. Seget. IY. lut. R. 229.2
   w Carlina sylv. O.L.Ericetis Palcuis, liccis Y.A. lut.R. 288. 2.
                                                                                         9 Iacea nigra lacin. O+L. Pascuis. I.Y. purp.
10 Scabiosa Offic. O*L. Seget. pascuis. I.Y. Carn. R. 374. 2.
   11 & 12. Petasites vulg. 0. *. L. Ripis. humidis. M. Apr. carn. R. 260.
 TAB.XVI. Fig. 1. Helenium. 0.*. Pratis humidis. I.Y. lut. R. 273. C. 7.
   2 Conyza vuly. O. +. L. Folsis aq. Y.A. lut. R. 262.5.
                                                                                         11 Scabiosa minor O Pascuis mont sicc YAR 3743.
   3Conyza minima L.Stagnis siccis. Y.A. lut R. 262.6. 12 Succisa Offic. O. * H. Prat. pascuis A. S., car R. 38 o. 21. 4Conyza carulea O. L. Pascuis sicc. Steril. Y.A. purp. dil R2701 TAB XXIII. Fig. 1. Bardana vulg. O. * L. Vijs &c. Com I'A R3321.
   5 Conyza palustris. Folsis aq Palustr. I.Y. lut.R. 363.10.6 Conyza palust. lacerata. Folsis aq. &c. I.Y. lut.R. 264.12.
                                                                                         2 Bardana Rosea OL Incultis YA R.332
                                                                                         3Bardana capite minore @L. Ruderatis YA.
   7Virga aurea latif. serrata. O. Ripis. Rarò! Y.A. lut.R. 279.6.

4 Bardana cap minore incano O. L. Vijs. ruder . Y.A. 8 Virga aurea angustif. serrata. O. Folsis, palustr. raro! lut. R. 5 Bardana cap majore incano O. L. Incultis Y.A. 9 Virga aurea major. O. Sylvis, Ericet. rarò! lut.

(38011. 6 Bardana cap araneoso. O. Montanis & c. Y.A. R. 332.2.
   noVirga aurea vuly. 0.*H. Ericet Sylvis &c. Y.A. lut.R. 278.1.
                                                                                          7Dipsacus sativvus @+Agris seritur I.Y.R. 382.1.
   nVirga aurea Cambria. O. Pascuis mont. lut. R.S. 813.
                                                                                          & Dips acus Sylv. 8+L. Humid incultis YA. R.382. 2.
 12Virga aurea alba acris. . Muris Hortis. A.S. albid. R. 2714.
TAB. XVII. Fig. 1. Jacobea vulg. 0.* L. Vijs incultis. &c. Com Y.A. lut.
                                                                                         9Dipsacus minor @L. Rivulis . aq YA.R. 382.4
                                                                                         10 Eryngrum marinum 0 * L. marit. aven YAR 384.3.
   2 Jacobea palustris O.L. Humidis, udis. &c.Y.A. lut. R. 285.4
                                                                                         11 Eryngrum campeftre O Rard A. S. pall R. 384.1
   Jacobea Senecionis fol. O.L. Vepretis Sepib. Y.A. lut R.285.2.
                                                                                         12 Sanicula Offic. O. * L. Sylvis, Sepibus M. I. alb R 475.04.
   4 Jacobea Cantubrigiens is O. Mont. sheril. & cretae. lut.R. 27.2.4. 5 Senecio vulg. O.*L. Cultis arvis. Areis &c. Com.R. 290.1. 6 Senecio mcanus. O. H. Arenos. agger. &c. I.Y.A. lut.R. 290.2. 7.8. Tussilago O.*L. Aq. & humidis. M. Apr. lut.R. 25.9.7. 9 After marit. flavus. O. Rupibus marit. lut.R. 26.8. 11.
                                                                                           SECT . W. HERBÆ Umbelliferæ Ray 387.
                                                                                       TAB. XXIV. 1. Sphondylium vulg. 6.+.L. Pascuis humid. KA. 2. Sphondylium laciniatum. Pascuis Y.A. alb. R. 408. 2.
                                                                                         3 Pastine a sativa @ + Hort. & Agris seritur Y.A. alb.R.410.2.
                                                                                         wTripolium majus.o.+L.Sallis marit.Y.A. purp.R. 270. 20.
    11 Tripolium minus. O. Sallis, marit. Y.A. purp R. 270. 10.
                                                                                         5 Tordylium majus Raro Y.A. alb.R.411.
6 Tordylium minus Raro Y.A. albic R.412.2.
   7. Tripolium nudum. O. Aq. Sallis. Y. A. hut. R. S. 80. 2.
```



7 Peucedanum vulg. O.+ Pahultr. marit A. flavesc R. 416.1. 9 Asperula O. +. H. Sylvis. R. 483. 1. 10 Rubeola repens carul. . L. Arvis. R. 483.3. 8 Crithmum marinum O.+. Clivis marit A. S. flaves c.R. 457.2. 9 Angelica sativa • * Hortis & c.Y.A. alb.R. 434.1. 10 Angelica aq. 9. H. Ad. rivul. & aq.Y.A. alb. R. 434.2. uAparine Offic. . * L. Sepib. Com. R. 484.1. 12Rubia Cynanchica. O. L. Collib. cretaceis. R.485. c. 9. 11 Coriandrum Offic. 0.*. Agris scritur . R. 470. SECT.VII. Planta Verticillata.RAY. 508. 12 Hipposelinum 0*. Rup marit. I.Y. albesc. R. 437. 1. TAB.XXV. Fig. 1. Scandix vulg. . I. Seget. M. J. alb. R. 428.1. TAB. XXI. 1. Serpylliun vulg. O. *.L. Palc. sicc. R. 521. 4. 2 Cicutaria vulg. O.L. Pomarijs Sepibus. Com A.M. alb.R. 429.1. 2 Serpyllum fl. amplo O.Mont. R. S. 1.2.2. 2. 3 Cerefolium sylv. . L. Sepibus Y.A. alb R. 431.6. 3 Serpyllum latifol. O.R. 521. 2 4 Serpyllum Cibratum . O. Mont. R. 522. 2. 4 Meum offic 0* Pratis Boreal Y.A. alb. R. 432.1. 50 enanthe aq. major.O.L. Pratis udis R. 441.1. 5 Mentha aq. verticillata O.L. Udis .R. 530. 10. 60enanthe aq. minor. OL. Pratis udis R441.2. 6 Mentha aq. vulg. O.L. Rivis. R.533. 12. 7Mentha fufca. O. Hort. &c. R.530. 2. 7 Oenanthe Cicutte facie. O.L. Rivulis R. 441.4. 8 Oenanthe Staphy lini folio. O. Rard. R. 442. 6. 8Mentha crispa.O.Rivis.R.531. 3. gMentha Piper ata lata R. 532. 7. 20Mentha Piper ata acuta R. S. 124. 5. 9 Bulbocastanum O.+.H. Pascuis aren Y.A. alb. R. 440.1. 10 Podag raria 0.+ L. Sepib Hort. alb R. 435. 4. 11 Amomum Offic 0.* I, Folsis lutosis . Y.A. alb R. 443.3. uMentastrum longifol.⊙.Riguis.R.532.9. 12 Selinum segetale 9.+L. Humid. Seget. alb. R. 443. 4 22 Mentastrum rotundifol. O. Udis. R. 532.7 TAB.XXVI. Fig. 1. Sium max. latifol O. L. Fluvijs Y.A. alb R. 443 TAB.XXXII. 1. Nepeta Offic. O. * L. Sepes . R. 548. 1. 2 Sium angustifol O. I. Rivulis YA. alb R 444. 4. 3 Sium repens Folsis aq. I.Y. alb R 444. 5. 4 Sium minimum O. L. Stagnis aq. M. I. alb R. 444. 6. 2 Pulegium Offic. O.* L. Palustr. R. 533. 1. 3 Prassium Offic. O. * L. Vijs, ruder. R. 556. 1. 4Ballote Dioscor. O.+L. Sepes. ruder .R. 571. 1. 5 Marubium 44. O.L. Rivis. R.535. 1. 6 Betonica Offic. O.*.H. Sylvis. R.550.1. Saxifraga major. O Sylvis. Y.A. alb R. 445. 1. 6 Saxifraga maj. laciniata. O Sylvis YA alb. 7 Galeopsis vera. O.L.R. 548. 8 Horminum sylv. O. +.L. Glareosis. R. 545. 12. 7 Saxisraga minor OH. Pascuis siccis Y.A. alb. R. 445. 3. 8 Saxifrag a minor laciniata O.H. Pascuis sicc. R. 445.4. 9 Saxifraga Cornubiæ O. Raro. g Clinopodium majus. O.L. Vepret. R. 558.17. 10 Carui Offic. 0.*. Pratis palustr. I.Y. carn. R. 446.1 10 Clinopodium minis. O.L. Mont. cret. R. 553.1. nPrunella Qffic.⊙.*L.Palcuis .R. 5511 11 Apium Scoticum O. Saxis marit. alb. R 447.1 12 Apium Offic. 9. * I. Udis , folsis marit A.S. alb. R. 447. 2. 12 Stachys vuly. Q. Raro. R. 554 1. TAB. XXVII. 1. Toeniculum vulg. 0. * I. Clivis marit R. 457. 1. TAB. XXXII. 1. Heder a terrestris. 0. * I. Sepes. R. 567. 1. 2Lamium rubrum. • * L.R.559.3. Com. 2Peucedanum minus. Rupibus Rard. alb R 460.5. 3 Apium petræumo Montosis rard alb R. 463.5. 4 Carota sativa S. X. Agris serituralb R. 465.1. 3Lamium rubrum fol incisis. O.L. Requiens R.560.6. 4Lamium perfol.majus. O.L. Hort Arvis. R. 560. 10. Daucus sylv. * X. Pas cuis. YA. alb. R. 465.2. 5 Lamium album. 0 * L. Sepib. dumet. R.559.1. Com. 6 Echinophora latifolia . Seget purp. R. 468.5. 6 Lamium luteum. O.H. Sylvis. umbros. R.560. 11. 7 E chinophora laciniata . Seget. raro. purp. 7 Melifsophyllon O. Sylvis raro. R.561. 13. 8 Echinophora tenuifolia . L. Seget . R. 467.3. 8 Lamium Cannab. vulg. O.L. Arvis. R.561.13. gAnthriscus vulg. . H. Dumet. Sepes Y.A. rub R 468.6. 9 Sideritis aq. O. + L. Thrvijs. R. 563. 3. 10 Anthriscus minor OI. Seges V.A. alb.R. 468.7. 13 Sideritis lutea. . Seget raro. R. 566.15. 11 Caucalis nodosa • L. Agger. agror. R. 468. 12 Caucalis tenuifolia • L. Ruder. agger. AM.R. 469.9. 11 Sideritis angustifol. . L. Seger. R. 566.16. 23 Sideritis fol. subrotundo. . L. Seget. R. 505. 12. TABXXVIII. Fig. 1. Cicuta aq. O Rivul. stagnis R. 450.1. TAB. XXXIV. 1. Calamintha Offic. O. * L. Vijs. Folsis. R. 569.1. 2 Cicuta vulg 🖲 * L. Agger. &c. Com.Y.A. alb. R. 451. 2. 2 Calamintha odore Pulegij. O.+L.Vijs. agger. R. 569. 2. 3 Cicuta minor OL. Hort. arvis YA. alb.R. 451.3. 3Bugula vulg. O. *. L. Sylvis, Pratis udis. R. 575. 1. 4 Cicutaria aq. O.L. Aq. comosis. alb.Y.A.R. 452. 6. 4 Bugula . Alpina O. Montosis . Ravo. R. 575. 2. 5 Seseli pratense O.* H. Pratis udis Y.A. lut. R. 453. 1.
6 Crithmum marinum O. Clivis marit. R. 469. 1. 5 Cardiaca 0*. Fimetis. R.571.1. 6 Scordium Offic. O.*. Palustribus. R.576. 1. 7Bupleurum minimum Pahuftr R. 474.4. 7 Scorodonia O.+ H. Dumens. Lylvis. R. 576.c. 29. 8 Perfoliata vulg. . L. Seget . R. 471. 8 Origanum vulg. 0.* L. Sepes. Dumet. R. 539. 4. 9 Chamæpytis vulg. ●* Agris.restil. R.573.1. SECT.V. HERBÆ Asperifoliæ RAY 493. 10 Cassida vuly. O.L. Fossis.aq. R.572.1. gBorrago Hort. ●.*.Hort. Muris R. 493. u Cassida minor. O.H. Paliultribus . R. 572. 2. 10 Buglosum sylv minus. I. Arvis R. 494.3. 12 Uerbena. vuly. 0.* Vijs. public. R.535-1. 11 Cynoglossum vulg. 8. * L. Ruder. R. 489.1

9Borrago Hort. •*Hort Muris R. 493.

10 Buglosum sylv minus • I. Arvis R. 494.3.

11 Cynoglosum vulg. • *L. Ruder R. 489.1.

12 Cynoglosum fol virenti • I. Vijs publ R. 491.8.

TAB XXIX. 1. Echium Vulg. • I. Vijs, Arvis.

2 Echium fl minore • Muris R. 5.119.

3 Buglosum marinum • Maritimis R. 500.13.

4 Pulmonaria fol. Echij • Sylvis Raro R. 485.3.

5 Symphytum majus. • *L. Rivis iossis aq. R. 505.1.

6 Lithospermum purp. • Raro R. 503.2.

7 Lithospermum vulg. • *L. Dumetis. R. 503.1.

8 Lithospermum arvense • L. Seg et. R. 503.5.

9 Echium Scorpioides aq. • L. Fossis. aq. R. 504.3.

10 Echium Scorpioides avens. • L. Arvis. Hortis. R. 504.1.

11 Echium Scorp: minus. • L. Ripis sicc. &c. R. 504.2.

12 Asperugo • Raro. R. 502.c8.

SECT.W. HERRÆ Stellatæ RAY 477.

TAB XX. 1. Cruciata hirsuta. O.H. Dumet. R. 479. 1.

2Rubia Sativa O. *. Hort. & Agris. Scritur. R. 408.1.

3Rubia syb. O. Rupibus. R. 408. 2.

4 Mollugo vulg. O.L. Dumet. Sepib. R. 481.1.

5 Gallium album. O.L. Udis. R. 481. 3.

6 Gallium alb. minus. Mont. & Udis. R. 482. 4.

7 Rubia crecta 4 drifolia. Mont. Raio. R. 478. 1.

8 Gallium luteum O. * L. Pascuis sicc. R. 482.1.

SECT. VIII. HERBÆ. Nasifloræ. RAY. 751. TAB. XXXV. 1. Anturhinum nunumum. 1. Culus. R. 760.37. 2 Antirrhinum medium. Arvis. R. 760. 31. 3Elatine fol. anguloso. . +. L. Seget. R. 759. 35. 4Elatine fol. rotundo. •* L. Seget. R. 759.34. 5Linaria vulg 0 x L. Pasc steril R. 752.1. 6 Linaria odorata O Sepibus Raro R. 756.20. 7 Pinguicula O. Vliginosis. R. 751.1. & Digitalis purp. $0 \times L$. Arenolis R.767.1. 9 Scrophularia vulg 0.*L. Dumet. Sepib R.764.3. 10 Scrophularia aq: 0. x. L. Folsis aq: R. 764.1. n Scrophularia Scorodonia fol. 0. Ripis aq: R. 766.x1. 12 Hyoscyamus vulg. 0 * L.Ruder. R. 711. 1 TAB. XXXVI. 1. Balsamina lucea . Ripis aq: raro R. 1328. C.S. 2Pedicularis luta . L. Pascuis R 760.1. 3 Pedicularis rubra elator . L. Aquofis R 770. 6 4 Pedicularis rubra vulg OH Ericetis udis R 770.5. 5 Euphrasia Offic •* L. Pascuis R 771.1. 6 Euphrasia major lutea Vliginos raro R. 772. 4. 7 Euphrasia pratens rubra • H. Prat. Seget R 772.3. 8 Euphrasia Ortoners is Udis raro.R.773.5. 9 Melampyrum hiteum . H. Dumet R. 775.4. 10 Melampyrum cristatum . Sylvis R 775.3.



```
. 11 Lentibularia major O Toisis aq. R. 1332. 2.
   11 Lentibularia major O Tossis aq. R. 1332. 2.

12 Lentibularia minor O Tossis aq. R. 3. 279.3.

TABXXXVII. 1. Viola purp. O * L. Hort. R. 1049.1. M. Apr.

2 Viola fl. pleno O Hort. Sylvis raro.

3 Viola alba O. Sepes agg cr. R. 1050. 2.

4 Viola sylv. O. H. Dumet. Sepes R. 1050. 3.

5 Viola aq rotundi fol. O. Udis Muscosis R. 1050. 4.

6 Viola rot. lutea O. Cambria.

7 Viola hirsuta major O. Dumet. R. 1051. 5.

8 Viola tricolor O. + Hort. Seget. R. 1052. 2.

9 Viola bicolor. O. L. Seget. R. 1053. 7.

10 Viola lacin. lutea O. Mont. udis R. 1052. 5.

11 Reseda vuly. Arvis Cret. R. 1053. 2.

12 Luteola O. L. Muris ruder. R. 1054. 6.

HERBE Sein echinato & stellato RAY. 578. SEC.
      12 Inteola © I. Muris ruder. R. 1053. 2.

HERBE Sein echinato & stellato RAY. 578. SECTIX.

HERBE Sein echinato & stellato RAY. 578. SECTIX.

TAB.XXXVIII. 1. Chelidonium minus O.* I. Udis R. 579.

2. Ranunculus dulcis O. H. Dumet. R. 584. 1.

3. Ranunculus prats crectus O. I. Pratis. Com. R. 383. 7.

4. Ranunculus prats crectus O. I. Pratis. Com. R. 383. 7.

4. Ranunculus erect. pallidus • Udis R. 382. 4.

6. Ranunculus repens vulg. O. I. Pratis udis R. 581. 1.

6. Ranunculus repens major. O. I. Fluv. ripis.

9. Ranunculus repens major. O. I. Fluv. ripis.

9. Ranunculus paluft. lavis. • I. Aquis. R. 585. 1.

10. Sinapi album. • I. Agger. folsar: R. 802. 2. lut.

10. Sinapi album. • I. Agger. folsar: R. 802. 2. lut.

11. Sinapi sativum. • *I. Hort: arvis R.

12. Sinapi arvorum. • I. Seget. R. 802. 1.

13. TAB.XXXX. 1. Ranunculus alb. Hedderaceus O Vadosis
      TAB.XXXIX. 1. Ramunculus aq. fol. varijs O.L. Stagnis R. 586.3.

2. Ramunculus aq. tenuifol. O.L. Stagnis R. 586.4.

3. Ramunculus aq. fol. Famiculi O.L. Aq camosis R. S. 136.5.

4. Ramunculus famicus major O.L. Vadosis R. 586.6.

5. Ramunculus famicus minor O.L. Uligines. R. 587.7.

7. Carda Muris D.L. Arvis udis R. 1222.6.5.
            7. Cauda Muris • I. Arvis udis R. 1332.c. g. &Flos Adonis • Seget. R. 596.1.

9. Malva Offic O. * I. Sepes. ruder R. 599.1.

10. Malva minor. OI. Semit. ruder R. 599.2.
        11Althea Offic: 0 * Maritimis R. 602. 1.
12 Area vulg O L. Uepret. Sepes R. 604. 1.
TABXL. 1. Caryophyllata vulg. 0. * L. Dumet. R. 606. 1.
              2. Caryophyllata fl. majore O. Sylvis R. 606.2.
3. Caryophyllata mont. pwp. O. Montosis R. 607.3.
4. Caryophyllata mont. prolifera O. Raro R. 607.4.
              5. Caryophyllata peninphyllea O. Scotia. R. 608.10.
6. Caryophyllata fol Chamædryos O. Montosis. R. 608.11.
7. Fragraria vuly O. H. Dumetis. R. 609.1.
             8Fr agraria stevilis O.L. Steril: pascuis R. 611.2.

9Ranunculus nem. albus O.H. Udis dumet R. 624.1.

10Puls atilla A. clauso O.Ericetis R. 634.2.
              11. Viorna. O.L. Sepibus R. 620.1.
       11. Viorna O.L. Sepibus R. 620.1.

12. Viorna fol. integro O.L. Sepibus.

TABXLI. 1. Pentaphyllum rubr. palustre O.R. 611.1.

2. Pentaphyllum villosum Palyber. S. 141. 2.

3. Pentaphyllum viulgare O.H. Semitus R. 611.2.

4. Pentaphyllum minus hivutum O.P. ascuis Ravo R. S. 142.4.

5. Pentaphyllum argenteum. O.L. Pasc. arenos R. 613. 14.

6. Pentaphylloides fragiferum O. Montos. Ravo R. 616.1.

7. Pentaphylloides Scoticum. O. Sylvis ravo R. S. 142.4.

8. Pentaphylloides E boracense O. Fluv: ripis R. 616.5.

9. Tormentilla vulg. O.* H. Dumet. & C.R. 617.1.
             9.Tormentilla vulg. 0 * H.Dumet & c.R. 617.1.
10.Tormentilla reptans 0 Arvis Sylvis R. 618.3.
              11, Argentina O. +. L. Udis . R. 617. c. 16.
        12.Moscatellina O.L. Umbrosis. 684.c. 27.

TAB.XIII. 1. Telephium vulg O.* I. Umbros. R. 689.1.

2.Rhodia radix O.Montos. R. 690.7.

3. Cotyledon vulg. O.* . Saxis & c. R. 1878.1.

4. Sedum majus O.* . I. Tectis R. 6872.
            6 Sedum luteum minus O.L. Tectis R. 691. 2.
6 Sedum lut scorpoidas O.L. Tectis R. 691. 3.
7 Sedum palufre purp. Mont. humid. R. 691. 4.
8 Sedum minus Vicentii O. Rupibus R. 5.152. 3.
9 Illecebra lutea Vic. Tectis R. 1041. 12.
             10. Tllecebra alba mitis O Rupes R. 1041. 14.
      11.Helleborus fl. viridi O R arò. R. 697. 2.
12.Helleborus niger vulg. O. Sepes rarò. R. 698. 3.
LAB.XI.II. 1 Caltha Palyfris O. L. Aq. R. 700. c. 9.
2.Ramunculus glossus. O Mont. R. 700. 1.
3. Juncus floridus O. L. Fossis limo fis. R. 701.
        3. Juncus floridus O.I. Foisis limofis .R. 701.
4. Aquilegia sylv. O. Dumetis R. 706.1.
5. Damasonium stellatum O.I. Stagnis R. 701.
6. Plantago aq: latifol. O.I. Stagnis aq. R. 618.1.
7. Plantago aq: longifol. O.I. Stagnis &c.
8. Plantago aq: minor O.I. Stagnis .R. 618.2.
9. Sagitta 6 riminea O.I. Fluvijs R. S. 619.
10. Sagitta latifolia. O.I. Rivul. R. 619.1.
        11 Sagitta angustifol OL R. 619.3.
```

```
Sect. X. Herbæ Bacciferæ Kay 652
   TAB XLIV 1. Bryonia alba 0. * L. Sepes R. 650. 1.
2Bryonia ngra 0. + L. Dumet K. 660. 2.
     3.Christophorian © Sylvis rare R. 662.1.
4.Ruscus © * H. Dumet R. 664.
5.Polygonatum vulg. © * Sylv. umbros R. 664.1.
6.Polygonatum A. maj. © Rupes Sylv. R. 664.3.
7.Lilium Convallium © * H. Sylv. umbros R. 667.1.
      8. Herba Taris O. +. Sylv umbros. R. 670.1
     9. Dulcamara O.* I. Aquos R. 6-2.1.
10. Solanum Hort. O.* I. Fimetis R. 672. 2.
     11. Solanum lethale O. Sepes. rnder. R. 679.1.
12. Asparagus Hort. O. *. Prat. marit. R. 683. 1.
   TAB.XLVI. 1 Barbarea O.L.* Minis agger R. 809. 8.

2 Barbarea pracox O. Folsis R. 809. 8. lut.

3 Erysimum vulg. O.* L. Ruder muris R. 810. 9.
      4 Erysimum latifol OL Agger ruder R. 810. 9.

5. Eruca aquatica OL Folsis aq. R. 808. 3. lut.

6. Eruca maritma Aren. marit R. 840. mir
             Eruca Monentis O. L. Ruder R. 807.4.lut.
      o Raphanus hort. . L. Serina R. 8041. pwp
      10. Rapishum albun D.L. Seget. Com. R. 805. 4.
      11. Rapistrum majus. Baro. R.806. 4. alb
      12. Sophia Chirwgorum. .* L.Ruder. R. 812. 10. lut.
  TAB. XIVI. 1. Nashurtium anwum O.L. Udis. R. 814.1.
      2. Nahm num aquaticum O.*. L. Vadosis . R. 816. 6. alb.
      3. Nashrutium aq. pracox. O.L. Vadosis. R.S. 172.7.

4. Nashrutium hirsut: mirus. O.L. Udis. R.815.5.

5. Cardamine prat. O.+ L. Udis. prat. R.814.2. can no

6. Cardamine ft. pleno. O. Raro. R.S. 171.2.
     7Cardamine inpatiens. Rup. udis. R.815. 4. alb. 8Cardamine triphyll. ©. Raro'. R.817. 11. 9Cardamine Bellidis. folio. ©. Rupis. raro'. R.817. 10. to Turritis vulg. © L. Murris. R.799 3.
      "Turrius major Q. Raro, R. 799. 5.
"Turius major Q. Raro, R. 799. 5.
  TAB XIXII I. Turritis invior. O.L. Aren. sicc. R. 800. 7.
A.Turritis invitor foliosa. O Muris. alb. Apr.
      3 Lunaria contortu major. Montib. udis R. 789.6.
     4 Lunaria contoria minori. Rupes. R. S. 165. 2. 5 Paronychia major. • Mont. R. 790.4 alb. 6 Paronychia vulg. * L. Siccis. R. 789.1. alb. 7 Paronychia b fida. • Siccis. alb. 8 Paronychia Hibernica aq. Lacubius. R. S. 181. alb. 6 Clabum contoria A. Angel Carlotte. R. S. 181. alb.
      9.Glastum sativum. Agris . Seritur . R. 842. lut.
     10L epidium rulg. O.*. Aquos. R. 8 28. 1. alb.
11Myagrum fylv. Seget. R. 8 20.1. lut.
12Brasica monospermos. Aren. marit. R. 8 3 8. alb.
TAB.XIX. 1. Cochleans Hort. ... *. Hort. mant. R. 8. 2. 2. 1. 2. Cochleans Murit. ... *. Salis mant. R. 8. 2. 2. 2. alb. 3. Cochleans Hederacea. Mant. R. 8. 2. 4. 6. alb. 4. Bursa Paltoris Offic. ... *. L. Vijs &c. Com. R. 8. 3. 8. 4. 5. Bursa Paltoris fol. Coronopi. ... L. Muris &c. alb. 6. Bursa Paltoris fol. unior. ... L. Muris Ag. g. er. alb. 7. Bursa Paltoris fol. integro. ... L. Muris &c. alb. 8. Raphanus aq. labfol. ... L. Aquis pigris. R. 8. 19. 3. lut. 9. Raphanus aq. nunor. ... L. Udis. R. 8. 18. 2. 10. Raphanus aq. fol. lacerat. ... L. Aq. pigris.
     10 Raphanus ay fol lacerat O.L.Aq. pigris.
11 Raphanus Ruficanus O.L. Folsis R. 818. 1. alb.
12 Raphanus Ruficanus fol lucerat. O.Aq. alb.
TAB.L.1. Nasturium fylv. O. Marit.R. 825. 2.
2 Nasturium petraum.L. Arenos. R. 827.9. alb.
    3 Nasturium petraum Cambriae. Rupibus. R. S. 174. 4.
4 Nasturium nunt parvam. Rupibus. R. S. 174. 5.
   4.Nathr tum Nont. parvan. Rupibus .R. 5. 174.5.
5.Nathr tum Valentinum. Raro'.R. 826.8.
6.Nathr tum verrucosum. L. Vijs. udis.R. 843. alb.
7.Thlafpi vuly. A.L. Seget. R. 830.1.
8.Thlafpi glabrum. Raro'.R. 5. 175. 5.
9.Thlafpi Diofeorid. +. Seget. &c. R. 831.5.
10.Thlafpi uranum. Montos. R. 831.2.
11.Thlafpi Glaft fol. Paleus mont. R. 832.10. alb.
    12Thlaspi perfol. minus . Saxosis. R. 832. 8.
```



```
12. Holosteum S. Alfine rulg. © 1. Com R 1030 20

Th. L. X. Alfine fol. Plantaginis • H Urginos & R 1032 5.

2. Alfine multicaulis • L Com Mur R 1031 1.

3. Alfine tennifolia • Seget R 1031 4

4. Alfine fl. pulchro • Mont R 1032 0

5. Spergula Sofetum • L Seget R 1034 19

7. Spergula narma • Sallis palustr R 1044 20

6. Spergula purp • H Arenos R 1034 18

9. Spergula purp • H Arenos R 1034 18

9. Spergula tetrapetala • H Pascuis Beril Aren R 1025 1.

10. Millegrana Graminea • R 1002 0

12. Millegrana minima • L Ericetis udis R 1020 7.

13. LX 7. Chauseillus vulg. • Mont Cret. R 1024 (3).

2. Chauseillus fel. meano • R 1005 7.

3. Chauseillus fel. poly • Pascuis

4. Chauseillus fel. poly • Pascuis

4. Chauseillus fel. putato • L & R 1015 7.

5. Hypericum pulchrum • L. Rupib R 1015 7.

6. Hypericum bishum • L. Rupib R 1021 19.

8. Hypericum bishum • L. Rupib R 1021 19.

8. Hypericum minimum • Pascuis Aren & R 1019 10.

9. Androsemum rulg. • * H. Dumet Sepib R 1020 16.

10. Androsemum Hypericoides • H. Dumet Sepib R 1020 17.

11. Alcyron vulg. • H. Rupis Ruft M 106 18.

2. Geum fol. curcinato. R 1047 1

3. Geum Hibernicum. R 1045 3

4. Geum Lambria. Rupib R 3. p. 507 2

5. Geum hirsutum. Mont humid R 1046 8.

6. Geum album Saxifrag a vulgo L * Pascuis fice R 1048

7. Geum hirsutum. Mont humid R 1046 8.

6. Geum mont liteum, Mont humid R 1040 6.

10. Geum mont liteum, Mont humid R 1040 8.

6. Geum mont liteum, Mont humid R 1040 1.

12. Parnaliia vulg. Palustrib R 1049 0.

14. Verbalcum lut fl. parve. R 1049 3.

3. Verbalcum narmm. R 1095 5.

5. Blattaria lutoa R 1095 1.

6. Salicaria fol. brevi.
1. Nervonica of pictar. Cambrine (i) R. S. S. S. S. Verronica and Olic. (i) s. L. Palcain sine. R. S. S. S. Verronica dybrate pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (i) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (ii) 1. Never hand for the pictar pack. obluga (iii) 1. 
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            3. Verbalcum niorum. R. 1095. 7
4. Verbalcum album. Vis R. 1095. 5
5. Blattaria lutea. R. 1096. 1.
6. Salicaria rulg. Ripis fluvior R. 1036. 1.
7. Salicaria fol. brevi.
8. Salicaria trifolia.
9. Lyfunachia lutea. Aquos R. 1021. 1.
10. Lyfunachia lutea. Aquos R. 1021. 1.
10. Lyfunachia olobosa., R. 1022. 5.
11. Pyrola rulg. * R. 1233. 1.
12. Pyrola rulg. * R. 1233. 1.
12. Pyrola mucronata. R. 1233. 2
13. Trientalis. R. 1101. c. 19.

TAB. LXIII. 1. Anagallis mas. • Seget. R. 1023. 1.
2. Anagallis famina. • * R. 1024. 2.
3. Anagallis famina. • * R. 1024. 2.
3. Anagallis lutea. 6. Sylv. Umbros. R. 1024. 4.
4. Anagallis lutea. 6. Sylv. Umbros. R. 1024. 4.
4. Anagallis lutea. 6. Sylv. Umbros. R. 1024. 4.
5. Nummularia rulga. 6. * R. 1099. 1.
6. Nummularia rulga. 6. * R. 1099. 1.
6. Nummularia rulga. 6. * R. 1098. 1.
10. Ros solis fol. volundo. 6. Aquos. R. 1055. 1.
10. Ros solis fol. rotundo. 6. Aquos. R. 1055. 1.
10. Ros solis fol. rotundo. 6. Vdis. R. 1100. 2.
12. Rorella longifol. max. R. 1100. 5.

TAB. IXIV. 1 Geranium columb. 6. L. Jepib & R. 1059. 18.
3. Geranium columb. 11. min. 6. L. R. 1059. 18.
5. Geranium columb. 11. min. 6. L. R. 1059. 18.
5. Geranium columb. 11. min. 6. L. R. 1059. 18.
5. Geranium columb. 11. min. 6. L. R. 1059. 18.
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   3. Geranium columb, minim • Pascuis steril,
4. Geranium columb, ft. min • L. R. 1059.18.
5. Geranium columb, ft. maj • L. Pascuis Arenos.
6. Geranium columb, discret fol. • Pascuis steril. R. 1059.21.
7. Geranium columb, discret fol. • Pascuis steril. R. 1059.21.
8. Geran columb, discret fol. ped. ft longos • R. 1059.23.
9. Geran hamatoko. • Ericetis mont & A. R. 1000.32.
10. Geran hamatolius incisis • Agger fossar
11. Geran hamatolius incisis • Agger fossar
12. Geran hamatolius incisis • Areno R. 1000.
12. Geran hamatolius Pussillum, Aren maux R. 1056.3.
2. Geran hawathum • R. 1057. m
3. Geran, thata fol. Arenos R. 1057. $.
4. Geran, nodosum Mont R. 1058.17.
5. Geran, Robert × Sepib. & R. 1058.17.
6. Geran, hatauboules, Praus humid R. 1061.33.
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         7. Geran, batrucheider, Prans humid R. 1061 33
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     7. Geran, Intrudivider, Pratis humid R. 1061-33
8. Geran, Intrudivider, Pratis Mont R. 1062-34.
9. Anthyllis maril. Arenos marit 1024-c.11.
10. Glaux maril. Salfis Paluftr, R. 1102-1.
11. Portulaca aquat. Udis R. 1035-25.
12. Plantaginella palustris. R. 1077-c.12.
TAB. LXVI.1 Allium Sylvestre R. 117-4.
2. Allium Sylv. Tennifolium. R. 1118-4.
4. Allium Sylv. Tennifolium. R. 1118-4.
4. Allium Sylv. Nucleis rariorib et majorib.
5. Allium Sylv. Amphicarpon &c.
6. Allium Sylv. Incomi. &c. Seget. R. 1119-3.
13. Allium Sylv. Incomi. &c. Seget. R. 1119-3.
14. Allium Sylv. Incomi. &c. Seget. R. 1119-3.
15. Allium Sylv. Incomi. &c. R. 1125-2.
16. Ornithogalum spicatum filtrenii R. 1153-11.
17. Ornithogalum vulgare. R. 1153-13.
18. Ornithogalum luteum. R. 1153-13.
```



TAB. LXVII. 1. Bulbofa Alp. juncifol. Cambria

- 2. Colchicum. Pratis R.1170.1.
- 3. Crocus * R.1176.17.
- 4. Hyacinthus Angl. Dumet Sep. R. 1159.20.
- 5. Hyacinthus Stellatus Cambria R. 1155.1.
- 6. Hyacinthus Stell. antumalis Rupil. R. 1157.13.
- 7. Gladiolus lacustris. R. 1325. c. 12.
- 8. Narcissus pallido. lut. angl. alt. colyee magn. brev. aurao.
- 9. Narcifsus Sylvestris. R.1131.5
- 10. Narcusus medio-luteus vulg. R. 1135.23.
- n.Iris Palustris lutea Rip. fluvior.R.1186 . 24.
- 12. Xyris. Dunet. Sep. R. 1190.7.

LAB.LXVIII.1. Orchis barbata fatid @ R.1212.1.

- 2. Orchis morio mus O Pascuis &c.R.1214.7.
- 3. Orchis morio fam. O Pascuis M. R. 121.1. S.
- 4. Orchis purpurea. O Pascuis sice. cret. R. 1215 . 14.
- 5. Orchis Pannonica. 3 Collib. Steril et Cret.R.1215.12.
- 6. Orchis odorata moschata @Y.R.1217.22.
- 7. Triorchis. O Pascuis Arenes Y.A.R. 1217. 26.
- 8. Orchis anthropophora. o Glareos. R. 1217.27.
- 9. Orchis *galea*. &c. ⊙ R.1218.29.
- 10. Orchis myodes minor O Agger. foss.R. 1219.32.
- 11. Orchis hermophrodition bifol. 9. R. 1221.40.
- 12. Orchis all. bifol .min. calcari. oblongo. R. 1221.

TABLEXIX.1. Orchis nugodes major. @ R.1220.33.


- 2. Orchis fuerflorade 3 Pascuis suc .R.1220.33.
- 3. Orchis sive Testiculus Sphegodes hirsut. fl.R.1221.39.
- 4. Orchis all. odorata. O Eborac.
- 5. Orchis palmata maj.mar. 3 Pratis himid R. 1223.1.
- 6. Orchis palmata famin. O Sylvis. I.R. 1223. 2.
- 7. Orchis palmata rul. O Pratis humid.R. 1224. O.
- 8. Orchis palmata min. @ Pascuis vice. R. 1224.7.
- 9. Pleudo-Orchis. 3 Palustr. R. 1226.3.
- 10. Orobanche major. 0 R. 1227 . 1.
- 11. Orobanche ramosa. 9 Seget.R. 1227.5.
- 12. Orobanche luppopitys. 3 Sylv.umlros R. 1229. 2.

- TAB. LXX.1. Orobanche radice dentata may. @ R. 1229.1.
 - 2. Nidus Airis. 6 Sylvis R. 1228.1.
- 3. Nídus Airis, purp. @ R. 1229. 3.
- 4 Calceolus Maria o Sylvis . R. 1239 . 12 .
- 5. Helleborine fl. utro rubente. @ Mont. R. 1231. 5.
- 6.Helleborine latifel.mont. o Sylvis caduis. R. 1230. 1
- 7. Helleborin e /l. allv. o Sylvis.M.I. R. 1231.6.
- 8. Helleborine palustris. O Udis &c. R. 1231.10.
- 9. Helleborine augustifolio. O Sylvis kr.
- 10. Bifolium Majus . Pratis . humid . &c . R . 1232 . 1 .
- 11. Bifolium minimum Ericelis udis . R. 1232 . 2 .
- 12. Bifolium palustr. Udis. palustr. R. 1232.4.

TAB. LXXI.1. Nymphæ alla o Fluvijs. R. 1320. 3.


- 2. Nymphæa all minima . Aquis pigris. R. 1320. 4.
- 3. Nymphæa lutea. @ Fluvijs. R. 1319. 1.
- 4. Nymphæa latea min. fl. fimbriato R. 1320. 2.
- 5. Aloe palustris. Fossis palustr. R. 1324. c. 7.
- 6. Filipendula @ Pratis & pascuis mont. R. 623.1.
- 7. Filipendula.unin. o Pascuis steril.R. 023.2.
- 8. Umaria O Ripis flucior. R. 624. c.3.
- 9. Thalietrum majus. o Ad rivul. Lag. R. 403.4.
- 10. Thalictrum munus. o Mont. cret. R. 404.5.
- 11 . Thalictrum mont.minus. O Rup. R. 3 . p. 249.
- 12. Thalictrum awnt. minim. 9 Rup. udis. R. 3. p. 249.
- TAB.LXXII.1. Valeriana Sylvestris major. 0 * R. 3 8 8.2.
 - 2. Valeriana Sylv.min. o Pratis Humidis. R. 388.3.
- 3. Valeriana Sylv. seu Palustris minor altera. R. 389. 4.
- 4. Lactuca aguina . Seget &r. R. 392.1.
- 5. Lactuca agnina fol serrates o Seget . R. 302.2.
- 6. Limonium o Salsis palustr. R. 395 . 1.
- 7. Limonium mia . 3 Clivir. Marit I.R. 395.
- 8. Caryophyllus mout. min. @ Palustr. marit. R. 1037.11.
- 9. Spargantum vantosum v Rip. fluvier kc. R. 151. 2.
- 10. Spargan. nen ramosum.o
- 11. Spargan. minimum & Stagnis & ricul. R. 1312.
- 12. Cuscuta mayor. Ericetts I.Y. R. 1003. c. 23.











Signifies ANNUAL
 Signif. PERENNIAL

S.N Sculp:



ENGLISH PLANTS with BLINK Flowers. T_AB . III. 3. Barren SCRREL. 4 Welsh SORREL 1. Common SORREL. 2. Sheeps SORREL. RAY Syn 31.10. ed. 57. 10.


RAY 178. L.S. 3 L 8. ed 56.8. RAY 180. 14. S. 3 1. 9 id 56.9.

LONDON Meadows Tune Vertue. WALES

5. Common ARSMART. 6 Creeping ARSMART. 7 Spotted ARSMART. 8. Spotted houry ARSMART. RAY 183.3.S.32.ed.58.2. RAY 183.2.S.32.3.ed58.3. RAY 182. S.32. ed 58. L



9. Silver ARSMART. 10. White branch'd ARSMART. x1 BOBARTS Arsmart. 12. Amphibious ARSMART.





..

•


*

ENGLISH PLANTS with BLINK Flowers TAB: III


1. Common PLAN TAIN. 2. Dented PLAN TAIN. 3. Hoary PLAN TAIN. 4 Rose & Spiked PLAN TAIN.

RAY 876.1. Syn. 125.1.ed 184.1.


RAY 876.1. Syn. 125.1.ed 184.1.



5Besom PLAN TAIN. 6. Common RIBWORT. 7. Mountain RIBWORT. 8. Childing RIBWORT. RAY 877.4. Syn. 17.6.3.ed. 185.3. RAY 877.7. Syn. 126.4.ed. RAY Syn. 126.7.ed. 185.7. RAY 877.1.




9. SeaPLANTAIN. 10.BUCKS-HORN PLANTAIN. XI. GreatBURNET. 12. Small BURNET.
RAY880.5. Syn1265. ed. 1855. RAY879.3. Syn. 126.6. RAY402.2. Syn. 162.1. ed. 100. RAY401.1. Syn. 162.2. ed. 100.
ed. 185. 6.






ENGLISH PLANTS with BLINK Flowers. TAB. V.






ENGLISH PLANTS with BLINK Flowers TAB. VI.


1. Tassel PONDWEED. 2. Horned PONDWEED. 3. Water STARWORT. 4. Long Water STARWORT.
RAY 190. 10. Syn35. 10. ed 62. 10. RAY 190. 13. Syn35. 11. ed 281. RAY 1323. cap. 6. Syn 208. 1. RAY Syn 208. 2. ed 280. 2.



6. Fine Feather PONDWEED. 7. Milfoil PONDWEED. 8. Knotted PONDWEED, 5.FeatherPONDWEED. RAY 1920.2.2. S.35. ed 2805. RAY 1322. L. Syn. 207. ed 279.



wLong Golden SAXIFRAGE. 9. Golden SAXIFRAGE. RAY 207.C.XI. I Syn38.L. ed.66. RAY 207. 2. Syn39.ed. 66. XL CORNWAL PENNY WORT. 12. Marsh PENNY WORT.





1,5

ENGLISH PLANTS with BLINK Flowers TAB. VII.

Frosty Sea ORRACH. 4. Jagged long ORRACH. 1. Delt ORRACH. 2. Delt Sea ORRACH. RAY 192.3. Syn36.1. edő2.1. RAY 193. 10. Syn36.4.ed.63.4. RAY 193.8. Syn36.5. ed.63.5. RAY 192.6. Syn36.3. ed.62.3.




5. Spear ORRACH.

6. Grass Sea ORRACH. 7. Sea PURSLAIN. 8. Heart Sea PURSIAIN.



9. Small Garden BLITE. 10. Allseed BLITE. 12. Wild SPINAGE. x1. Stinking ORRACH.




Sutton Nicholls sculp.




ENGLISH PLANTS with BLINK Flowers TAB. VIII.

L. RAND'S Oak BLITE. 2. Frost BLITE. 3. BUDDLE'S Fig BLITE. 4. BUDDLE'S RoundBLITE.


RAY 198.8. Syn. 36. ed 63. 1.



g. Thick Shining BLITE. 6. Sharp pointed BLITE. 7. Maple BLITE. 8. Broad pointed BLITE. RAY Syn.ed. 64.5. RAY 197.9. Syn.37.3. ed. 64.3.




9. Sea BEET. 10. Wall PELLITORY. x1. Tree STONECROP. 12. Shrub STONE CROP. RAY 204. 3. Syn. 38. ed. 65. RAY 206. 1. Syn. 38. ed. 66. RAY 199. 17. Syn. 38. ed. 65. 10. RAY 199. 16. S38.7. ed. 65. 9.





ENGLISH PLANTS with BLINK Flowers. TAB. IX.

1. Small GLASSWORT. 2. Prickley GLASSWORT. 3. Small joynted GLASSWORT. 4. Woody joynted GLASSWORT. RAY 108. 14. Syn 374 ed 64.7 RAY 212. 2. Syn 39. 3. ed 67. 3. RAY syn 39. Led 67. L. RAY 212. Syn 39. ed. 67. 2.





5 Bastard ToAD-FLAX. 6. Upright KNAWELL. 7. Spreading KNAWELL. 8. PEPPER 6 rass.

RAY 399.c.7 Synbied 99. TABERN. 505. RAY. 213.1. Syn40.1. ed. 68.1. RAY 1325.c.w. Syn209.ed 281.



9. LADIE S MANTLE. 10. Small LADIES MANTLE. XL Silver LADIES MANTLE. 12 PARSLEY PIERT.
RAY 208. 1. Synzg. 1. ed. 66. RAY Hist Plantar. 202. 2. RAY 209. 4. Synzg. ed. 66. RAY 209. c. 14. Synzg. ed. 67.






ENGLISH PLANTS with BLINK Flowers TAB. X.


1. Common KNO TGRASS. 2. Thick-set KNO TGRASS. 3. Short leav'd KNO TGRASS. 4. Narrow KNO TGRASS. RAY 184. 1. Syn 33.ed59.

TABERN. Vol. 2. 202 Ic. 833. TABERN. V. 2. 203 Fig.




5. SeaKNOTGRASS. RAY 185. 2. Syn33. ed. 59.2 6.NEW TONSKNOTGRASS. 7.WhorldKNOTGRASS. 8.VALENTIA.Knotgrass.

RAY214.3.Syn40.ed. 68. RAY215.8.




9. Smooth RUPTURE-WORT: 10. Hoary RUPTURE-WORT XI. Sea HEATH. 12. BLINKS.




Sutton Nicholle sculp.




FNGLISH PLANTS with LETTUCE FLOWERS. TAB. XI.
1. Common Mouse-EAR. 2. WELCH MOUSE-EAR. 3. Narrow Mouse-EAR. 4. Brad Mouse-EAR.
RAY 242.1. Sun. 46. ed. 75. RAY. 241. 19.
RAY 244. 2.



5. Narrow Sincoth MOUSE-EAR. 6PLUKENET'S, NOUSE-EAR. 7. Common DANDELION. 8. Narrow DANDELION. RAY 245.8. Syn. 45. ed. 76.4. RAY syn. ed. 2.p. 75. x1. RAY 244.3. Syn. 46.1 ed. 76.1. RAY Syn. 46.2. ed. 76.2.




9. Common rough DANDELION. 10. SANDY rough DANDELION. x1 Rough Branched DANDELION. 12. Smooth branched DANDELION. 12. Smooth





ENGLISH PLANTS with LETTUCE FLOWERS. TAB. XII.

L. Comon HAWKBIT. 2. Jagged HAWKBIT. 3 Small Jagged HAWKBIT. 4. Dandelion HAWKBIT. RAY 230.1. S.42, Led. 71.1.



5. Buddle HAWK-BEARD. 6. Comon HAWK-BEARD. 7. Succory HAWK-BEARD. 8. Castor HAWK-WEED

RAY 234.16. S. 43. 4. ed. 72. 4.

RAY 232. 6. S. 43. 3. ed. 72. 3.



Charlon
Greenwich & C.

Vell

KENT

Jan

KEN

KENT

JAN

KEN

KENT

JAN

KEN




ENGLISH PLANTS with LETTUCE FLOWERS. TAB. XIII.

LBroad HAWK-LUNG 2. Round HAWKLUNG 3. Broad dented HAWK-LUNG 4. French HAWK-LUNG RAY 239. 7. S. 45. 6. ed. 73. x1. RAY Syn. edit. 2. p. 74. 6. RAY Syn. ed. 2. pag. 74. 7. RAY 239. 9. S. 44. 4. ed. 74. 4.



6 WEICH hoary HAWK-LUNG 7 Broad hairy HAWKLUNG 8 Long havry HAWKLUNG g Narrow HAWK-LUNG 6. WELCH hoary HAWK-LUNG RAY 240 x L S 45 5 2 74 5. RAY 239 8 Syn 45 ed 75 8. RAY 238 2 S 44 Led 74 L.








ENGLISH PLANTS with LETTUCE Flowers. TAB. XIV.

1. Round leav'd SOW THISTLE. 2. Broad SOW THISTLE. 3. Narrow SOW THISTLE Prickle dented SOW THISTLE
PLUKENET. Tab. 61. Fig. 5. RAY 225. 7. Syn. 42. 4. ed 71. 4. PLUK. Tab. 62. fig. 4. Dodon Pempt 643. fig. 3.




5. Prickle jagged SOW THIS TIE. 6. Corn SOW THIS TIE. 7. Tall Marsh SOW THIS TIE. 8. Smooth Broad SOW THIS TIE. RAY 225.6 Syn. 42.3. ed. 70.3. RAY 226.10 Syn. 42.5 ed 71.5. RAY 226.11 Syn. 42.6 ed 71.6. RAY 224.2 Syn. 42.2 ed. 70.2.



9. Smooth Jagged SOW THIS TIE. 10 BUDDLE'S Jagged SOW THIS TLE. XI Wild SUCCORY. 12. NIPPLE WOR T.

RAY 224.1. Sun 42.1. ed 70.1.

RAY 256.4. Syn 47.ed 77. RAY 256.6.2. Syn 47.ed 77.



n Nicholls souly



ENGLISH PLANTS with LETTUCE Flowers &c. TAB. XV.


2.1



5. Ivy LETTUCE. 6. Narrow GOATS-BEARD. 7. Comon GOATS-BEARD. 8. Purple GOATS-BEARD. RAY 252.1. Syn41.5. ed 70.5. RAY 252.6. RAY 252.1. Syn46. ed. 76. RAY 252.3. Syn47. ed. 77.




9. Small SWINES SUCCORY. 10. Common CARLINE. XI BUTTER-BUR Leaves. 12 BUTTER-BUR in FLOWER.
RAY 229. 2. Syn. 44.12. ed. 78. RAY 288. 2. Syn. 247. ed. 80. RAY 260. Syn. 47. ed. 78.






.






ENGLISH PLANTS with STAR-FLOWERS. TAB. XVII.


1. Common RAGWORT. 2. Marsh RAGWORT. 3. Groundsel RAGWORT. 4. CAMBRIDGE Ragwort
RAY 284. 1. Syn. 50. 1. ed. 82. 1. RAY 285. 4. Syn. 50 3. ed. 82. 3. RAY 285. 2. Syn. 50. 2. ed. 82. 2. RAY 272. 4. Syn. 49. 3. ed.



G. Common GROUNDSEL. 6. Cotton GROUNDSEL. 7. COLTSFOOT.
RAY 290.1. Syn.50.1. ed. 83.1. RAY 290.2. Syn.50.2. ed. 83.2. RAY 259.1. Syn.47. ed. 78.



9. Golden SAMPIRE. 10. Great Sca STARWORT. x1. Small Sea STARWORT. 12. Naked Sea STARWORT.
RAY 268. x1. Syn. 48. 1. ed. 79. 1. RAY 270. 20. Syn. 49. 2. ed. 80. 2. RAY 270. 20. Syn. 49. 2. ed. 80. 2. RAY 270. 20. Syn. 49. 2. ed. 80. 2.







.

160

ENGLISH PLANTS with naked FLOWERS & Downey SEED TAB XVIII.








1 1

.


è

ENGLISH PLANTS with DAISIE FLOWERS. TAB. XIX.





ENGLISH PLANTS with BALD or NAKED FLOWERS. TAB. XX.




,就是我们,我们就会不是不是一个。 我们就是我们的,我们就是我们的,我们就是我们的,我们就是我们的,我们就是我们的。 我们就

ENGLISH CAPITATED PLANTS with JAGGED HOLLOW Flowers TAB.XXI.

1. Mush Thistle. 2. Welted Thistle. 3. Gray Thistle. 4. Marsh Thistle.


RAY 308.1. Syn.53.1. ed86.1. RAY 309.2. S53.2. ed862. RAY 309.3. S53.2. ed.86.3. RAY 309.4. S53.4. ed86.4.



5. CommonTHISTLE. 6Dwarf.THISTLE. 7. Spear.THISTLE. 8. Fryars. THISTLE. RAY.310.6.5.3.5.ed. 87.5. RAY.310.6.5.53.6.ed.87.6. RAY.310.1.5.54.7.ed.87.7. RAY.311.3.5.54.8.ed.87.8.




9. Milke. THISTLE. 10. Cotton. THISTLE. 11. Star. TH. STLE. 12. Yellow THISTLE. RAY312.1.5.54.9. ed. 87.9. RAY313.1.5.5411. ed. 87.12. RAY317.1.5.5412. ed. 87.12. RAY317. 45. ed. 88. 13.





ENGLISH CAPITATED PLANTS with lagged Hollow Flowers TAB. XXII




* I* \

5 (1 A) (2) [fiv | fiv) [anner [anner

.


l. BURDOCK. RAY.332.2. |2.Rose.BURDOCK | 3. Small headed BURDOCK4 Small woolly headed BURDOCK.



5. Great woolly headed. BURDOCK. 6 Cobweb headed BURDOCK. 7. Manured TEASEL 8. Wild-TEASEL RAY. 332.2. S. 55. ed. 88. 2. RAY 382. 1. S. 59. ed. 96. RAY 382. 2. S. 59. ed. 96.



9 Small wild TEASEL 10. Sea ERINGO x1. Common ERINGO 12. SANICLE RAY 382. 4. S. 60. 3. ed. 97.3 RAY 384. 3. S. 60. 2. ed. 97. 2. RAY 384. 1. S. 60. 1. ed. 97. 1. RAY 475. c. 4. S. 72. ed. 115.





1 . 1

ENGLISH PLANTS with UMBELLIFEROUS Flowers TAB.XXIV.

2 COW PAR SNEP. | 2 Lagged Cow. PAR SNEP | 3. Garden PAR SNEP. | 4. Wild. PAR SNEP. | RAY. 400. 2 Syn. 63. ed. 102. | RAY. 400. 2 Syn. 63. ed. 102. | RAY. 409. Syn. 63. ed. 102.




5. Great HARTWORT. 6. Small HARTWORT. 7. HOGS. FENNEL. 8 SAMPIRE.

RAY. 411. 1. 2. Sozedio 2. RAY. 416.1. Syn. 63. ed. 102. RAY. 45.7. 2. S. 69. ed. 111.



9.ANGELICA. 10.Water.ANGELICA. 11.CORIANDER. RAY. 434. 25.64.ed. 104. RAY. 470. S. ed. 114. RAY. 437.1. S. 65. ed. 104.







ENGLISHUMBELLS with long and round SEED.

TAB.XXV.


IVENUS Comb 2.COWWEED. 3.Wild.CHERVIL. 4.SPIGNEL or MEU.

RAY 428.15yn.64.ed.103.RAY.429.15yn.64.ed.103.RAY.431.6.Syn.64.ed.103.RAY.432.1.Syn.64.ed.104.





9. EARTH NUT. 10. GOUT WEED. 11. Barard stone PARSLEY. 12. HONEWORT.
RAY. 440. 1. Syn. 65.ed. 105. RAY. 435.4. Syn. 65.2.ed. 104.2. RAY. 443.3.5.66.ed. 106. RAY. 443.4. Syn. 66.ed. 106.






ENGLISH UMBELLS with small round SEED TAB. XVI

1.BROAD WATER-PARSNEP Narrow WATER-PARSNEP Creeping WATER-PARSNEP. Least WATER-PARSNEP.

RAY 443.5. Syn. 663. ed. 107.3. RAY 444.4. S. 66.4. ed. 107.4. RAY 444.5. S. 665. ed. 107.5. RAY 444.6 S. 67.6. ed. 107.6.




5. Great; broad SAXIFRAGE. | 6. Great Cut SAXIFRAGE. 7. Small broad SAXIFRAGE. 8. Small Cut SAXIFRAGE. RAY445.1. S. 67.1. ed. 108. 1. | Hist. Oxon. | RAY445.3. S. 67.2. ed. 108. 2. | RAY445.4. S. 67.3. ed. 108.3.






ENGLISH UMBELLS with ROUGH SEED. TABXWIL

1 Common FENNEL. 2ROCK PARSLEY. 3 Mountain STONE PARSLEY. 4 Common CARROT RAY 457.1. SYN. 69. edm. RAY 460. 5. SYN 69 edm. RAY 463. 5 SYN 70. edm. RAY 465.1 SYN 701 ed.




9 Rough PARSLEY 10 Corn Rough PARSLEY. 11 Knotted Rough PARSLEY. 12 Fine Rough PARSLEY. RAY 468. 6 SYN713. edn3.3. RAY 468.7 SYN713 ed. 113.3. RAY 468. SYN71. 5. ed. 1145. RAY 469.9. SYN.71. 6 ed.






ENGLISH UMBELLS & PLANTS with Rough Leaves. TAB. XXVIII.


1 Great Water HEMLOCK. 2 Common HEMLOCK. 3 Lesser HEMLOCK 4 The Water HEMLOCK RAY 450.2 SYN68ed 10 9. RAY 451.2 SYN68 Led 110. LRAY 451.3 SYN68ed 102, RAY 452.6 SYN 69 ed 110.



5 Meadow SAXIFRAGE 6 Prickly SAMPIRE 7 Least HARES EAR 8 THOROW WAX RAY4531SYN.69.ed.110. RAY4691 SYN 69 ed 111. RAY47.44 SYN 72 ed 115. RAY 47 1 SYN 72 ed 115



9 Common BORAGE 10 Small Wild BUGLOSS x1. Great HOUNDS TONGUE 12 Green Houres TONGUE RAY 493 SYN.76.ed.120. RAY 494 3 SYN.75.ed.119 RAY 489 1. SYN.751 ed 119.1 RAY 491.8 SYN.752.ed 119.2



PARTITION OF ALLES


SEL MINATO

To Wall and Oak


ENGLISH PLANTS with Rough LEAVES, & comoulty 4 SEED TAB XXIX

**Nipers BUGLOSS. 2. Small Vipers BUGLOSS. 3. Sea BUGLOSS. 4. Long BUGLOS S. Cowslips.


RAY 498. L. Syn. 75. edu 9. RAY Syn. edit. 119. RAY 500. 13. S. 76. 2. edu 9. RAY 489. 3. Syn. 75. edu 18.




5. COMFREY. 6.Blew GROMILL. 7. Shop GROMILL 8.Bastard GROMILL.
RAY 505.1.S.77.ed121. RAY 503.2.S.76.ed.120.2. RAY 503.1.Syn.76.1.ed.120.1. RAY 503.5.Syn.76.3.ed.120.3.



9. Water SCORPION GRASS. 10. Field SCORPION-GRASS. 11. Small SCORPION-GRASS. 12. German MADWORT. RAY 504.3.5.77.3. ed. 121.3. RAY 504.1.5.77.1. ed. 121.1. RAY 504.2.5.77.2. ed. 121.2. RAY 502. c8.5.76. ed. 120.










ENGLISH Whirl PLANTS with Hooded & Lipt FLOWERS. TABXXXI

160mmonMOTHER of THYME. 2Large flow red MOTHER of THYME. 3Upright broad THYME. 4LEMON THYME.


RAY 5.21.4. Syn. 78. 1. ed. 1.2.1. RAY 5.21.4. Syn. 78. 2. ed. 1.22.2. RAY 5.21.2. RAY 5.21.2.



5. Mater WHIRL-MINT 6. Common Water MINT 7. Red MINT 8. Crofs Whirl MINT RAY 530. 1. Syn 78. 1. ed 123. 1. RAY 533. 12. Syn 78. 3. ed 123. 4. RAY 530. 2. Syn 78. 2. ed 123. 3. RAY 531. 3. Syn . edit. 124. 6.



oEssexRepperMINT 10EALES Repper MINT 11Long HORSEMINT 12Round HORSEMINT RAY5327. Syn.79.5. ed. 124.2 RAY Syn. ed. 124.5. RAY532.0. Syn.79.6. ed. 124.3. RAY532.7. Syn.79.7. ed. 124.4.




The state of the s

त्याः। = व्यापः व्यापः

1 / 45F (15

ENGLISH Whart PLANTS with hood & light FLOWERS. T.IB. XXXII.


LNEP or CAT-MINT. 2 Pensur Royal Pudding grafs. 3. White HOREHOUND. 4. Stinking HOREHOUND. RAY. 548. L. Syn 80. ed. 125. RAY. 533. L. Syn. 79. ed. 125. RAY. 556. L. S. & L. ed. 128. RAY. 571. L. Sim. 84. ed. 132.



5. Water HOREHOUND. 6. Wood BETONY. 7. HEDGE NETTLE. 6. ENGLISH CLARY. RAY.535. 15. 79. ed. 125. RAY. 550. 1. S. 80. ed. 127. RAY.548. Syn. 80. ed. 127. RAY.545. 12. Syn. 80. ed.




9. Great Wild BASIL 10. Small Wild BASIL 11. Common SELF-HEAL. 12. B.y. HOREHOUND RAY, 550, 17. S81, ed. 120 RAY, 552, 1. Sept. Sept. 128. RAY, 551, 128. RA





ENGLISH PLANT Swith Hood & Lipt TLOWER S. TAB XXXIII.


1. Ground Dry, or ALEHOOF. 2. Red ARCH-ANGEL. 3. Red cut leaved ARCHANGEL 4. Great HENBIT.
RAY 567. 1. Syn. 131. ct. 83. RAY 559. 3. Syn. 129. ct. 82. 2. RAY 560. 6. Syn. 129. 3. ct. 82. 3. RAY 560. 10. S. 129. 4. ct. 82.4.



g.WhiteARCHANGEL. 6 Vellow ARCH-ANGEL: 7. BaftavdBalm 8. Purple NETTLEHEMP. RAY559.1.S.129.1.e4.82.1. RAY560.11. Syn. 129. 5. e4.82.5. RAY561.13. Syn. 129. 65.82.6. RAY561.13. S. 129. 7 Syn. 82.7.



9. Clowns ALL HEAL: 10. Yellow IRONWORT: 11. Narrow IRONWORT: 12. Round IRONWORT: RAY563.3. S. 130. 1. e.t. 83.1. RAY566.15. S. 130. 3. e.t. 83.2. RAY566.16 S. 130.4 e.t. 83.4 RAY566.16 RAY566.16 S. 130.4 e.t. 83.4 RAY566.16 RAY566.16 RAY566.16 RAY566.16 RAY566.16 RAY566.16 RA





* ENGLISH Whirl PLANTS with Hood or Lipt FLOWERS. TAB XXXIV.

1. Common CALAMINT. 2 Reld CALAMINT. 3. Common BUGLE. 4. Mountain BUGLE.


RAY569. 1. Syn. 83. cd. 131. 1. RAY569. 2. Syn. 83. cd. 132. RAY575. 1. Syn. 84. ed. 133. 1. RAY575. 2. Syn. 85. 2. cd. 133. 2.




5.MOTHERWORT 6. SCORDI.UM. 7.Wood SAGE. O. Wild MARJORAM. RAY571.1.S. 05.ed. 134. RAY576.c.30S.05.ed.133. RAY576.c.29.Syn.05.ed.133. RAY539.4.Syn.80.ed.126.1.




9.GROUND-PINE · 10. HOOD-MINT · 11. Small HOOD MINT · 12. VERVAIN · RAY573.1.S. 84. ed. 132 · RAY572.1.S. 84. ed. 132 · RAY572.1.S. 84. ed. 132 · RAY535.1. Simil 79. ed. 125 ·






ENGLISH CAPSULES with SNOUT FLOWERS. TABANN.


i. Small FO AD -FLAX² Smalt SNAPDRAGON. 3 Pointed FLUELLIN. 4Round FLUELLIN. RAY 760. 35 Small FlueLLIN. 4Round FLUELLIN. RAY 760. 35 Small FlueLLIN. RAY 759.35 Sno5.3. ed. 160.3. RAY 759.34 Sno5.4. ed. 160.4.



5.Comon TOADFLAX. 6.Bluijhweet TOADFLAX. 7.BUTTERWORT. 8. Purple FOXGLOVE. RAY752.1.S.105.1.ed.160.1 RAY756.20.S.105.2.ed.160.2. RAY751.1.Syn.104.1.ed.159.1.RAY767.1.Syn.106.ed.161.



9 Brown IIG-WORT 10. Water IIG-WORT 11. Sage IIG-WORT 12. Comon HENBANE RAY 764.3. S. 106.2. ed 161.2. RAY 764.1. S. 106.1. ed 161.1. RAY 766.11. Syn. ed 161.3. RAY 711.1. Syn. 101. ed 155.







Programme Committee Commit

ENGLISH CAPSULES with SNOUT FLOWERS. IABXXXII


2. Quick in Hand Touch me note Yellow RATTLE 3Tall Red RATTLE 4Comon Red RATTLE
RAY 1328. c. v. S. 209. ed. 282. RAY 769. 15. 106. 1. ed. 162. 1. RAY 770. 6. Syn. 106. 3. ed. 162. 3. RAY 770. 5. 5. 106. 2. ed. 162. 2.



5.Comon EYE-BRIGHT 6.Yellow Mar/h EYE-BRIGHT, Red EYE-BRIGHT 8.ORTON rat EYE-BRIGHT RAY772.4. Syn. 107. 4 ed. 163.4. RAY772.3. S. 107.2. ed. 162.2. RAY773.5. S. 107.3. ed. 162.3.



9lomon COW-WHEAT. 10. Gefted COW-WHEAT 11. Great BLADDERS NOUE 2 Small BLADDERS NOUE RAY 775. 4. S. 108. 2. ed. 164. 2. RAY 775. 3. S. 108. 1. ed. 164. 1. RAY 133 2. 2. Syn. 207. 2. ed. 279. 2. RAY STN OP S. 208. 3. ed. 279. 3.




The state of the s


TIAM COOK

ENGLISH HERBS with Heel'd FLOWERS.&c. TAB. XXXII


1. Purple VIOLET. 2. Double VIOLET. 3. White VIOLET. 4. Dog's VIOLET.
RAY 1049. 1. S. 152. 1. ed 214. 1.
RAY 1050. 2. S. 152. 2. ed 214. 2. RAY 1050. 3. S. 152. 2. ed 214. 2. RAY 1050. 3. S. 152. 2. ed 214. 2.



5. Bog VIOLET. 6 Yellow Rock VIOLET. 7. Havy Wood VIOLET. 8. Garden PANSIE.
RAY1050.4. S. 15.2.4. ed 214.4. RAY1052.6. S. 153. 10. ed 215.10. RAY1051.5. S. 152.6. ed 215.6. RAY1052.2. S. 152.7. ed 215.7.



9. Corn PANSTE. 10. Great Yellow PANSTE. 11. Base DYERS-WEED. 12. Comon DYERS-WEED. RAY 1053.7. S.153. 9. ed 215. 9. RAY 1052. 5. S.153. 8. ed 215. 8. RAY 1053. 2. ed 216. 1. RAY 1054. 8. S.153. 2. ed 216. 2.






ENGLISH PLANTS with Rough HEAD S, as CROWFEET. TAB XXXVIII.


1. PILEWORT. 2. Sweet Mood CROWFOOT. 3. Upright Meadow CROWFOOT. 4. Bulbous CROWFOOT. RAY.579. usyn. 85. ed. 134. RAY.584. usyn. 85. 6. ed. 135. RAY.583.7. Syn. 86. 6. ed. 135. RAY.581. 2.5. 862. ed. 135.2.



g. Pale upright CROWFOOT. 6. Doodys Trifid CROWFOOT. 7. Creeping CROWFOOT or Butter cups. 8. Thames CROWFOOT RIX.581.1. Syn.86.1.ed.135.1.



9.RAY.8 Hairy CROWFOOT. 10.Corn CROWFOOT. 21 SMALAGE CROWFOOT. 12 Joy CROWFOOT. RAY.583. 10.S.86. 5. ed. 135.5. RAY.585. 1.S.87.7. ed. 136. RAY.585. 1. Syn. 87. 8. ed. 136. 1. RAY.586. 2. Syn. 87. 9. ed. 136. 2.




The state of the s




ENGLISH PLANTS with Rough Heads &c. TAB.XXXXX

1. Water CROWFOOT with various Leaves. 2. Fine Water CROWFOOT. 3. Fine to inced Water OROWFOOT.

RAY. 506. 3. Syn. 87. 10. ed. 136. 3. RAY. 586. 4. S. 87. 11. ed. 136. 4. RAY. Syn. 24. 2. ed. 136. 5: 4 Fennell Water CROWFOOT.



5. Great SPEAR-WORT. 6. Small SPEAR-WORT. 7. MOUSE TAIL. 8. ADONIS Hower Red Mithes RAY, 587.8. Syn. 88. 14. ed. 137.8. RAY, 587.7. S. 88. 13. ed. 1377. RAY, 1332. C. v. S. 209. ed. 282. RAY, 596. 1. Syn. 88. ed. 137.




9. Common MALLOW. 10. Dwarf MALLOW. 11. Marsh MALLOW. 12. Vervane MALLOW RAY599.1. Syn. 88.1. ed. 138.1. RAY599.2. Syn. 88.2. ed. 138.2. RAY.602.1. S. 89. ed. 138. RAY.604.1. Syn. 89. ed. 139.





ENGLISH PLANTS with Rough HEADS & Naked SEED. TAB. XL.

1.AVENSHert BENNET. 2. Great flowred AVENS. 3. Mountain AVENS. 4. Childing AVENS.
RAY. 606. 1. S. 89. 1. ed. 139. 1. RAY. 606. 2. S. 89. 2. ed. 139. 2. RAY. 607. 3. S. 89. 3. ed. 139. 3. RAY. 607. 4. Syn. 89. 4. ed. 139. 4.




5. Cinquefoil AVENS. 6 Germander AVENS. 7. Common STRAWBERY. 8. Barren STRAWBERRY. RAY. 60 8.10.5. ed. 2,p.13 9. 6. RAY. 608.5.89.5. ed. 13 9.5 RAY. 609.1. Syn. 90. 1. ed. 140.1. RAY. 611. 2. Syn. 90. 3. ed. 140



9. Wood ANEMONY. 10. PASQUE Flower 11. TRAVELLERS JOY. 12. Plain TRAVELLERS JOY






ENGLISH PLANTS with Rough Heads & Naked SEED. TAB. XLI.


1PurpleMar/ACINQUEFOIL. 2.IRISH Mar/hCINQUEFOIL. 3. Common CINQUEFOIL. 4. Smallsrough CINQUEFOIL.
RAY. 611. 1.2. 90. ed. 140. RAY. Syn. ed. 12. 2. 141. 2. RAY611. 2. S. 90. ed. 141. 3. RAY612. 7. Syn. 90. 4. ed. 141. 5.



5. Tormentill CINQUEFOIL. 6.Upright WELH CINQUEFOIL. 7. SCOTCH CINQUEFOIL. 8. Shrub CINQUEFOIL. RAY 613.14. S.90.3.ed.141.4 RAY 616.1. Syn.91.1.ed.141.1. RAY Syn.91.4.ed.142.4 RAY 616.5. S.91.2.ed.1423.



9.TORMENTIL SEPTFOIL. 10PLOT STORMENTIL .11. SILVERweed wild TANSEY. 12. MOS CAT ELL.
RAY 617. 1. Syn. 91. ed. 14. 2. RAY 610. 3. S. 91. 2. ed. 14. 2. 2 RAY 617. c. 16. Syn. 91. 3. ed. 14. 2. 3 RAY 684 c. 27. S. 98. ed. 15. 1.





ENGLISH HERBS with 5 leaved FLOWERS & Star Heads. TAB. XLII.


LORPINE or Live long. 2. ROSE-ROOT. 3. NAVEL-WORT 4. HOUSELEEK. RAY.690.7. S. 99. ed. 152. RAY. 1878 . 1. Syn. 99. ed. 153. RAY.687.



5. Comon Yellow SENGREEN . 6. Grooked Yellow SENGREEN 7. PurpleMar In SENGREEN . O. Brigto LSENGREEN.



9STONEGROP or Wall Pepper, 10 Mild White STONE CROP. 11. Green BEAR S. FOOT. 12. SETTER-WORT. RAV104112. S.150.3. ed. 15 3.5 RAV1041.14. S.150. 4 ed. 15 3.6 RAV. 697. 2. S. 100. 1. ed. 15 3.1 RAV 698. 3. S. 100. 2. ed. 15 3. 2.




.

MUSEUM A. MOSEUM A. MOSEUM


8

ENGLISH PLANTS with STAR & ROUGH HEAD S,. TAB. XLIII.


1. MaJh Marigold. 2. Globe CROW-FOOT. 3. Water GLADIOLE. 4. COLUMBINES.



5. STARWater PLANTAIN. 6. Broad water PLANTAIN. 7 Narrow Water PLANTAIN & Smallwater PLANTAIN.




9. Grafs ARROW-HEAD. 10. Broad ARROWHEAD. 11. Marrow ARROW-HEAD. 12. Small ARROW-HEAD.
RAY Syn. 92. cd. 143. RAY 619. 1. S. 92 cd. 143. RAY 619. 3. Syn. 92. cd. 143. RAY Syn. edit. 2. p. 143.





ENGLISH PLANT S bearing BERRIES.


White BRIONY 2.Black BRIONY 3.Herb CHRISTOPHER 4. Butchers -Broom



5.SOLOMONS SEAL 6.SweetSOLOMONS-SEAE, 7LILY-CONVALLY &HERBPARIS.ONEBERRY RAY 664.15.96.1.ed.147.1. RAY 665.3.5.96.2.ed.148.2. RAY 667.1.5.96.1.ed.148.1.RAY 670.1.5vn.96.ed.148.



BITTER-SWEET 10. Common NIGHT SHADE 11 DWALE. Deadly Night/hade. 12. Wild SPARAGUS
RAY 672.1. S. 96.1. ed. 149.1 RAY 672.2. Syn. 97.2. ed. 149.2. RAY 679.1 Syn. 97. ed. 150. RAY 683.1 S. 97. ed. 150.






.

ENGLISH HERBS with 4 Leaved FLOWERS & Long Tods TAB XIV.

1.JACK by the HED GE 2 TREACLE WORMSEED. 3.WALL FLOWER, or STOCK 4. ScaSTOCK GYLLY FLOWER. RAY 792.c. 9.S. 109. ed. 165 RAY 811. 2. Syn. 113. 1. ed. 170.1 RAY 782. 1. Syn. 108. ed. 164 RAY 780. 3. S. 108. ed. 164.



5.WhiteThorow COLEWORT 6. Sea COLEWORT. 7. RoundTURNEP. 8. Long TURNEP. RAY797. 20. RAY796.13.S.109.ed166RAY 800.1.S.110.ed.167.RAY800.2.S.110.2.ed167.2



9.Wild NAVEW. 10.White MUSTARD. 11. Comon MUSTARD. 12.Wild MUSTARD. RAY802.2.S.110.ed.167.RAY802.2.S.111.3.ed.168.3.RAY803.3.S.111.ed.167.RAY802.1.Syn.111.ed.168.2.






ENGLISH PLANTS with 4Leaved FLOWERS & Long Pods. TAB. XIVI.

1. Winter CRESS. 2 Early Winter CRESS. 3 Hedge MUSTARD. 4. Broad Hedge MUSTARD.


RAY 809.8. Syn. 11. 2. 2. ed. 169.2. RAY 809.8. RAY 810.9. S. 112.3. ed. 169.3. RAY 811.3. S. 113.2. ed. 170.2.



5. Water ROCKET. 6 Sea ROCKET. 7 Man ROCKET. 8 Wall ROCKET.
RAY 808.3. S. 112.1. ed. 169.1 RAY 840. c.14. S. 118. ed. 176. RAY 809.5. S. 112.2. ed. 169.2. RAY 807.4 S. 112.1. ed. 169.1.




9. Garden RADISH 10. White CHARLOCK. 11. Great White CHARLOCK. 12. FLIX-WEED.
RAY804.1.S.1111.ed168.1. RAY805.2.S.111.2.ed168.2. RAY806.4. RAY812.10.S113.3.ed170.3.






ENGLISH HERB Swith 4 Leaved FLOWERS & flender Pods. TAB. XIMI.


1.Bitter CRESS. 2.Water CRESS. 3.Early Water CRESS. 4. Hairy CRESS.
RAY 814.1.Syn.114.1.edi71.1.RAY 816.6.S.115.6.edi72.0. RAY.3.413.1.S.115.7.edi72.7.RAY 815.5.S.114.4.edi71.4.



5.Lady Smock or CRESS. 6DoubleLady CRESS. 7.Impatient CRESS. 8.Trofoil CRESS. RAY814.2.S.114.2.cd171.2. RAY817.12. RAY815.4S.114.3.cd171.3. RAY817.11.



9 Daisy CRESS. 10. Tower MUSTARD. 11. Great Tower MUSTARD. 12 Daisy Tower MUSTARD. RAY 817.10. S. 114. 5. ed. 17 2.5. RAY 799. 3. S. 110. 1. ed. 166. RAY 799. 5. RAY 799. 6. S. 110. 2. ed. 166.





ENGLISH HERBS with 4 Leaved FLOWERS, Short Pods & Husks. TAB. XIVID


1. Thale CRESS. 2. Broad Thale CRESS. 3. Wreathed CRESS. 4 Small wreath CRESS. RAY 800.7. Syn. 110. ed. 166. RAY Syn. 241. ed. 166. 2. RAY 789. 6. Syn. 109. ed. 164. RAY Syn. 109. 2 edit 2. p. 165. 2.



5. Speedwell CRESS. 6. Whitlow CRESS.
RAY 790.4. Syn. 109.3. ed. 165.3. RAY 789.1. S. 109.1. ed. 165.

7.NotchedWhitlow CRESS.8IrijhRujh CRESS.

RAY Syn. ed. 2. p. 281.




g. WOAD. w. PEPPR-WORT.DITTANDER. u. GOLD of Pleasure. 12. Sea Grape COLEWORT.
RAY 842. Syn. 118. ed. 176. RAY 828. 1. Syn. 117. ed. 174. RAY 820. 1. S. 115. ed. 176.






ENGLISH HERBS with 4 leaved FLOWERS & Short Cods. TAB.XLX

1. Garden SCURVY-GRASS. 2. Sea SCURVY-GRASS. 3. Jvy SCURVY-GRASS. 4. SHEAPERDSPURSE. RAY822.1. S.11g. ed. 173.1. ... RAY823.2. Syn. 116. ed. 173.2. RAY824.6. Syn. ed. 173.4. RAY838.4. S.118.6. ed. 176.7.



5.RaggedSHEAPERDS-PURSE. 6.SmallSHEAPERDS-PURSE. 7.WholeSHEAPERDS-PURSE. 8.Broad WATER RADISH.
RAY819.3.S.115.ed.17.2.1.



9. Small Jagged WATERRADISH. 10. Great Jagged WATERRADISH. 11. Broad HORSE-RADISH. 12. Jagged HORSE-RADISH. RAY 818.2.

RAY 818.2.


RAY 818.2. cd.72.2.






ENGLISH HERBS with 4 Leaved Flowers & round flat Cods. TAB. L.


1. Narrow Wild CRESS. 2. Shepherds CRESS. 3. WELLH Rock CRESS. 4. Fine CRESS. RAY 825. 2. Syn. 116. 1. ed. 1741. RAY 827. 9. S. 116. 2. ed. 174. 2. RAY 3. p. 413. 3. S. 17. 4 ed. 1744. RAY. S. 236. ed. 174. 5.




g. Valencia CRESS. 6. Wart or Swines - CRESS. 7. Common Cow CRESS. 8. Smooth Cow CRESS. RAY 826.8. Sur. 3. edi74.3. RAY 843. c.20. Syn. 119. edi77. RAY 830.1. S. 117.1. ed. 175.1. RAY Syn. 11 8.5. edi75. 5.



9. Penny CRESS. 10 Hoary CRESS. 11. Wood CRESS. 12. Small thorow CRESS. RAY 8 31.5. S. 117. 3. ed. 175. 3. RAY 8 31. 2. S. 117. 2. ed. 175. 2. RAY 8 32. 10. S. 117. 4. ed. 175. 4. RAY 8 32. 8. S. 2 3 6. ed. 176. 6.






ENGLISH HERBS with 4 Leaved FLOWERS & small Cordated Pods. TAB. 51.


1. Broad vpright SPEEDWEL. 2. Comon Male SPEEDWEL. 3. Meadow SPEEDWEL. 4. Stalkt. SPEEDWEL.

RAY845. 1. Syn. 119. 1. ed. 17. 7.1. RAY851.7. S. 121. 13. ed. 179. 13. RAY850. 3. S. 121. 11. ed. 179. 11. RAY850. 4. S. 121. 12. ed. 179. 12.





5. Ltaft spiked SPEEDWEL 6. Smooth SPEEDWEL 7. Garden SPEEDWEL. 8. Wall SPEEDWEL. 9. Ruc SPEEDWEL.

RAY 846.3. Sug. 2. ed 177.2. RAY 848.6.S 119.3 ed 178.3 RAY 847.3. Sug. 4. ed 178.4. RAY 847.3: S1205 ed 178.5 RAY 848.5: S120.6. ed 178.6.




10. Jvy SPEEDWEL. 11. ComonBROOKLIME. 12. Long BROOKLIME. 13. NavrowBROOKLIME.
RAY848. 4. S. 120.7. ed. 178.7. RAY852.1. S. 120.8. ed. 178.8. RAY852.4. S. 120.9. ed. 179.9. RAY852.5. S. 120. 10. ed. 170. 10.





ENGLISH Plants with Four-leaved FLOWERS. TAB. LII.


1. Wild POPPY. 2. Corn POPPY 3. Long POPPY. 4. Yellow POPPY.
RAY 854. 3. S. 121. 1. edi80. 1. RAY 855. 4. S. 121. 2. ed. 180. 2. RAY 850. 7. S. 122. 5. ed. 180. 5. RAY 850. 0. S. 122. 6. ed. 180. 6.



Rough round POPPY. 6. Rough long POPPY. 7. Sellow HORY POPPY. 8 Purple HORN-POPPY. RAY855.5.5.122.4. ed. 180.4. RAY856.6.5.122.3. ed. 180.3. RAY857.12.5.122.7. ed. 180.7. RAY857.14.5.122.8. ed. 181.8.



9. CELANDINE. 10.Rose WILLOW-HERB. 11. Hairy great WILLOW-HERB. 12. Hairy Small WILLOW-HERB. R. 858.15. Syn.122.9. editing. R. 860.15.123.1. ed. 181.1. RAY 861.4. Syn.123.2. ed. 182.2. RAY 861.5. S. 123.3. ed. 182.3.





ENGLISH HERBS with Fore-leaved FLOWERS. TAB. LIII

1. Oreat Smooth WILLOW HERB. 2. Ray WILLOW HERB. 3 Midle Smooth WILLOW HERB. 4. Small Smooth WILLOW HERB. R4Y 861.6. S. 123.4. ed. 182.4. R. 862.9. Syn. 124 ed. 182.7. R4Y 861.7. Syn. 123.5. ed. 182.5. R4Y 862. 8. Syn. 124. 6. ed. 182. 6.




5. Wood SPURGE. 6. Warty SPURGE.



7. Broad SPURGE.



8. Sca SPURGE RAY. 805.10.5.124. 2. cd.184. 4



9. Corn SPURGE. RAY868.23.S.125.5.ed.184.6.R.869.26.S.125.7.ed.184.5. RAY869.27.S.125.8.ed.184.8.R.869.2.S.125




10. Sun SPURGE.



n.Round SPURGE.













FNGLISH HERBS with BELL-FLOWERS. TABLIN.

1. Sreat BINDWEED. 2. Small BINDWEED. 3. Navrow BINDWEED. 4. Sea BINDWEED. RAY 725. 6. S102. 1. ed. 157. 1. RAY 725. 7. S. 10 2. 2. ed. 157. 2. R. 725. 8. Syn. 10 2. 3. ed. 157. 3. R. 726. 12. S. 10 2. 4. ed. 157. 4.




5. RAMPIONS. 6. Round BEILFLOWER. 7. Tall round BEILFLOWER. 8. by BEILFLOWER. RAY 739. u.S. 103.4. ed 158.1. R.740. 12. S. 103. 5. ed 158.2. R.741.17. S. 103.6. ed 159.3.




9. Giant THROAT-WORT. 10. Small THROAT-WORT. 11. Great THROAT-WORT. 12. Small VENUS LOCKING -GLASS.
[RAY 732. L.S. 103. 1. ed. 158. 1. R.735. 18. S. 103. 3. ed. 158. 3. R.732. 4. S. 103. 2. ed. 158. 2. R.743. 4. S. 104. 7. ed. 159.






ENGLISHHERBS with g Leaved hollow FLOWERS. TAB. IV.


Horned RAMPIONS. 2. Scabious RAMPIONS. 3 Marsh GENTIAN. 4 Dwarf GENTIAN. RAY744. 2. Syn. 104. 9. edit 9. 1. R. 744. 2. S. 104. 10. edit 9. 2. R. 710. 6. S. 101. 1. edit 9. 1. R. 719. 7. S. 102. 2. edit 9. 2.




5. BUCK-BEANS. o. Greek VALERIAN. 7. Small CENTURY. 8. Sellow CENTORY. R. 1099. c. 14. Syn. 161. ed. 226. R. 1102. 20. S. 163. ed. 228. R. 1092. 1. Syn. 159. ed. 224. R. 1093. c. 10. S. 160. ed. 224.




9. Manured FLAX. 10. Wild FLAX. 11. Narrow FLAX. 12. Turging FLAX. R. 107 2.1. S. 157. ed. 220. R. 107 3. 2. S. 157. 2. ed. 220. 2. R. 107 4.13. S. 157. 4. ed. 221. 5. R. 107 5. 21 S. 157. 5. ed. 221. 6.






ENGLISHPLANT Swith 5 Leaved FLOWERS. TAB. 50.

1. Wall PINK. 2. Wild PINK. 3. Single PINK. 4. Moss PINK. R. 988. g. S. 138. 2. ed. 199. 2. RAY 988. 6. S. 138. 1. ed. 198. 1. R. Syn. 242. ed. 199. R. 1504. 11. S. 141. 6. ed. 202. 19



5. Deptford PINK. 6. Lagged PINK. 7. Great PERIWINCKIE. Small PERIWINCKIE. R. 991.4. S. 139.3. ed. 199.3. R. 1000.6. S. 139.3. ed. 200.3 R. 1091.2 S. 242. ed. 223. 2. R. 1091.1 S. 159. ed. 223.




9. Yellow PRIMROSE. 10. OX-SLIPS. 11. COW-SLIPS or PAGILS. 12. BIRDS-EYN. R. 1080. 1. S. 138. 1. ed. 222. 1. R 1081. 4. S. 138. 2. ed. 222. 2. R 1081. 5. 138. 3. ed. 222. 3. R 1084. 9. S. 139. ed. 223.




A. M.

ENGLISH Plants with 5 leaved Flowers ._ TAB. 57.


1. Sea CAMPION. 2.Bladder CAMPION. 3. SOPE-WORT. 4. Hollow SOPE-WORT.
RAY 998.3. S. 139. 1. ed. 199.1. R. 998.1. Syn. 139. 2. ed. 199.2. R. 999. 2. S. 140.6. ed. 200.6.
RAY 998.3. S. 139. 1. ed. 199.1. R. 998.1. Syn. 139. 2. ed. 199.2. S. 140.6. ed. 200.6.




g. White CAMPION. 6.Red CAMPION. 7BroadWhite CATCHFLY. 8 Red German CATCHFLY. R.994.2.5.140.8.ed.200.7. R.994.3.5.140.9.ed.200.8. R.1001.5.5.140.n.ed.201.10. R1000.1.5.141.4.ed.201.13.




9 DENTS Small CAMPION. 10. Night CAMPION. 11. Spanish CATCH-FLY. 12. COCKLE.
R.996.13.S.140.10.cd. 201.9. R.994.5. S.141.13.cd. 201.12. R.1002.9.S.141.15.cd. 201.14. R.998.1.S.139.4.cd. 200.4.





ENGLISH PLANTS with & Leaved, Forked FLOWERS. TAB. 58.


1. Great STITCH-WORT 2. Middle STITCH-WORT . 3. Lefser STITCH-WORT . 4. Bogg STITCH-WORT. Rozo 201. 2. R. 1030. 21. S. 147. 8. ed. 209. 10.



5Broad Huiry STITCHWORT. 6 Narrow Hairy STITCHWORT-Buddles Hairy STITCH-WORT & Mouse car STITCH-WORT.

R.10 29.14. S.146. 3. cd. 208. 5. R.10 29.15. S.146. 4. cd. 208. 6. R.

R.10 29.14. S.146. 3. cd. 208. 5. R.10 29.15. S.146. 4. cd. 208. 6. R.




9. Snowdon STITCHWORT. 10 Welfh STITCHWORT. 11. Marsh CHICKWFFD. 12. Comon CHICK-WFFD. R.Syn. 147. 5. ed. 208.7. RAY. S. 147. ed. 209. 11. R. 1030.19. S. 147. 6. ed. 208.8. R. 1030. 20. S. 147. 7. ed. 209. 9.






ENGLISH Chick-weeds, Spurreys Pearl-worts &c. TAB. 59.


1. Plantain CHICKWEED. 2. Wall CHICKWEED. 3. Grafs CHICKWEED. 4. Mount CHICKWEED. R. 1032.5. S. 148.2. ed. 209. 2. R. 1031.1. S. 147.1. ed. 209.1. R. 1031.4. S. 198.3. ed. 209.3. R. 1033. x. 1. S. 148.4. ed. 209.4.



5 Sand CHICKWEED. 6. Corn SPURRY. 7. Sea SPURRY. 8. Purple SPURRY. R. 1032.9.8.148.5.ed 210.5. R. 1034.19.8.148.6.ed 210.8. R. 1034.20.8.148.8.ed 210.10. R. 1034.18.8.148.7. ed. 210.9.




9. RAY Stitch-wort. 10. PEARL-WORT. 21. Plots PEARL-WORT. 12. Leaft RUPTURE-WORT R. 1025. 1. S. 145. 1. ed. 206. 1. R. 1026. 5. S. 145. 2. ed. 206. 2. R. 1026. 6. S. 145. 3. ed. 2073. R. 10267. S. 146. 5. ed. 207. 5.






ENGLISH PLANTS with 5 Leaved FLOWERS. TAB. 60.


1. Comon DWARF Ciftus. 2. Kendal DWARF Ciftus. 3. Brent DWARF Ciftus. 4 Jersey DWARF Ciftus. RAY 10.13.1. S.142.1.ed. 202.1. R.1015.7. Syn.142.2.ed. 203.2. RAY. Syn.143.3.ed. 203.3. R.1012.2.S. 238. ed. 203.



5. Comon S. IOHNS-WORT. 6. Upright S IOHNS-WORT. 7 Evay-blade S. IOHN WORT. 8. Ground S. IOHNS-WORT. R. 1018. 5. S. 143. 1. cd. 203 1. R. 1019. 11. S. 143. 2. cd. 203. 2. R. 1021. 19. S. 143. 5. cd. 204. 5. R. 1019. 10. S. 143. 3. cd. 205. 3.




9 TUTSANor PARKleaves. 10 Titt an S. TOHNS-WORT. 11 S. PETERS-WORT. 12 Hoavy S. PETERS-WORT. R. 10 20.16. S. 143.6.cd 204.6 R 10 20.17. S. 143.4.cd 204.4 R 10 19.12. S. 144.7. cd 204.7. R 10 20.18. S. 144.8. cd. 205.8.





ENGLISH Plants with 5 leaved Flowers . _ TAB. 61.


1. Garden PRIDE. 2. Round PRIDE. 3. Iryh PRIDE. 4. Welsh PRIDE. RAY. Hist. Plant 1046. 6. RAY. Pl. 1047. 1. RAY 1045. 3. R. 3. P. 507. 2. S. 151. 2. ed. 213. 2.



5. Hairy PRIDE. 6. White PRIDE. 7. Trifid PRIDE. 8. Rue PRIDE.
R. 1046. 8. S. 151.1. ed. 213.1. R. 1048. S. 151.3. ed. 213.3. R. 1043.21. S. 151.6. ed. 213.4. R. 1043.25. S. 151.7 ed. 213.5.



9 PURPLE Heath SENGREEN. x.RAYS Yellow SENGREEN. x.WhiteHousleek. 12 Parnajs Grass
R. 1940.6.S. 150.2.ed. 212.2. R. 1042.20.S. 150.5.ed. 212.3. R. 1040.1.S. 150.1.ed. 212.1.R. 1049.c. 4.S. 151.ed. 213.







1. Comon MULLEIN 2. Small MULLEIN 3. Sage MULLEIN 4. White MULLEIN RAY 1094. 1. S. 160. 1. ed. 225. 1. R. 1094. 3. S. 160. 2. ed. 225. 2. R. 1095. 7. S. 160. 4. ed. 225. 4. R. 1095. 5. S. 160. 3. ed. 225. 3.




¿Yellow MOTH-MULLEIN 6. Spiked LOOSE-STRIFE 7. Short LOOSE-STRIFE 8. Trefoil LOOSE-STRIFE RAY Hijt. Plant. p. 1096. 1. R. 1036. 1. S. 149. ed. 212.








ENGLISH Plants with 5 leaved FLOWERS. TAB. LXIII.






ENGLISH Cranebills, with 5 leaved FLOWERS.





ENGLISH Cranesbills with 5 leaved Flowers &c. TAB. 65.






ENGLISH Herbs with Bulbous Roots. TAB. 66.

2. Flowring Crow GARLICK. 3. Twinheaded GARLICK. 4. Raven GARLICK. 1. Crow GARLICK 7. Holm GARLICK. 8 RAM SONS. 6. Purple LEEKS. 6.CornGARLICK. RAY Syn. 165.6.ed. 230.6. RAY 1119.3. Syn. 164.3. RAY 1126.2. Syn. 165.4. ed. RAY 112.12. Syn. 165.5. 9 Laucashire ASPHODEL. 10 Green spiked Bethlem Star. 11. Common Bethlem Star. 12. Yellow Bethlem Star. RAY 1153.9 Syn. 166.2.ed. RAY 1153.13. Syn. 166.3.ed. 231.3. Syn.169.1. RAY115.11. Syn.166.1.ed.




ENGLISH Herbs with Bullous Roots.


4.English HYACINTH. 3.True SAFFRON. 1. Lewyd's Snowdon BULB. 2. Meadow SAFFRON. RAY1170.1.Syn.167.ed.232.RAY1176.17.Syn.167.ed2,32.RAY1159.20.Syn.167.2.ed. RAYSyn.160.ed.233.



8.Hornsey DAFFODIL 6 Autumn Star HYACINTH. 7 Lake HYACINTH. 5. Star HYACINTH.



11. Yellow Flower de -luce. 10.Primrose DAFFODIL. 12 Sanking Gladdon. 9. English DAFFODIL. RAY 1131. 5. Syn. 166. 1. ed 230. 1. RAY 1135. 23. Syn 166. 2. ed. RAY 1186. 24. Syn. 168. 1. ed. RAY 1190. 7. Syn. 168. 2. ed. 234. 3.





ENGLISH Orchis'es with double or single Roots. T.13.68.

1. Goat-Stones. 2. Spotted Male Pool-stones. 3 Female Fool-stones. 40val Inne Orchis
RAY 1212.1. Syn. 169. 1. ed. RAY 1214.7. Syn. 169. 2 ed. RAY 1214. 8 Syn. 170. 3. ed. RAY 1215.14. Syn. 170. 5. ed. 236. 3. 236. 4. 236. 5. 235.1. put Specks Tune (Chalkey ground Fl. mid Hay Purple speckt Comon Sin Tujures Fl. Iune 5. Little Purple Orchis. 6. Sm. all Yellow Musk Orchis. 7. Triple Ladies - traces. 8Ris as Man-Orchis.

RAY 1215.12 Syn. 170. 4. ed. RAY 1217. 22 Syn. 170. 6. ed. RAY 1217. 26. Syn. 171. 7. ed. RAY 1217. 27 Syn. 171. ed.

237. 6. 237. 7. 237. 9. BRusa Man-Orchis. I'L small, blujh speckt. On barren Chalky hills! dry, & Fl.yellon Iuly. In barret ground. White Aug tsc. Chalkey drži ground Wfields in inwelpitts. 12 Leser Butterfly-Orchis. "Butterfly-Orchis. 10 Common Fly-Orchis. 1718. ed. RAY1219.32. Syn.171.10.ed. RAY1221.40.Syn.172.13.ed.RAY1221.Syn.edit. 238.14. blackish FL. Iune Thickets K Hilly




ENGLISH Orchis'es with double, or handed Roots. & Rapes. 7413.

2 Humble Bee Orchis . 3 Green-wing'd Humble-Bee. 4 Sweet White Orchi 1 Great Hly-Orchis. RAY 1220.33 Syn: ed. RAY 1220.38 Syn.17 1.11.RAY 1221.39 Syn.172.12.
238.1. ed.238.12 Gravil pits & Pastures Tune · dry soil. In some Patures. May 5 Male Hand-Orchis 6 Spotted Hand-Orchis. 7 Red spurd, Hand-Orchis. 8 Frog Hand-Orchis RAY 1223.1 Syn. 172.14. RAY 1223. 2 Syn. 172.15. RAY 1224.6 Syn. 173.16. RAY 224.7 Syn. 173.17. ed 239.15. ed 239.16. Purplespeckt Fl. Purple May Kc. Wet boggs Meadows Tune Woods & Meads. Iun. Wet Me ads Fastu res. Fl. Green vedon Tune Dry Pasture 9 Zeeland Dwarf Orchis. 10 Broom-Rape . 11 Branched]
RAY 1226. 3 Syn. 173. ed. RAY 1227.1. Syn. 173. 1. ed. RAY 1227. 5
240. 12 Plotts Rape 11 Branched Rape. RAY 1229.2. Syn. 174. 2 Fl.Rusty Tun. &c. Broom field Heath & lone. Strair c. Iune &c Shady dark Woods FlBurplish In Corn & Flux Fl. Greenish
Tune
Bogs about
Cambridge




•


•





5 Blackish Helleborine. 6 Greenish Helleborine. 7 White Helleborine. 8 March Helleborine. 5.3. ed.RAS 1231.10 Ngn.175.5. ed.



oLong, nointed Helleborine . 10 Common Tway blade . 11 Least Twablade . 12 Marsh Tway blade. n 1751.ed.RAY 1232. 218yn.176.2.ed.RAY 12324 Syn.176.3.ed.









TAB. 72. ENGLISH Miscellany Plants.

1. Water Valerian. 2. Marsh Valerian. 3. Small floured Marsh Valerian. 4. Corn-Sallet.

RAY 388.2. Syn. 60. RAY 388.3. Syn. 61.2. RAY 389.4. Syn. 61.3. ed. 98.3. RAY 392.1. Syn. 61.

ed. 98.2. RAY 39 2. 2. Syn. 61. 2. RAY 39 5. LSyn. 61.ed. RAY 39 5. Syn. 61. ed 99. RAY 1037.1. Syn. 149. ed 98. 2. Salt Mars hes. 1 x Mountains Fl. Purp. Iun.Y. Amongst (Tl.Bhush 9 Branched Bur-Reed. 10. Unbranched Bur-Reed. 11. Least Bur-Reed. 5.1. RAY 1311. 2. Syn. 205. 2. RAY 1312.3. ed 277.3.

on Heath's H. Blush Iun.Y.



PLANTS already Engraved in Mr. Petiver's ENGLISH HERBAL.

Herbs with Stamineous, Threddy or Blink Flowers.

SECT. I.

Blink Flowers.

SECT. I.

Male Hops Ray Hift. Plant. 156. Syn. 52. T. I. F. I.
Female Hops. Ray Hift. & Syn. id. Tab. 1. Fig. 2.
Male Homp. Ray H. Pl. 158. 1. Syn. 53. Tab. 1. Fig. 3.
Female Homp. Ray id. Tab. 1. Fig. 4:
Male Dog's Mercury R. 163. 1. Syn. 53. I. T. 1. 5.
Female Dog's Mercury R. 163. 1. Syn. 53. I. T. 1. 5.
Female Prench Mercury R. 163. 2. S. 54. T. 1. 7.
Female French Mercury R. 163. 2. S. 54. T. 1. 7.
Female French Mercury R. 163. 2. S. 54. T. 1. 8.
Great Nettle Ray 160. 4. S. 54. T. 1. 10.
Roman Nettle Ray 161. 5. S. 54. T. 1. 10.
Roman Nettle Ray 161. 5. S. 54. T. 1. 10.
Roman Nettle Ray 165. 1. S. 55. T. 1. 12.
Great Water Dock Ray 177. 4. Syn. 55. 1. Tab. 2. 1.
Great Water Dock Ray 177. 5. Syn. 56. 4. T. 2. 3.
Small Burdock Ray. 174. Syn. 56. 4. T. 2. 3.
Small fharp Dock Ray 174. Syn. 56. T. 2. 5.
Bloodles Dock Tab. 2. Fig. 6.
Gold Dock Ray 174. Syn. 56. 6. Tab. 2. 4.
Blood vein'd Dock Ray 174. 8. Syn. 56. 6. Tab. 2. 8.
Broad Dock Ray 174. 8. Syn. 56. 5. Tab. 2. 11.
Trailing Buck-wheat Ray 182. 1. Syn. 57. 1. Tab. 2. 11.
Trailing Buck-wheat Ray 182. 1. Syn. 56. 8. Tab. 2. 12.
Common Sorrel Ray 178. 1. Syn. 56. 8. Tab. 3. 1.
Sheep's Sorrel Ray 186. 14. Syn. 56. 9.
Barren Sheep's Sorrel T. 3. 3.
Welfh Sorrel Ray 181. 3. Syn. 58. 2. Tab. 3. 6.
Spotted Arjmart Ray 182. 2. Syn. 58. 3. Tab. 3. 6.
Spotted Arjmart Ray 183. 3. Syn. 58. 2. Tab. 3. 6.
Spotted Arjmart Ray 183. 3. Syn. 58. 2. Tab. 3. 6.
Spotted Arjmart Ray 184. 6. Syn. 58. 4. Tab. 4. 1.
Deuted Plantain Ray 87. 5. Syn. 184. 2. Tab. 4. 5.
Common Plantain Ray 87. 5. Syn. 185. 4. Tab. 4. 1.
Deuted Plantain Ray 87. 5. Syn. 185. 5. Tab. 4. 5.
Common Ribwort Ray 877. 5. Syn. 185. 5. Tab. 4. 6.
Mountain Ray 880. 5. Syn. 185. 5. Tab. 4. 6.
Mountain Ray 880. 5. Syn. 185. 5. Tab. 4. 6.
Mountain Ray 880. 5. Syn. 185. 5. Tab. 4. 6.
Mountain Ray 880. 5. Syn. 185. 5. Tab. 4. 6.
Mountain Ray 880. 5. Syn. 185. 5. Tab. 5. 6.
Event Pondweed Ray 188. 1. Syn. 60. 2. Tab. 5. 5.
Round Pondweed Ray 188. 1. Syn. 60. 1. Tab. 5. 4.
Long Pondweed Ray 188. 1. Syn.

Narrow Grass Pondweed Tab. 5. Fig. 11.
Fine Grass Pondweed Ray 189. 7. Syn. 61. 7. Tab. 5. 12.
Fennel Pondweed Tab. 5. Fig. 13.
Tastel Pondweed Ray 190. 10. Syn. 62. 10. Tab. 6. 1.
Horned Pondweed Ray 190. 13. Syn. 281. Tab. 6. 2.
Water Starwort Ray 1323. Syn. 280. 1. Tab. 6. 3.
Long Water Starwort Ray Syn. 280. 2. Tab. 6. 4.
Feather Pondweed Ray 191. Syn. 279. Tab. 6. 5.
Fine Feather Pondweed Tab. 6. Fig. 6.
Millfoil Pondwed Ray. 191. 2. Syn. 280. 5. Tab. 6. 7.
Knotted Pondweed Ray 1322. 1. Syn. 280. 5. Tab. 6. 7.
Golden Saxifrage Ray 207. 1. Syn. 66. Tab. 6. 8.
Golden Saxifrage Ray 207. 2. Syn. 66. Tab. 6. 10.
Cornwal Pennywort Ray. Syn. 211. 14. Tab. 6. 11.
Marsh Pennywort Ray 1323. Syn. 280. Tab. 6. 12.
Delt Orrach Ray 192. 3. Syn. 62. 1. Tab. 7. 1.
Delt Sea Orrach Ray 193. 10. Syn. 63. 4. Tab. 7. 2.
Frosty Sea Orrach Ray 193. 8. Syn. 63. 5. Tab. 7. 3.
Jagged long Orrach Ray 192. 6. S. 62. 3. Tab. 7. 4.
Spear Orrach Ray 195. 3. Syn. 63. 8. Tab. 7. 7.
Heart Sea Purstain Ray 196. 2. Syn. 65. 11. Tab. 7. 10.
Stinking Orrach Ray 196. 2. Syn. 64. 6. Tab. 7. 11.
Wild Spinage Ray 198. 11. Syn. 64. 6. Tab. 7. 11.
Wild Spinage Ray 198. 11. Syn. 64. 6. Tab. 7. 11.
Wild Spinage Ray 198. 1. Syn. 64. 8. Tab. 7. 12,
Rands Oak Blite Tab. 8. 1.
Froft Blite Ray 198. 8. Tab. 8. 2.
Buddle's Fig Blite Tab. 8. 3. Stinking Orrach Ray 198. 11. Syn. 64. 6. 1ab. 7. 11. Wild Spinage Ray 195. 1. Syn. 64. 8. Tab. 7. 12, Rands Oak Blite Tab. 8. 1. Froth Blite Ray 198. 8. Tab. 8. 2. Buddle's Fig Blite Tab. 8. 3. Buddle's round Blite Tab. 8. 4. Tbick finning Blite Ray Syn. 64. 5. Tab. 8. 5. Sharp pointed Blite Ray 197. 9. Syn. 64. 3. Tab. 8. 6. Maple Blite Ray 8. 7, Broad pointed Blite Tab. 8. 8. Sea Beet Ray 204. 3. Syn. 65. Tab. 8. 9. Wall Pellitory Ray 206. 1. Syn. 66. Tab. 8. 10. Tree Stonecrop Ray 199. 16. Syn. 65. 9. Tab. 8. 11. Shrub Stonecrop Ray 199. 16. Syn. 65. 9. Tab. 9. Fig. 1. Prickly Glasswort Ray 198. 14. Syn. 64. 7. Tab. 9. Fig. 1. Prickly Glasswort Ray 198. 14. Syn. 67. 1. Tab. 9. Fig. 3. Woody jointed Glasswort Ray 212. 2. Syn. 67. 1. Tab. 9. Fig. 3. Woody jointed Glasswort Ray 211. 1. Syn.67. 2. Tab. 9. 4. Baftard Toad-flax Ray399. 7. Syn. 99. Tab. 9. Fig. 5. Upright Knawell Tab. 9. 6. Spreading Knawell Ray 213. 1. Syn.68. 1. Tab. 9. 7. Pepper Grass Ray 1325. 10. Syn. 281. Tab. 9. 8. Ladies Mantle Ray 208. 1. Syn. 66. Tab.9. 9. Small Ladies Mantle Ray 202. 2. Tab. 9. 10. Silver Ladies Mantle Ray 202. 2. Tab. 9. 10. Silver Ladies Mantle Ray 209. 4. Syn. 66. Tab.9. 11. Parsy Piert Ray 209. 14. Syn. 67. Tab. 9. 12. Common Knotgrass Tab. 10. 2. Short leaved Knotgrass Tab. 10. 3. Narrow Knotgrass Tab. 10. 2. Short leaved Knotgrass Tab. 10. 3. Narrow Knotgrass Ray 185. 2. Syn. 59. 2. Tab. 10. 5. Newton's Knotgrass Ray 214. 3. Syn. 68. Tab. 10. 7. Valentia Knotgrass Ray 215. 8. Tab. 10. 8. Smooth Rupture-wort Tab. 10. 10. Sea Heath Ray 1716. 15. Syn. 314. 5. Tab. 10. 11. Blink's Ray. 1035. 26. Syn. 211. 14. Tab. 10. 12.

N. B. The Figures of all these Plants done on Copper Plates are ready to be delivered upon paying 10 s. in Hand, and in May next, on depositing a Guinea they shall be made up 600. and 10 s. more, when they amount to One Thousand Trees and Herbs, all which shall be of the Growth of Great Britain and Ireland. Such Generous Patrons as will be pleased to give one Guinea in Hand, instead of 10 s. to forward this Work, shall have a TABLE Dedicated to them. Any Person subscribing for Five Books, shall have a Sixth Gratis.

€ ,

1

X.

.11

51,

. . . .

EL-07

Hortus PERUVIANUS Medicinalis:

OR,

The SOUTH-SEA Herbal.

Containing the Names, Figures, Use, &c. of divers Medicinal Plants, lately discovered by Pere L. Feuillèe, one of the King of France's Herbalists.

To which are ADDED,

The Figures, &c. of divers American Gum-Trees, Dying Woods, Drugs, as the Jesuits Bark-Tree and others, much desired and very necessary to be known by all such as now Traffick to the South-Seas, or reside in those Parts.

TAB. I. Medicinal PERU Plants.

I G. 1. Peru Chilca or Long Flezbane.

Conyza frut. fol. angust. nervosis L. Feuillée p. 750.

Conyza Afric. humilis, fol. angustioribus nervosis, Floribus umbellatis Inst. Rei Herb. 455.

It has a pleasant Smell, the Indians make a Tea of Grenothen their Stemach

it, to strengthen their Stomach.
2. Manga Paki or Round Peru Fleabane

Conyza fol. subrotundo utrinque acuto F. p. 749.

This is a sweetning Herb, Astringent and Vulnerary, the Natives drink a Tea of it in Cholick Pains, Dissenteries, and other Fluxes. Grows 4 Feet high with blewish Flowers, in the Valleys about Lima the Capital City of Peru.

3. Paica Jullo. Bidens Mercurialis folio, fl. radiato F.744. Its Flower-rim white. Grows about Lima and Callao. 4. Cuiez-bane. Bidens fol. trinervi lanceato, fl. singulari &

The Flower-rim composed of 6 yellow Petals. Grows in the floney Valleys of Lima. It's a purging Plant, but rarely used by reason of its Violence, they also think it a Poison, because it kills a House Animal call'd Cuiez in Peru and Chili.

5. White, Mugwort Bident. Bidens Arthemisia folio, st. albo

radiato F. 745.

Found in the Plains about Lima. 6. Heart-leaved Maracoc.

Granadilla pomifera, Tilia folio F. 720.

Its Rays crimion with a white Circle in the middle. Grows in the Vallies about Lima in divers Places, it's also nursed in their Gardens.

7. Trisid, spotted Maracoc. Granadilla fol. tricuspidi, obtuso & oculato F. 718.

Observed in a Garden at Malambo, in the Suburbs North

8. Caigua or striped Melon. Momordica fructu striato lævi.

The Natives eat its Finit in their Broths which much refreshes and cools them, the Heats there being very excessive. A. and B. are 2 small Beetles which feed on the Leaves of this Plant.

9. Lima Pumkin.

Melongena Laurifolia, fr. turbinato variegato F. 735. Its Flower white with a purple Star, the Fruit much eaten at Lima but apt to caule Fevers, which are there difficult to cure. C. its Fr. Slit.

10. Purple Bladder Nightsbade

Alkakengi amplo flore violaceo F 724.

They boil 3 or 4 of these Berries in White-wine or Water, and drink it with wonderful Success in Stoppage of Urine or Gravel.

11. Ancoacha or yellow Mallow.

Malva lutea, calyce simplici, obtuso Carpini solio, pediculis

florum piælongis F. 75c.

Its Fower yellow with a purple fringed bottom and cordated Petals. A Tea of this is wonderful in Alis of the Stomach, the Indians pick and make a Poultifs of it, which they apply to tipen Swellings and use it as an Universal Remedy. Grows in moist Ground and by River-sides North of Lima.

Capraria Peruviana Agerati foliis, absque pediculis F. 764.

Pere Feuillée says, the Virtues of this Heib (which are the fame with the China Tea) was not known in Peru till 1709. 12. Lima Tea.

N. B. This feems to be the same or little different from Dr. Sloan's West-India Tea, which he has described in his Nat. Hist. of Jamaica p. 209. pl. 25. who says it's call'd in Curassow Cabritta and Capraria because the Goats there seed on it, which last Name our Author here calls it by. J. Petiver.

TAB. II. Medicinal Peru and Chili Plants.

Fig. 1. Ima Tobacco.

Nicotiana minor, fol. cordi formi, tubo Floris pralongo F. 717. Its Flower of a greenish yellow. Grows in Valleys.

2. Ray leaved Bignon.
Bignonia fl. luteo, foliis radiatis, & elegantissime dissectis F.731.
Found in a very dry Place at the Foot of high Mountains about

3. Broad Clove-strife. Onagra Laurifolia fl. amplo pentapetalo. F. 716.
* Small Female Clove-strife.

* Small Female Clove-strife.

Onagra minor, fl. luteo pentapetalo.

The Flower yellow and L. like the last, but both smaller.

The Indians highly esteem these 2 Shrubs, making a Poultise of the L. pickt, which sweetens, mollisses and dissolves, sit therefore to dissipate inveterate Tumors and Buboes which are very common in those Parts. These grow by River sides.

4. Log rooted yellow Wood-sorrel.

Oxys luteo flore, rad. crassissima F. 734.

Grows on the Mountains of Peiu, &c.

5. Spotted Mountain Pelegrina.

Hemerocallis st. purpurascentibus maculatis F. 711.

Hemerocallis fl. purpurascentibus maculatis F. 711.

The Beauty of this Flower is, in having 3 of its Petals spotted, the rest plain, and therefore the Spaniards call'd it Pelegrina or Curious.

6. Ylo Hounds-tongue.

6. Ylo Hounds-tongue.

Cynoglossum fol. nervosis acutissimis F. 765.

Grows in the Valley of Ilo near Arica about a Foot and an half high, with blewish Flowers.

7. Malla or Peru Cress-Violet.

Cardamindum quinquishdo folio F. 756.

This elegant Plant the Spaniards call Paxarito because its Flower is composed of 2 particular large yellow L. extended like the Wings of a Bird. Grows about Lima.

8. Lima Flower fence or Tara.

8. Lima Flower fence or Tara.

8. Lima Flower fence or Laru.
Poinciana spinosa Feuillée. 572.

The Dyers use the Husks of this Fruit to dye black, they soke some of the Pods all Night in Water, mix a little Allum with it, and boil it to a due Thickness, which makes a very sine Black and strong Ink.

Medicinal Chili Plants.

9. Dafie Ragwort or Nillque.
Jacobea Leucanthemi vulg, folio F. 759.
Its Flowers yellow. They drink a Tea of this after the cold
Fit, which abates the hot in intermitting Fevers. Grows on

the ragged Sea Banks of Chili, &c.

10. Chili Duthoa or Floripondio.

Stramonoides arboreum oblongo & integro folio, frudulævi F.761.

Grows in Chili about a Yard high, with a white Flower for very fweet, that a fingle Plant is enough to perfume a whole Gaiden. They apply the L. to hasten the Suppuration of Tumors which it does effectually. N. B. This Plant bears the Flower of a Stramonium or Thorn apple, but differs in its Fruit being smooth, there is a fort of this in the Hort. Malabaricus Vol. 2. Tab. 22 11. Chili Nightsbade

Solanum Chenopodoides, acinis albescentibus F. 721.

The Natives were ignorant of the Virtue of this Plant until the Negroes came amongst them who were subject to a certain Disease which kill'd them in their Prime, it was an extraordinary Extension of the Anus attended with a Fever, which was so mortal, that many of them dy'd before they discover'd this Remedy. They juice the tops of this Plant, mixing it with Rose water and a little Allum, which taken cures their Ail, stops the Fever, and they become very well. With this suice they also bathe their Eyes, which takes away all Pains Juice they also bathe their Eyes, which takes away all Pains and Dimness in them. This Sovereign Plant grows about a Yard high, on the Mountains of Valparaiso, &c.

12. Oak-leaved Bittersweet.

Its Flower purple, found with

Solanum foliis Quercinis. F. 722. Its Flower purple, found with the lake, and grows near 2 Yards high.

TAB. III. Medicinal Chili Plants.

Fig. 1. Carlet Avens or Quellgon.

Caryophylata fol. alatis, fl. ample coccinee F. 726. les an Aperitive Herb, which the Natives make a Tea of to keep their Bodies in order. Grows about half a Yard high

Pentaphylloides Alcea minori folio, fl. purp. F. 763.

Grows about a Foot high, with a purple Flower, on the Banks of the River Plata.

3. Panama Centaury.

Centaurium minus purp. patulum F

Contain minus purp. patulum F. 747.

This Plant is very bitter. They make a Tea of it which is Aperitive and Sudorifick, it fortifies the Stomach, kills Worms, is Aperitive and Sudorifick, it fortifies the Stomach, kills Worms, cures intermitting Fevers and the Jaundice, it's also given with very good Succels in Rheumarisms, &c. They take it as hot as may be in Bed, covering themselves close, to provoke Sweat. N.B. The History of the Academy Royal des Sciences A.D. 1707. p. 65. 8°. gives great Encomiums to this Plant, which they say has a fragrant Smell like Natural Balsam, and is so great a Sweetner of the Blood, that it's a specifick in Pleurises, and in Fevers they use it instead of the Jesuits Bayk. Its sound plentifully about Panama and divers other Places, that which grows on the Mountains is esteem'd the best. J. P.

4. Upright Purple Wood sorrel.

4. Upright Purple Wood forrel. Oxys roleo flore erectior F. 733.

They make a Tifan or Broath of this Herb, which is opening its L. are tart and refreshing, they abate Choler and cool the Heat of the Blood. Grows on the vast Plains along the Ri. ver Plata.

116. Light or Purple striped Day-Lilly.

Hemerocallis flor. purpuralcentibus striatis F. 710. Grows by River fides.

7. Salfilla or Purple climing Lilly.

Hemerocallis scandens, flor. purpureis F. 713.

The Natives use the Root of this and attribute the same Virtues to it as Sarfaparilla, they infuse it all Night in cold Water, and drink it for Pains in the Stomach alto, and find great Relief by it.

8. Illeu or Chili Spiderwort.

Bermudiana cærulea, Phalangii ramosi sacie F. 715.

Grows on Mountains.

5. Tellow Star-leaved Salfa.
Salfa fol. radiatis, floribus fubluteis F. 716.
The Author fays this Plant is very well known in Europe for its Use in Venereal Diseases.

To. Nuil or Gland-flag.

To in Nuil or Gland-flag.

To in Nuil or Gland-flag.

This Flowers all white. The Natives drink a Tea of it in Stoppage of Urine proceeding from the Gravel, it's also good to expel Wind. Grows on Mountains and dry Places. 11. White Gavilu with a yellow Lip.

Epipadis st. albo F. 727.

Grows 3 Foot high, in the same places with the last, and has the like Virtues.

Epipaliu fl. vitelcente & variegato F. 727.

Grows on the fides of Mountains a Yard high.

13. Gavilu with a large yellow Flower.

Epipallu amplo flore luteo F. 729.

The Indian Women newly laid in, mix the Juice of this Plant with their Broths to cause their Milk to return, which

TAB. IV. Medicinal Chili Plants.

Fig. 1. Fig. 1. The Indians car this Herb in their Soops to refresh them. Growt in moift Ground and by River fides.

2. Bloody, ftreakt Rampion.

Rapuntii facie, fol. finuatis, flore amplissimo sanguineo & striato F. 729. ... Rivers and in moist places 2 Foot high.

3. Tupa or Chili Cardinal Flower. Rapuntium fpicatum foliis acutis F. 739.

Its Flower red. Grows on Mountains. The Root and Bark yields a venemous Milk, which will endanger the Eyes like Spurge, the very Smell of the Flower causes Voiniting, and the whole Plant is reckoned a violent Posson.

4. Tall Chili Violet.

Viola arborescens, Origani acuto solio F. 738.

A Tea of this is opening, its Flower blew and without Scent, Grows by River sides a Foot and an half high.

5. Blew Chili Milkwort.

Polygala cærulea, angustis & densioribus Foliis F. 721.

The Natives make a cold Infusion of this Herb all Night in Water and drank in the Morning proves a strong Diuretick and eases Pleuritick Pains. Grows on Mountains.

6. Chili Panke.

Panke Anapodophylli folio F. 741.

They eat the raw Stalks peeled, which are of a sweetish agreeable Taste, they also drink a Tea of its Leaves, which very much refreshes them in violent Heats. The Tanners boil the Roots of this with their Skins which very much thickens them, it also yields a black Dye. Grows in moist boggy Places and by River sides. ces and by River sides.

7. Chili Llau-panke.

Llau-panke amplissimo Sonchi folio F. 741.

They apply the Juice of this Plant to ease the Pain and stop the immoderate Flux of the Piles, they use also the Grounds of it Poultise-wise. The Dyers mix it in their Compositions to stain Black. Grows about a Yard high.

8. Pichua with Purstain Leaves.

Tithymalis perennis Portulace folio F. 707.

The Inhabitants of Chili, &c. min fome Drops of the Milk of this Spurge in a large Glass of Water or Broth, which moderately and kindly purges them, for which Cause they call it the weak or Female Pichua. Grows commonly on the Sea Coasts in floney and dry Places.

9. Veiny heart leaved Spurge. Tithymalus fol. trinerviis & cordatis F. 709.

Grows with the last and purges like it.

10. Male Pichua Spurge. So call'd because it works more violently both upwards and downwards. It differs from the others, but grows in the same Places.

11. Chili Ortiga.

Ortiga urens, Acanthii folio F. 757. Grows in Valleys about a Cubit high. N.B. Of this Species is the Frutex pungens & urens Marcgr. 79. Fig. or the stinging Palma Christi of Waser 27. p. 237.

11. Dyers Itiu.

Peryclymenum fol. acutis, flor. profunde diffectis F. 760. Grows about 4 Yards high, with Scatlet Flowers on the sides of Mountains. They use the Chips of this Wood with the Roots of Panke at Fig. 6. mixt with a black Earth call'd Robbo, these these they have the state of the Wood with the state of the at Fig. 6. mixt with a black Earth call'd Robbo, these they boil in Water to a just height, with which they dye their Stuffs of a sine Black, which fades not like those of Europe.

12. Pillabilcum or Chili Polypody.

Polypodium 1ad. squamosa F. 755. Filix Dadyloides perelegans Mus. Petiver 215. Ray Vol. 3. p. 72.

A Tea of this opens Obstructions and the Natives drink of it, when their Stomaches are oversharged. Found on the side of the

when their Stomachs are overcharged. Found on the fide of a Mountain near Pincho in Chili. It also grows about Mount Scrado on the Coast of Africa.

13. Chili Guilno or Furging Oat-grass.

Gramen Bromoides Catharticum F. 705.

Gramen Bromoides Catharticum F. 705.

They make a Tea of its Roots which they drink hot in a Morning which purges them; its Taste like its Operation little different from Sena. Grows in marshy and wet Places.

TAB. V. American Gum-trees and others very desirable. Fig. 1. Dellium. Is said to flow from the Trunk of a Thorney Tree, call'd Bdellia growing in Arabia and India, its Leaves are like the Oak, the Fruit resembles a Fig and of a pretty good Relish. The Gum of a bitterish Taste and turns yellow upon the Tongue, the best comes in oval Drops, is fragrant, reddish and transparent, it's used both externally and internally, being aperitive, sudorifick, directive and discussive, it hastens Birth; provokes Terms and resists Poison. fifts Poifon.

2. Cancamum. Resembles several forts of Gums or Rosins of different Colours clotted or sticking together, it's said to come from Arabia, Pomet says it was brought him from America A. D. 1686, by Dr. Brifet of 4 different Colours clinging together, issuing from a Tree of a moderate height, Lemery says it strengthens the Stomach and Bowels, kills Worms and stopens Obstructions of the Splan.

fays it strengthens the Stomach and Bowels, and opens Obstructions of the Spleen, &c.

3. Bailam Copaiba, by some Colocai and by the Portuguese Gamemolo. Marcgrave says the Wood is very red, which they saw into Boards for divers Uses, its L. 4 or 5 Inches long and oval

shell'd, with a Pilbert-Kernel, which bruited the like reage. Pijo adds that the Fruit ripens in April which the Brasilians eat, by sucking the Juice and spitting out the Skin; the Monkeys also are very greedy of them.

This Balm is yellower than Turpentine, and of a different Smell and Taste. It comforts the Stomach and Bowels, sweetens the whole Mass of Blood and is an admirable Vulnerative respective powerfully and stops Flurge of all kinds. smell and latte. It common to the stationers the whole Mass of Blood and is an admirable Vulnerative, provokes Urine powerfully and stops Fluxes of all kinds, it's fudorifick and cures most Diseases where Sweating is proper and equal, if not excells all other Natural Balsams. It's said the Natures learnt the vulnerary Virtue of this Balm by observing the Hogs to repair to it when wounded.

4. Caranna. Flows from the Trunk of a Tree like a Pain, which grows plentifully in New Spain. Mr. Lemery says it's fo call'd from Carthagena in the Spanish West Indies where it comes, where it is called by the Brassians Ibirapitanga Which grows plentifully in New Spain. Mr. Lemery says it's a thick large Tree with a reddish and thorney Bark, the call'd from Carthagena in the Spanish West Indies where it L. small and blunt of a sine shining green. Its Flower little, sweet and of a beautiful red, with Pods stat and prickly. This Wood is used amongst the Dyers, and the Stationerrs make a

which grows plentifully in New Spain. Mr. Lemery fays it's so call'd from Carthagena in the Spanish West Indies where it grows in plenty. It's so famed a Cephalick, Arthritick and Vulnerary, that it's usually said what Tacamahac cannot cure, Caranna can. Hernandez says it's a large Tree with longish

L. growing Star-wife.

5. Copal. It's brought from New Spain and other Parts of America. Pomet says this Gum flows from the Trunks and Branches of several large Trees growing on the Mountains with

Fruit like our Cucumbers but of a dark grey, in which is a mealy Flower of a very good Taste.

6. Elemi. This Gum, tays Pomet, slows from the Cut Trunk and Branches of a midling Tree with long and narrow silvery L. and red Flowers, the Fruit of the Colour and Shape of Olives, and therefore by some named wild Olives. It's of singular Use in all Wounds and Frastures in the Head and Scull.

7. New Baifam. In Colour and Shape like Tolu, but of a less agreeable Smell, its boil'd from small red cluster'd Fruit, whose L. are very large and broad. Grows in St. Domingo and

the other Ifles.

8. Ballam Peru or Natural Balfam. This Pomet afferts, is made by boiling its Bank, Branches and Leaves in Water, feurmaing off the Fat which is this black Balm. It's used as a

great Pedioral, particularly in Astma's.

from the Bodies of large thick Trees II. Lacam. 1.32

Aldersgate-street, London. May 21. 1715.

oval, bears a midling s leaved Flower, its Pods roundish, thin shell'd, with a Filbert-kernel, which bruifed smell like Pease.

Piso adds that the Fruit ripens in April which the Brasilians eat, by sucking the Juice and spitting out the Skin; the Monkeys also are very greedy of them.

This Balm is yellower than Turpentine, and of a different pours. It's so famous amongst the Indians, that they see in all Pains whatever. pours. It's so famous in all Pains whatever.

Wood is used amongst the Dyers, and the Stationerrs make a red Ink of it.

14. Campechy, Logwood or Brafiletto.

In the Bay of Campechy it grows plentifully and thither our English go to cut it, it's also found in other Parts of New Campechy and Campechy and there are whole Forrests of it. Spain as at Sanda Cruce where there are whole Forrests of it. Pomet and Lemery are both nistaken in making this Tree the same with our Jamaica Pepper, or All Spice., which we much better know than the Logwood, whose Flowers and Fruit we are as yet Strangers to.

15. Nephritick Wood. By reason it's a Sovereign Remedy for the Stone, Gravel, and Difficulty of Urine, it's also good in Obfructions of the Liver and Spleen. Grows in New Spain and chiefly about Mexico. Hernandez says it's a large Shrub with little roundish Leaves, its Flower yellow, long and little, growing in Spikes. The Mexicans call this Tree Coatli and Tla-

ning in spines. The mexicans can this free comm and tiapalz patli.

16. Jefuits Bark or Quinquina.

This Figure is coppied from Pomet, who fays it grows on the
Mountains near the City Loxa in the Province of Quito in
Peni; its Size about that of a Cherry tree; the L. round and
indented, its Flower reddish, with a Pod including an Almondlike Vermal covered with a thin Skin. Its Bark is much here great Pedoral, particularly in Asima's.

9. Sarcocol o Fleso-glue. Is a granulated Gum said to flow from a little prickly Shrub growing in Arabia, Persia, &c. It agglutinates Fleso. heals spongey Wounds, and stops Hemorrhages, it's also a very good Ophthalmick, taking away Spots, Speeks, &c. hindring Deflucion into those Parts.

10. Liquid Amber. Comes by Incision from the Bark of a large Tree with I-y Leaves growing in Virginia and Mexico, where it's call'd Ocosolt or Ococol: It's very fragrant, control en and mollisies the Hardnels of the Womb, is good in Speaks, Recumentation and Weaknels of the Nerves.

11. Laurum 12.2. Troop the Bodies of large thick Trees

Fames Petiver, F. R. S. Lond.

Books, Tracts, &c. Published by the Author.

E Piscibus Fluviatilihus Anglicanis. An Account of our Fish water Fishes, viz. such as are found in Lakes, Miss, Poots, Ponds, Eraks or Rivers. Price 6d.

Cochlearum & Grustaceorum Anunclium Britannicorum Ca-

talogus. Price 8 d.

De Scarabæi: Anglicanu. In which are named and described near 250 different forts of English Beetles in a Classical Method Price 1 s. 6d.

Price 1 s. 6d.

De Bombyliis Anglicanis. Of English Humble Bees, Pr. 2d.

Mr. Ray's Method of English Plants illustrated. Price 1 s.

English Plants, with stamineous, thready or Blink Flowers,
No. 120. whose Figures are already Engracen. Price 2d.

Mr. Ray's British Plants. Illustrated with 600 Figures, in
Folio Copper Plates. On which are Engraven their Names to

each, with References to his History and Synopsis of British Plants, as also the Places were growing, Colour of their Flowers, and Time of Flourishing, if Annual or Perennial, Common or Rare, Medicinal or otherwise. The Flower, Seed, Vessel, &c. Figured from the Life, and added to each different kind. Price One Guinea and an half.

The Englis or Latin Engraven Catalogues of the abovesaid

600 Plants. I s. Ed. cach.

Labels for Specimens of English Wild Plants. 1 s. 6d.

Labels for Medicinal English and Foreign Plants, 1 s. 6d. and if defired, the Specimens theinfelves.

Botanicum Londmerse, o London Herbal. Giving the Names Descriptions and Vitties, &c of tuch Plants about London, as have been observed in the several Monthly Herborizings made for the Use of the young Apothecaries and others, Students in the Science of Botany or Knowledge of Plants. Price 2s. 6d. Catalogues of Sheppy, Kentish and Glocestershire Fossils. Pr. 6d.

A Voyage to the Levant, viz. Sardinia, Sicily, Cyprus, Scanderoon, Rhodes, Stanchio or Coos, Samos, Scio and Smyrna. Giving an Account of each Place, their Inhabitants, Language, Coini, Weights and Meafures, their Provisions and Prices, Animals, Vegetables, &c. to which are added some Plants lately discovered on the Islands of Chio and Samos. Price 8d.

An Account of several Collectors of natural Rarities in Ita-

Price 6 d.

De Animalibus Crustaceis Caudatis. Of divers Crustaceous Animals, as Lobsters, Crawfish, Pawns, Shrimps, &c. extracted from P. Bellonius and G. Rondeletius with Remarkson them. Pr. 6d.

A Catalogue of the more Rare Plants, now and lately growing in the Physick Garden at Leyden. Price 8 d.

Catalogus Plantarum Indigenarum Hispania & Italia, quonum Icones à Jacobo Barrelier nuper incisæ sunt. Price 4 d.

Plants observed on the samous Mountains and other Places about Geneva by the Celebrated Mr. Ray and other Botaniss.

Price Is. Madam Merian's History of Surinam Insects, abbreviated and

methodized, with fome Remarks on them. Price 1s.

The Virtues of several Sovereign Plants found wild in Maryland with Remarks on them. Price 6d.


Pterigraphia Americana. In which are the complete Figures

of near 200 different forts of rare American Ferms, with divers Musbrooms, Coralls, Sponges, Shells, &c. Price 10s.
Gazophylacium Natura. Containing the Figures of above 1200 different Animals, Plants, Fossils, &c. from all Parts of the World, with their Names, &c. in 100 Folio Copper Plates.

Price 1 1, 10s. The Catalogues and Descriptions separate 2s. 6d. Price 1 l. 10 s. The Catalogues and Descriptions separate 2 s. 64. Musai Petiveriani Centuria X. Figuris. Price 2 s. 6d.

N. B. Divers of these Trads are now so very scarce, that of fome of them, there are not 20 left.








Planta Medicinales & PERU. Medicinall PERU Plants. TAB. I.





Plantæ Medicinales e PERU & c. Medicinall PERU Plants. TAB. II



















Aromata INDIX, Radices & Gummi. INDIAN Spice, Roots & Gums. Felium Indum





.

.

JAMES PETIVER

his BOOK,

For a Collection of whatever Trees; Shrubs, Herbs, Grasses, Rushes, Ferns, Messes, Sea, or River-Weeds, &c. you shall find.

Directions for the gathering of Plants.

Before to gather that part of each Plant which hath either Flower, Seed, or Fruit on it, but if none, gather it as it is, and if the Leaves which grow near the Root of the Plant, differ in Shape from those which grow at the top, (which frequently happens in Herbs) be pleased to add two or three of those lower Leaves to compleat the Specimen. And of each Tree, Shrub, Herb, &c. be pleased to Collect 3 or 4 Samples.

- 2. On whatever Tree, Shrub, Herb, &c. you find any ripe Seed, or dry Fruit, as Nuts, Podds, Heads, &c. Besides the Specimen gather some of the Seed or I ruit apart, with two or three of its Leaves (and Flowers if it have any) putting them into a piece of wate Paper, marking it with the same Mark or Number you do the Specimen. The like to be observed if you find any peculiar. ty in the Wood, Bark, Root, Gum, Rozin, &c. of any Tree or Herb.
- 3. Wherever you go alboar, or into the Fields or Woods, carry with you the Collecting-Book (to gather the Sample or Specimens in, which you must shift into this Book the same Day, or within two or three at farthest after you have gathered them) also a Knife, Pen and Ink, or Pencil, three or four Sheets of waste Paper (with some Thread) to wrap the Fruit or Seed in.
- 4. Entitle each Collection according to the Place and Time you gather them in, viz. These Plants were gathered at the day of Anno Dom.
- 5. If to any Plant you can learn its Name, Vertue, or Use, it will very much add to the Illustration of it,

And to the Satisfaction of

Your most Obliged Friend and Servant,

James Petiver.



Plantarum ITALIÆ, Marinarum & Graminum, ICONES, NOMINA, &c.

TABULA PRIMA.

Fig. 1. Alga spiralis maritima Bocc. Ray H. Plant. 1850. Ad Oftia Tyberina, sequentibus.

Ad Offid Typerina, requestious.

2. Fungus Amanita, marinus purpurascens Barrel. pl. 1310.

3. Fungus Amanita, marin. gilvo-luteus Barrel. pl. 1311.

4. Dens Lupi Piscatoribus Bar. 1312. Sea Wolfs-tooth. It's a white or fleshy Fungose Body, about 2 Inches long with a po-

rose Bottom.

5. Penna marina carnea Barrel. 1313.
6. Penna marina purpurea 114. These Two are like Feathers in Quill and Plume, its base thorney, by which it fastens to other Rodies.

7. Penna marina spinosa rubra 1315. More fiesby and thicker

8. Penna marina spinosa grisea obsoleta 1316. More clammy 9. Fucus niger vermiculatus & viscosus 1317. Found on the Rocks near Pale Fort. than the rest.

10. Fucus Latifol. Albicans, crispus, pilosus 1318.

often whitish. The Fishermen eat it with Oyl and Vinegar.

12. Fucus fruticans Coralloides subluteus major 1320. It eats like the last, but grows much thicker.

13. Eadem minor 1321. Its Flesh coloured and yellowish, they

mix it with Sallets.

14. Fucus crispus virens & è gilvo luteus 1322. It's soft, and eaten while fresh; its upper Tusts thick set, curl'd and green; the lower broad and black colour'd.

15. Fucus crispus loricatus nigricans 1323. It's 3 Inches long, thick branched and covered with curl'd black Hairs

16. Fueus crispus niger, crasso & grumoso caule 1324. Its Staks 4 Inches long, black and scaley, often ending in netted

Tab. II. Orallinæ, & c. Maris Mediterranei.
Fig. 1. Fucus crispus niger tenuifolius 1325.

Fig. 1. Fucus crispus niger tenuifolius 1325.

2. Fucus fruticans tenuifolius atropurpureus 1326.

3. Corallina cristata minima 1328. It's soft and joynted, growing in Tusts of divers colours, as white, black, purple, greenin, &c.

4. Corallina minima bifida 1329. Grows in longish Tails or Skains of very fine pa e Threads much divided and ending forked.

5. Corallina minima Faniculacea 1330. Its Threads finer than

the last and not so closely set.

100 tast and not so closely set.

6. Corallina minima crispa, albida 1331.

7. Corallina crustacea bisida 1332. It's brown with pale tips.

8. Corallina sistulosa fragilis crassion 1333. Ray 65. pl. 2. Of a stess of the colour, each Joynt haif an Inch long.

9. Corallina Fistulosa fragilis subtilior 1334. Ray 66. Much less than the last and whitish, its tips swelled and pounc'd.

10. Corallina Astaci corniculation amula 1225. Ray 70. 20.

10. Corallina Aftaci corniculorum æmula 1335. Ray 79. 20. 11. Corallina fruticola pennata 1336. A fort of yellowish Sea Feather, with crooked, inch long, joynted L. facing one way.

12. Corallina capillacea è cine eo fusca 1337. Its L. very fine

12. Corallina capillacea è cinereo fusca 1337. Its L. very fine and soft, growing in Tails like Spikenard.

13. Corallina susca Faniculacea & mollior 1338.

14. Corallina reticulata 1339. This brownish netted Sea-moss, grows to other Plants, Shells, &c.

15. Lithophyton marinum albicans 1340.

16. Lithophyton cinereum rugosum 1341. CB. pr. 153. pl. 2. Both these and most of the foregoing are found about the Tybers Mouth and on the Shores of the Tyrrhene Sea.

Tab. III. Gramina Italiana. Italian Grasses.

Fig. 1. Ramen spica Briza majus Barr. 1154. Grows in Hedges and dry Places.

2. Gramen Spica Briza minus Barr. 1155. Common in Fields.

3. Gramen Sparteum spicatum, utriculis Festuca, majus 1156.

4. Gramen Sparteum spicatum, Lolii utriculis, minus 1157. Differs from the last in baving less and bearded Tusts.

5. Gramen Palustre locustis Eruca-formibus 1158. Its Spike near a Foot long and stiffs, the back of each Tust is round, prickly and composed of 4 chasses Rowes.

6. Gramen Loliaceum spurium hirsutum, aristis geniculatis, maximum 1150. The whole Spike a Sam or longer with lander

maximum 1159. The whole Spike a Span or longer, with flender awned Husks, its Leaves and Stalks hairy.

7. Idem medium seu vulgare 1160. Grows with the last but differs in being less and having narrower Leaves and senderer Spikes.

8. Gramen Loliaceum junceum majus 1162. It's of a pale green. Grows on Mountains near haif a Front high. 9. Gramen Loliac. Junceum minus 1163. Grows with the last, but spreads on the Ground and is less.

10. Gramen junceum, nodosum, minimum, capillare 1164. Grows in Hilly Meadows and but little taller than the Figure.

11. Gramen Caryophyllatum Loliaceum repens 1167. The

lower Leaves broad, Stiff and pale, its Stalks two Cubits tall.

tower Leaves broad, July and pate, its Statks two Cubits tall.

12. Gramen Alopecurinum viride & molli spica 1179. It's fost downy Spikes 2 Inches long and hairy.

13. Gramen Cistatum perenne majus 1186. Its Spikes 4 Inches long with party-coloured Glumes or Chaff of pale and green, in which and being slenderer it differs from the next.

14. Gramen Cristatum annuum minus 1187.

15. Gramen Alopecurum pratense panicula seriosa -- 90

15. Gramen Alopecurum pratense, panicula sericea t188. 16. Gramen Typhinum prat. tuberosum, spica asperà 1189. ab. IV. Rainen Typhin. arvense rad. tuberosa, spica aspera, Tab. IV. Rainen Typhin. arvense rad. tuberosa, spica aspera, Fig. 1. Conga 1190. Its Roots more fieshy and less threddy than the last.

2. Gramen Typhinum fupinum tuberosum, spica non aspera 1191. It's broad Leaves, crooked Stalks and Jmother Spikes distinguish it from the rest.

Gramen Typhinum fupinum Italicum minus 1192. Head's foft and short.
4. Gramen Typh. maritimum, longius radicatum 1193

5. Gramen Typh. Phalaroraes majus bulbol, aquat. 1200.

6. Idem minus 1201. More flender and prickly than 5.
7. Idem Spica pilosa 1202. In watery Ditches with the 2 last.
8. Idem Arvenie 1203. On the Walls of Kome, &c.

9. Gramen proliferum ci ipumi majus 1211.

10. Gramen proliferum, panicula janthina minus 1212.

11. Gramen panicula crifpa 211. These 3 about Rome.

12. Gramen Etanthemum Palustre, tenui & sparsa Briza panicula 1216. Grows on wet Banks by River sides.

12. Gramen Phalysoidestinaria Rivas panicula minus 222.

13. Gramen Phalaroides sparsa Briza panicula minus 1211. 14. Gramen Phalar sparsà panicula, minimum angustif. 1218.

15. Gramen Tremulum majus, erecta panic. citrinum 1222. 16. Gramen Tremulum minus, erecta collectaq; panic. 1223. Tab. V. Ramen Phalaroides majus, acerosa erecta prig. 1. I nicula 1224. Grows on Mountains.
2. Gr. Phalar. majus acerofum, nutante spica 1225. Ramen Phalaroides majus, acerosa erecta parva pa-

3. Gr. Phalar. minus, erecta spica, obliquis aristis 1226. 4. Gr. Phalar. sparsa & nutante Briza panicula, obliquis aristis, minus 1227.

5. Gr. Avenaceum angustifol. panicula pyramidali 1233. 6. Gr. Sparteum Bromoides, brevioribus utriculis, nutante pa-

niculâ 1234. It's a rough dry Glass.
7. Gr. Spart. Brom. longioribus & erectis utriculis 1235.
8. Gr. Spart. Brom. panic. spicatâ, majoribus utriculis 1236.
9. Gr. Sparteum five Ægylops Spartea villosa 1237. 10. Gr. Festuceum Myurum elatius, spica heteromalla gra-

11. Scirpus maritimus capite glomerato 1246. Ray 1298. 17.


12. Scirous marit, minor, graciliore spica 1240. 113
13. Scirous capitulo singulari, rad. tomentosa 1249.
14. Cyperoides latifolium minus 1256.


15. Cyperoides latifol. typha erecta minus 1257. Shady Hills.
16. Cyperoides angustitol. cypha nutante minus 1258. Found in Woods.

L O N D O N:

Printed for James Petiver in Aldersgate-street. MDCCXV.

Where may be had Mr. RAT's English Plants Illustrated with 600 Figures on Folio Copper Aquatilium Animalium Amboina, Icones & Nomina, price 10.5. Plates. Price 1 l. 12 s. 6 d. Pterigraphia Americana; In which are Figured near 200 rare Sorts of American Ferns &c. Pr. 10 s. Hortus Peruvianus Medicinalis, or South-Sea Herbal; Containing the Figures, Names, &c. of divers Medicinal Herbs and Gum Trees from those Parts. Price 2 s. 6 d. Musai Petiveriani Centuriæ X. Figuris. Price 2 s. 6 d.









Musei marini, Corallina &c. Mauis MEDITERRANEI. Levant Sea Mosses & c. TAB
















RARIORUM JCONES

Aliarum minus vulgarium CATALOGUS.

A Jacobo Petiverio, S. R. S. Lond.

Tab. 1. Fig. 1. Alix Ægypt, latifol. ex alis Julifera nobif.
Calaf & Ban Ray Hift, Plant. 1424. Chabr.
66. f. 4. IB. 220. f. Salix Ægyptiaca Park. 1433, fig. 6.
2. Sebestena sylv. CB. 446. 2. Frudu minore serotino Nob.
Ray 1555. c. 25. Var. Chabr. 12. f. 6. IB. 198. f. 2. Park. 552.2.
3. Tamarix Ægyptia arbor CB. 484. 3. Ray 1705. 3. Tamariseus Ægyptia gallas serens Park. 1479. 4.
4. Balsamum Syriacum Rutæfolio CB. 400. 1. R. 1755. c. 13.
Chabr. 24. 3. IB. 298. f. Park. 1528. c. 80. f.

4. Balfamum Syriacum Rutæ folio CB. 400. 1. R. 1755. c. 13. Chabr. 24. 3. IB. 298. f. Park. 1528. c. 89. f.
5. Abavo arbor, rad. tuberofa CB. 434. 10. R. 1371. 1.
6. Cucumis Ægypt. rotundifolfus CB. 310. 5. Phyt. 618. 5. Chabr. 135. f. 2. IB. 248. fig. Ray 646. 4.
7. Melo Ægyptius CB. 311. 5. phyt. 620. 4. Chabr. 134. f. 4. IB. 245. f. Ray 945. 4.
8. Kali Ægypt. fol. valde longis hirfutis CB. 8. p. 289. Ray 1915. pl. 1. Chabr. 543. f. 5. Park 179. 3. & 1285. f. 4.
9. Matricaria inodora CB. 134. 5. Phyt. 223. 4. Ray 357. 2. Chabr. 374. f. 5. IB. 131. f. Park. 85. 8.
10. Loto affinis Ægyptiaca CB. 332. 1. Ray 922. 1. Chabr. 159. f. 4. IB. 361. f. Park 849 3.
11. Melilotus Ægyptiaca, corniculis reflexis minor CB. 331.

159. f. 4. IB. 361. f. Paik 849 3.

11. Melilotus Ægyptiaca, corniculis reflexis minor CB. 331.

3. Phyt. 666. 45. Ray 953. 13. Park. 719. 5.

12. Lenticula palustris Ægypt. sive Stratiotes aq. sol. Sedi majoris latoribus CB. 362. 6. Ray 1324. c. 8. Chabr. 567. f. 2.

1B. 787. Park. 1249. f. 2.

13. Stratiotes Ægyptiaca fol. obtusis Nobis. Ray 1324. cap.

8. 2. Park. 1249, f. 2. inferior.

14. Herba Ægypt. sol. Junci, succo slavo urente Nob. Felsel tavil Ægyptiis CB. 412. 5. Chabr. 126. f. 4. IB. 188. f.

15. Cyperus rotund. Oriental. major CB. 13. 1. Phyt. 13. 1.

Th. 208. 1. Ray 1294. pl. 13. Chabr. 194. f. 5. IB. 502. sig.

16. Papyrus Nilotica sive Ægyptiaca CB. 19. 1. Phyt. 44. 1.

Theatr. 323. 1. Ray. 1320. c. 2. Chabr. 195. 5. IB. 460. f. 1.

Park. 1207. f.

Park. 1207. f.

17. Gramen Dactylon Ægyptiacum CB. 7.1. Phyt. 90. p. 20. Theatr. 116. 1. Ray 1272. 3. Chabr. 180. f. 5. IB. 14. f. Paik. 1294. f. 7.

Tab. 2. Fig. 2. Icus folio Mori, fructu in caudice ferens CB. 459.1. Ray 1439.1. Chabr. 9. f. 5. IB. 124.

fig. Park. 1294. f. 1.
2. Oenoplia spinosa CB. 477. 2. Ray 1534. 4. Chabr. 51. f. 3. IB. 39. f. 1. Park 1442. f.

3. Lycio affinis Agyptiaca CB. 479. 10. Ray 1629. 10. Chabr. 52. f. 6. IB. 62. f. 2. Park 1011. f. 12.
4. Lycium Indicum alterum CB. 479. 8. Ray 1128. 9. Chabr. 53. f. 4. IB. 62. fig. 1. Park. 1011. f. 10.
5. Myrobalanus Ægypt. spinosa, foliis geminis Nob. Park. 1678. f. 2. Vesl. Obs. 61.

6. Hippomarathum Sphærocephalum Ægypt. Alpin. Exot. 198. f. Ray 459. 4. Park. 884. 6.
7. Marum Ægyptiorum Alpini Exot. 253. f. Park. 1674. f. 10.

Veil. Obs. 79. t.

8. Cucumis Ægyptius reticulatus seu Lussa Arabum Vest.
Obs. 48. sig. Ray 640. 2. GAZ. NAT. Tab. 11. sig. 2.

9. Hyoseyamus albus Ægyptius Alpin. Exot. 192. f. Vest.

10. Convolvulus Ægyptius Vell. Obs. 74. fig. Omel hessen

Arabibus. Ray 720. pl. 1.

11. Lysimachia lutea Ægyptiaca Ray 1022. 4. Cassabel darira Alpin. Exot. 194. f. Vest. Obs. 63. Park. 189. f. 2.

12. Galega Ægypt. siliquis articulatis CB. 352 2. Ray 926. c. 3. Park. 1099. t. 6. Sesban Ægyptiis.

13. Galegæ assinis Sophera dicta CB. 352. 3. Ray 911. 3. Park 1089. f. 8.

14. Arum Ægyptium minus Vest. Obs. 44. caule maculoso

15. Nymphea seu Nusar Ægyptium Vest. Obs. 43. c. 34. Nym phea alba major Ægypt. sive Lotus Ægyptia Park, Lotus Ægyptia Alpin. Exot. 214. s. Ray 1321. 5. Vest. Obs. 43. s.

Plantarum ÆGYPTIACARUM albis odoratis Nob. Chabr. 416. f. 6. 1B, 256. f. CB. 223. 9. Phyt. 421. 7.

Plantarum Ægyftiacarum aliarum Catalogus.

Cimum Ribau Ægyptiis Alpini 53. Vest. 132. Ray 187i. 2. Anguria Ægyptiaca CB. 312. 3. Phyt. 622. 3.

2. Anguria Argyptiaca CB. 312. 3. Laft C22. 3. Park. 773. 3. Ray 644. 4.
3. Melonis Egyptin alterium genus CB. 313.6. Phyt. 620.5. Chajan Egyptiis Alpini 45. Vefl. 115.
4. Braffica fpinofa CB. 311.3. pt. 54. f. Phyt. 178.16. Matth. 369. f. 6. Chabr. 271. f. 5. 1B. 835. f. 2. Park. 269. f. 3. Ray 797. 18. Alp. Exot. 200. f.
5. Lubinus pereoriums pentaphyllos CB. 248. 5. Vefl. Obf. 70.

5. Lupinus peregrinus pentaphyllos CB. 348. 5. Vest. Obs. 70. f. Pona Ital. 102. f. Alp. Exot. 322: f. Park. 397. f. 3. & 1075.

5. Ray 859. 2.

6. Thlaspi Alexandrinum CB. 108. 2. pr. 50. 10. Phyt. 165.

6. Thiatpi Mexandrinum CB. 108. 2. pt. 30. 10. Thyt. 105.
12. Park. 849. 3. Ray 834. 20.
7. Cicercula Agyptiaca Cluf. 236. f. 1. Lathyrus angustif. fem. maculofo CB. 344. 8. Ray 896. 15.
8. Phaseolus niger Lablab Alpini 31. f. Vest. 75. f. Obs. 24. Phaseolus Agypt. nigro semine CB. 341. 1. Cluf. 227. f. Pan. 732. Chahr. 140. f. 4. IB. 270. f. Ray 888. 25.
9. Phaseolus Agypt. sem. rusto CB. 341. 2.
10. Datura Agyptia Vest. Obs. 38. f. Contarena Alp. Exot. 167. Ray 748. 2.

167. Ray 748. 2. 11. Hyofcyamus Ægyptius Cluf. 83. f. 2. Park 463. f. 4. Ray

712. 5. 12. Beidelfar Ægypt. Alpin. 36. f. Vesl. 86. f. Apocynum Ægypt. st. stellatis croceis spicatis, folliculis magnis cordatis Nob. Humidis propè Alexandriam.

13. Beidel Star Medicis Arabum Offar & el Usar dictum Vest.

Obs. 28. f. Apocynum Hort. flor. umbellatis reflexis, folliculis hirtis Nok.

hirtis Noh.

14. Secamone Ægypt. Alpini 53. f. Vefl. 134. f. Ray 1087. 2.

Apocynum angusto Salicis folio CB. 303. 3. Park. 385. 3.

15. Melochna Alpini c. 28. f. Vefl. 93. f. Ray 1068. 10. Corchorus CB. 319. Phyt. 632 Lob. Ic. 585. 1. Belg. 597. f. Obs. 269!

16. Faha Ægyptiaca, Dioscoridis affinis CB. 196. 8. Chabr. 562. f. 3. IB. 775. f. Park. 374. f. ult. Ray 1322.

17. Felfel tavil Ægypt. Piper longum Ægyptium Vest. Obs. 35. f. Euphorbium Ægypt. marit. Juncis ramosis Nob.

18. Gr. stellatum Ægyptium Vest. Obs. 33. fig.

19. Capparis non spinosa, frustu majore CB. 480. 3. Park. 1023. 3. Ray 1630. 3. Cappar Alexandris Alp. 52. Vest. 129.

20. Sebestena Domestica, fol. latioribus & crassiotibus, frustu majore Nob. Alpin. 13. f. Vest. 30. f. Obs. 12. CB. 446. 1. Chabr. 12. f. 5. IB. 198. f. Park 552. 1. Ray 1555. c. 25.

21. El-hanne Ægypt. Alcanna Avicenna Alpin. 20. f. Vest. 47. f. Obs. 16. Ligustrum Ægypt, angustifolium CB. 476. 5. Chabr. 41. f. 5. IB. 541. f. Park. 1446. f. 3. Ray, 1603. 3.

22. Acacia Ægyptiaca vera Nob. Alpin. 7. f. Vest. 5. C. Obs. 8. f. opt. Chabr. 92. f. 1. & 2. IB. 429. t. Park. 1548. f. 1. Acacia fol. Scorpoidis leguminosa CB. 392. 1. Ray 976. 1.

23. Carub vel Carnub Ægyptiis Alpin. 4. f. Vest. 9. Obs. 5. Siliqua dulcis CB. 402. 1. Chabr. 89. f. 1. IB. 413. f. Park. 236. f. 1. Ray 1718. c. 2.

f. 1. Ray 1718. c. 2.

Planta Ægypti à D. Lippi observatæ; ex Lutetia nobis transmisse à Clarissimis Botanicis D. Seb. Vaillant & D. Lud. Stofchio.


24. Sonchus Ægyptiacus giganteus Nob. Bot. Hort. 2. pl.22.
25. Senecio Ægypt. fol. Matricaria Bot. Hort. 2. pl. 85.
26. Echium ferox Ægypt. fore carneo Boeib. 7. p. 79.
27. Hormmun Ægypt. frutic. nanum, Sideritidis folio.
28. Stachys Ægypt. talviæ minotis fol. fl. purpurafcente.
29. Linaria Memphitica, hastato fol. tricuspidato, fl. luteo.


30. Raphanistrum Ægypt. Siliquis recurvis.
31. Geranium Ægypt. Althew fol. glauco crassiori.
32. Fubugo Ægypt. fol. tereti bidigitato.
33. Alhagi Ægypt. fol. longiori, aculeis brevioribus.
N.B. Vide Planta alia Ægyptiacæ in Petiveriana II. No.465. &c.

Ab Autore nuper editi.

T. PETIVERIANA 1, 2 & 3. five Collectanea NATURA.
Domi Foisique Autori Communicata. Lond. Fol.
2. Papilionum Britannicorum Icones, Nomina, & c. Figuris plusquam 80. Tabulis Ameis incisi. Londini MDCCXVII.
Folio.

Horrus Peruvianus Medicinalis: or the South Sea Herbal Fig. Encis. Fol.








Plante AGYPTIACA Pr. Alpini & Veslingij Egyptian Trees & Herbs 7/1BII.





PTERI-GRAPHIA AMERICANA

Jones continens plusquam 400 FILICUM variarum Specierum, Viz. Arborescentes, Scandentes, Spinosas, Floriferas, aliasque perraras; nec non Muscos, Lichenes, Fungos, Corallia, Spongias, aliaq non pauca Submarina sui adjeuntur Crustacea, Testacea, aliaque Animalia ferè omnia ex Insulis nostris Charibbæis Viz. Antego, Barbados, S. Christophers, Nevis, Jamaica &c. Figuris Anais, Folio incisis, XX Tabulis. A IACOBO PETIVER. Soc. Regal. S. LOND.

```
FILICES Seminibus maculis rotundis.
                                                           60 . . rotundis planis 1.5.
                                                           61 . Taxi foly's major 2.5
      Phyllitides folijs integris.
1 Phyllitis maculata, amplificino folio Tab 6 fig. 8.
                                                           62 . . . minor 2. 6.
                                                           63 . mas Ensiformis major 1.3.
2 . . . breviori & crassiori folio 6.1.
                                                           64 . . . vix serrata 1.4.
3 . . latifolia, pediculo & cofta spinosis 4.14.
                                                           65 . Betonicæ folio 1.7.
   . . folio longo leviter undulato 6.9.
                                                               . . . . vix serrata 7.15
  .... & marginato 6.10.
                                                           57 . . auriculata, serris bifidis 2.9.
  .... apice palmato 6.11.
  angustifolia 6 3.
                                                                  Filiculæ seu Filices minores.
8 .... maculis majoribus 6.2.
                                                           68 Filicula Rutaceo folio 1.12.
  .... lineis maculata 10.3.
                                                           69 . pinnis planis, caule hirt 4.11.
     Phyllitides ramosæ.
                                                              . . dentatis, caule squamoso 9.6.
                                                           70
10 . ramosa, maculis majoribus 6.13.
                                                           71 . . lunulatis 3.15
                                                           72 . alis inferioribus pendulis 1.9.
11 . . . minoribus 6.14.
                                                           73 . . . auriculatis 1.8.
12 . . maculata scandens, caulibus squamosis 4.15.
13 .... angustifolia 4.16.
                                                           74 . . laceratis, tota translucens 3.14.
14 . mac. 5 folia. pediculis squamosis 4.13.
                                                                Trichomanes punis simplicibus.
      Polypodij seu Filices pinnis alatis.
                                                           75 Trichomanes folio cordato 1.11.
                                                               . . triangulo dentato 1,10.
15 Polypodium maximum 7.8.
16. .. auriculatum 7.5.
                                                            77. argenteum ad oras nigrum 9.16.
17 . . . pentaphylloides 7.6.
                                                               Adianti pinnis serè angulosis & membranaceis.
18 . . . trifoliatum 7.3
19 · auritum majus laciniis crispis 7.7.
                                                           78 Adiantum ramosius repens spinosum 11.5 & 6.
                                                           79 . . frutescens, pinnis subrotundis 5.12.
80 . . altisimum pinnis globuliferis 2.12.
20 . . tricuspidatum, oris pilosis 7.4.
21 . scandens Quer cus folio. caulibius Lycopodii 15. 2.
                                                           81 . . alis longis crispis 5.13.
22 . minus vulgaris facie 7.9.
 23 - . Afplenij folio 1.14.
                                                           82 . . pinnis Quercinis ramis villosis 4.5.
                                                           83 . mas, caule squmoso 5.3
 24 ... brevi pendulum 1.15.
                                                           84 . . . & pediculo pulverulento 5.7.
 25 ... crifpo, peudulum 13.12.
26 . . . villoso 7.16.
                                                           85 . . pinnis acutis 5.9.
 27 . longifolium, radice dulci 7. 10.
                                                               . . . subrotundis 5.10.
 28 . maculis minoribus 7.10.
                                                           87 . . . crispis, lunulis albicantibus 5.11.
                                                           88 . . . speciosis 5.14.
 29 . . squamulis argenteis 7.11.
30 . crifpum Struthionis pennis 3.8.
                                                           89 . Trichomanoides summo crenatum 1.13.
                                                           90 . . . & crifpum 10.8.
 31 . Lonehitidis folio 7.14.
 32 · · · undulato minus 1.16
                                                           91 · · pinnis cristatis 5.8.
 33 ... angustissimo pendulum 14.1.
34 · vulgare, summo trisido 15.3.
                                                           92 . . . latis 5.5.
                                                           93 · · alis radiatis 14.6.
 35 Taxi folio pendulum 14.1.
                                                           94 . pinnis hexagonis furcatis 10. 2,
 36 . pendulum subtus pilosum 12.3.
 37 · serratum majus, cofta alata 7.13.
                                                                   Adiantuli pinnis clavatis.
 38 . minus, fol. integris & serratis 14.2.
                                                            95 Adiantulum clavis simplicibus 14.7.
                                                           96 . . . capillacevm 10.7.
        Filices ramosæ Fæminæ dictæ.
                                                           97 . clavatum Crithmi fol.majus 5.15.
 39 . ramosa, aculeata repens 4.6.
                                                           98 . . . minus 5.16.
 40 . . . piramidalis pinnis parvis 4.12.
                                                            99 · triphyllum 11.4.
                                                           100 triphyllum repens 11.9.
 41 . . arborescens, pinnis dentatis 1.1 & 2.
 42 . . . . latis caudice spinoso 4.1.
                                                           101 . membranaceum Lichenoides 13.5.
 43 · · Cicutæ folijs 2.8.
                                                                  Darea seu Filices pyxidatæ
 44 . . globulifera major 2.11.
 45 .. pinnis dentatis, caule piloso 4.9.
                                                           102 Darea ramosissima scandens 12.5.
 46 .... & coftis hir sutis 2.7.
                                                           103 . Tilicis maris facie n. 8.
 47 . . . arborescens caudice & caule spinosis 4.7.
                                                           104 :Timbrigensis facie 13.13.
 48 . . . . humilis & spinosa 4.8.
                                                           105 · affinis digitata 15.5.
 49 . . . latis laceratis 3.13.
 50 . . . gladiatis, caudice spinoso 5.1.
                                                                  FILICES Seminibus striatis.
 51 . . furcata. Taxi folije 5.4.
                                                           106 Phyllins major, margine crenato 6. 7.
                                                           107 .folio, geminato dentato 13.11.
     Filices minus aut non ramosæ Mares dictæ.
                                                           108 ramosa margine membranaceo 12.2.
 52 . dentata caule & costa villofis 4.10.
                                                           109 . . . & strys undulatis 7.1.
                                                           110 Lonchitis striata 3.2.
111 . auriculata & crenata 3.1.
 53 · pinnis aculeatis 1.6.
 54 · acutis planis majoribus 2.1.
 55 . . . . minoribus 2, 2,
                                                           112 · · ramofa · caule squamosa 5.2.
 56 ... parvis 2.3.
57 ... ad alas foliosa 5.5.
                                                           113 Filix Striata pinnis crenatis, major 3.3 .
                                                           114 . . . minor 3.4.
 58 . . latis summo dentatis 2.10.
                                                           115 Adiantum streatum, nigri facie 3.7.
 59 ... apice piramidali 2.13.
                                                           116 . . alis laceratis 3.5
```

117 latioribus 2, 14.



```
Agarici, Boleti Tungi, Tuberæ &c.
118 Adiantum striatum, cacumine bisecto 3.6.
119 ... pentaphy lhum, humile 11.7
                                                                         190 Agaricus amplifsimis squamis 17.3.
120 Richomanes striatum, vulg. facie 2.15.
121 pinnis bifidis. 2.16.
IILICES sem. medio, vel margine striatis.
                                                                         191 . cinereus & sericeus 17.15.
                                                                         192 . lutous Clypeiforms transfixus 16.13.
                                                                         193 niveus Brassicam crispam referens 13.6.
                                                                                . Rosco-niveus, strys aureis rugosus 17. 2.
122 Phyllitis angustifolia, lineis catenatis 6. 4.
123 · lineata, hine inde dentata 6. 5.
124 · apice trifido. 10. 6.
                                                                         194
                                                                         195 Boletus cancellatus purpureus 17.11.
                                                                          196 . Phalloides, pilicolo villoso 17.9. G.H.
                                                                          197 . . rugosus, pedic . fiftuloso 17.10.I.K..
198 Coralloides fusca & laciniata 16.14.
125 . aspera, furcis lineatis 6. 6.
126 . lineata, Graminis fol. longi/simo 14. 3.
                                                                          199 . ramosilsuna purpurascens 16.15.
127 . ramola and margines pulverulenta 6.15.
                                                                          200 Fungus aureus laciniatus 17.7 E
201 . coccineus, squamosus & globosus 17.4 AB.
202 . crenatus, tenuissimus & niveus 17.6 D.
128 Lonchitis juxta nervum pulverulenta 3.9.
129 lineata, pinnis longifsimis 6.12.
130 ... ad başin foliosa 12.1.
131 . ramosa, limbo pulverulento 3.10.
                                                                          203 . In latus totus niveus 17.5 C.
                                                                          204 Amgoides Cyathiforme coccin.oris pilosis 13.8.
 132 Filix lineata arb. caudice aculcato 4.2.
                                                                                  . candidum, intus villosum 17. 4.
133 . . auleata, apice pyramidali 4. 3.
134 . Lonchitidis pinnis 3. 11.
                                                                          206 18. purpureum 16. 16.
                                                                          207 . retreulatum Flabelliforme 17. 8. F.
208 . rugatum Infundibuliforme & pwp. 13. 7.
135 . Fæmina vulg. facie 3.12.
136 Adiantum limbo nigro præcinctum 9.15.
137 Trichomanes avgenteum, ad oras nigrum 9.16.
                                                                         209 . scutellatum purp. deforis villosum 15.10. & 17.5.
                                                                          210 Lycoperdon coronatum 17.12.L.M.
        FILICES Pulverulente.
                                                                          211 Tubera candida mollia 17. 14.N. & 13.
138 Phyllitis pulverulenta angustifolia, caule crinito 8.7212 Testiculorum forma, majora 16.17.
139 . . . & oris crinitis 8.10.
140 . . glabra, & rigida 8.8.

SUBMARINA viz. Corallij, Spongis
                                                                                 . . . minora 13.0. 10. A. SUBMARINA riz. Corallij, Spongiæ, Fuci &c.
 141 .. fol. angustis trifidis 8. 9.
                                                                          214 Corallium album, nodulosum 18.5.
 142 . . ramosa aurea 8.5.
                                                                          215 . . stellatum , ramosum 18.4.
216 . . Fungoides , ovale 20.16.
 143 . . nodosa major 6. 16.
 144 . . nodosa minor 9.7.
145 . . & crimita latissimofol. 13. 14.
                                                                          217 . . . crispum 20.17.
218 . . virgulis minoribus 18.6.
      .... longifol. 9. 14.
                                                                          219 . porosum, latum planum 18.13.
220 . . . muricatum 18.7.
 146
 147 . . lanfolia villosissima 9. 13.
 148 Hemionitis max. pulrerulenta 5 folia 7. 2.
                                                                                 . . Typhoides muricatum 18. 18.
 149 Lonchitis pulverulenta scandens Citrij fol. 8. 6.
                                                                          221
                                                                          222 . . . stellulatum 20. 15.
223 . Roseum fossile 20. 20.
 150 . Vaniliæ folio 15. 1.
151 . pinnis Ensiformibus 8. 4.
                                                                           224 . rubrum crifpun 16. 9.
 152 . . . rix crenatis 8. 3.
                                                                           225 Alcyonium Cornu Cervi referens 16.8.
       . Calamifera, pinnis serratis 9.8.
                                                                           226 Afteria Coralloides, stellis maximis 18.2.
 154 Ofmunda pulverul. pinnis vix serratis 8. 1.
                                                                                      . majoribus 18.3
             scandens, sarmentis capillaceis 8. 2.
                                                                           228 Lithophyton Polypody ramosi facie 16.7.
  156 Filix albissimo pulvere subspersa 9.11.
                                                                           229 Madrepora cristata wens 16.10.
 157 . pulverul. ramosa, pinnis parvis 2.4.
158 Filicula pulverul. Crithmi face 9.4.
                                                                           230 Millepora perforatu 20.18.
                                                                           231 Spiculum Barbaderse Lapideum 20.21.
           pendula, alis digitatis 11.1.
                                                                           232Epicorallum reticulation vulg. 18.8.
  160Adiantum pulverulentum aureum 9.9.
                                                                           233 virgulis punctatis & asperis 18. 11. 234 . . alatis 18. 12.
  161 . . alis infra auritis 9.10.

FILICES in racemulis Floridæ.
                                                                                 .. muricans majoribus 18.9.
                                                                           235
236
  162 Osmunda racemifera Phyllitidis folio 8.13.
                                                                                 . . . minoribus 18.10
                                                                           237 Spongia dura, compressa foraminosa 18. 14.
  163 . . . vix crenato 8. 15.
                                                                           238 Typhoides for aminos a 18.19.
  164 . spicis oeminis 14.5.

165 . . . villosa 8.16.

166 . pinnis parris 8.14.

167 . Adianti nigri fa cie 9.1.
                                                                           239 . . ramosa 19.1.
                                                                           240 ... apicibus cavis 19.2.
                                                                           241 . . . . acutis, furis 19. 3.
242 . . maculata 19. 8.
243 . . muricata & ferè cara 19. 9.
       . . Asphodeli radice 9. 2.
  168
  169 . . . . Matricariæ folio 9. 2.A.
       ... Cicutæ folio 9 3.
... verticillata 12.4.
FILICES Milcellaneæ.
                                                                           244 . . ramosa & nodulosa 19.5.
  170
                                                                           245 . fibrillis reticulatis 18. 20. 246 . . . cava longa 18. 21.
                                                                                 foliata aspera 19.4.
                                                                           247
                                                                           248 .Infundibuli formis 19.6.
  172 Filix spinosa, sern. hunulatis 4.4.
  173 . Conifera scandens, pinnis varys 9.12. 249 . Labri formis 19.7.
174 . macul. scandens, fol. rotund & longis glabris $250 Corallina Opuntioides minor 20.19.
                                                                           251 Muscus marinus pennatus 14.4.
252 . triagulais 18.15.
253 . frutelcens, fruttu trcapsulari 16.11.
254 Sargazo dentatu rulg. 16.12.
255 . longifolia 19.10.
256 . latifolia 19.11.
  175 . . . fol. ovalibus & longis pilosis 10.5.
176 . mayinifera, Granij folio 8.12.
177 . Ranunculi folio, prolifera 8.11.
178 Ophioglosum Palmatum 11.2.
   179 . vulgaris facie 10. 4.
            Musci, Lycopody, Hepatice &c.
                                                                           257 . . angustifolia 19.12.
258 . argustifolia plana 19.13.
  180 Muscus denticulatus erectus 16. 5. 181 . . repens 12. 6.
                                                                            259 Fucus vesiculis alatis 18.17.
  181 . repens 12. 0.
182 . max. Linariæ folys 16. 4.
183 Lycopodium altismum 3. 16.
184 . Coridis fol. flagellis longis 16. 3.
185 . erectum, Sabinæ folio 16. 2.
186 . ramofismum crilyum 16. 1.
187 Hepatica capite palmato 15. 6.
188 . digitata Europides 11. 2.
                                                                            260 · velicarius, summo plants 18.16.
261 Ovarium JAM. resicis 10t: compressis 18.1.
262 Vesicaria corona serrata 17.1.
                                                                            ANIMALIA Ciriltacea, Teltacea, Insecta & c.
```

188 . digitata Hingoides n. 3. 180 Lichen cornua Damæ referens 16.6.

263 Perca BARBAD. cestá lineatá 20.3. 264 Lacerta BARB unquibus fuscis 20.1. 265 . Cæciliæ facie 20.2.




366 Cancellus BARB. brachy's tuberculosis 20.4. 386 Clavicula BARB. ore pulvinato 10. 11. 387 Rhombus BARB. fascia albida 12.7. quadratus 20.5. rug olus, pedibus hirtis 20. 6. 388 Sphoudyhus LAM. tessellata 12.8. 389 Tridaena Campeach .parva perelegans 15.11. .. ruber, lævis 20.7. 370 .. muricatus, compression 20.8. 390 Ichneumon max tripile 13.1. 371 . Arancoides 20.12. 391 . *majus* caudâ *lunatâ 13.* 2. 372 Mantis marina RARBADENSIS 20.20. 392 . affinis, antennis clavatis 13. 3. 373 Squilla BARB. ovalis 20. 9 . acutis 13.4. 393 . . 374 Araneus BARB. parvus, pedik maculosis 20.13. 394 .alis maculatis 11.10. 375 Capricornus BARB. Scaber, fuscus 20.14. 395 Musca Apiformis major 11. 12. 376 Cochlea BARB. compressa, minor 14. 13. 396 , , minor 11.13. 377 .IAMAIC. verruculata 10. 9. 10. 397 . . minima ñ. n. 378 .. umbilico patulo 14.11. 398 . alis nubulosis 12.12. 379 .. ore tridentino 14.14. 380 Fibula JAMAICENSIS 14.12. .. ore tridentino 14.14. 399 . . . anten . acutis 12. 14. 400 . . . minor 12.11. 381 Nerita Bass. alba, radis coracinis 14.12. 401 . . . minima 12.15. 382 ... vndis nigris 15.13. 402 . . . ano acuto 12.13. 384 Olivaris IAM. strijs capillaribus 20. 22. 386 . . . vndatis & radiatis 12.9. 403 Vespa impennis e croceo nigro 11.14. 404 . minor 11. 15. 405 Scolopendria BARB. preelonga pedibus plumofis 406 Perla major 15.7. 411 Musca alis opacis & vitreis 15.8. 407 . . maculus vitreis 14.10. 412 Cochlea IAM. ore lacerato x. 13. 408 . cornubus clavatis 15.9. 413 File: Androgymus, maculatus 9.5 409 . triangularis 14.9. 414 Muscus marinus dentatus 14.4. 410 . maculis vitreis 14.8. Cochleæ AMERICANÆ in Museo Nostro, quarum ICONES in Gazophylacio NATURE non Figurantur. 415 Pecten IAM strijs lævibus Mem. 1 List 168. 464 C. alba cun undis fuscis G. Searl. 416 P.Anteg. minor, undis fiwcis. 465 Musculus striatus minor, cardine serrato 1.365. 417 PBARB. albus minor, strijs lunulatis M.3. I. 177. 466M striat min rostro acuto latior L 362.201. 418 P.B. alb.min str. squamofis M.4. I. 176. 467 id. longior L. 362. 202. 419 Sphondy his Amer. max ruber 468 Pholas Virginiana L. 434. Car. 21. 420 SBARB rugis minoribus M.g. L. 215.50. 469 P. Campeach &c. cardine foraminoso L. 432 Car. 20. 421 S.B. rug. major. sulcatus M 6. L. 215.51. 470 P. CAR. & c card fert bidente L. 431 Car. 22. 422 SAmer rostro internè dextro M.7. 471 Solen Marianus latus I. 421. Car. 13 423 SAmer roltro int. sinistro M. 8. 472 Curassavicus latus, radijs biriis albis H. Ruysch. 424 Berberi IAM. auritus L. 224.59 473 Buccinum minus Anteg fascy's nigris M.17. 125 B IAM. tenuis Mg L. 223.57. & 8. 474 Cochlea IAM. alba, crassa M. 2. I. 571. 426B IAM. cardine serrata L 227. 475 C. IAM. flavicans, crassa M.3 I. 566. 427 Glycimeris I.AM striates perceases M 13. L 238. 476 C. aure magno fusco CAR. 29. 428 G. LAM. lare striates M x1 I, 240. 477 C. aurita fascijs catenatis CAR. 28. 429 id minor maculatus L. 245. 478 C. Amer. undis flavis M.4 430 GBARB. min striat. maculatus M 12 L. 243. 479 C. IAM. verrucis fasciata M. 10. 431 Mytuhus IAM. albus crassus L. 236. 480 Nerita Amex. e nigro & rubro undata M 12. 432 M.I.alb tenuior I, 2,29 481 Trochus IAM miner alte umbilicatus M 22. 433 M.I. quadratus vulg. M 16.L 368. 482, TIAM min. mar moreus M. 23. 434MI. quadr. min fuscus M17. L367. 483 TIAM maj coftis sinuosis, & c M2c L. 646. 435 M. BARB. ovalis fuscus M. 14. L. 231. 484 Vermiculus BARB. tortilis fuscus. 436 Lunata Anteg. lævis mac.pene 4 dratis L. 270. 485 Patella Amer. alba rugosa.M.3. 437 L IAM albida, leviter fasciata L. 273. 486 Proftrata Barb. alba crafsa rugosa L 544.31. 438 L. IAM. major. fascijs dense elatisM 2.1. L. 286. 487 P. roft. Barb. cancellata parva M. 6. 439 L. IAM. maj fasc plurimis tessellatis M. 22. 488 Concha Veneris Virg. ventre flaves c. &c. M. 14. 440 L.BARB. minor to sellata M. 18. 489 Cylindraceus Amer. albus parvus M. 28 441 L. BARB. minfasc. rarior elatis M 19. 490 C.BARB vulg undis marmoratis M. 19 442 L. BARB min fasc plurimis clatis M 20 L. 278. 191 Olivaris IAM. fasciis eleg. maculatis &c.M. 23. 443 L.BARB. min fasc. latis & clatis 492 Persicoides IAM maculatus M. 29. 444 Bucardites MARIANA lævis. H. Iones. 493 P. Barb fasciatus M. 26. L. 834. 445 Tellina cuneata Anteg fasc & undata T. Grigg. 446 Pectunculus IAM. ex toto hunulatus M. 25 L. 323. 494 P.Barb. fiyew M. 27 495 Buccinum BARB dent fascys nodosis M.32. 447 P.BARB. Compressus, lavis, intus flavus T. Walduck 496 B. Campeach fasciatum crassum M. 13. 497 B. IAM. fasciatum tenue M. 14. 449P.AMER. Compr. eleganter imbricatus M. 26. 450P.BARB. Cancellanus M. 27. L. 337. 498 B. IAM. triangulare M. 15 451P. IAM. minor. quasi cancellatus 499 Murex bilinguis IAM vulo M.3. 452PAMER.cavus, strijs undatis. Carter 500M. IAM. variespinosus M.7. 453P BRASIL alb. undis capill majoribus M. 23. CAP 17. 501M.AMER. valde rugosus M. 8. 454P. LAM. rubesc. undis capill tenuissimis M. 24.C.18. 502 Buccinum ampullac. & c muricatum minus CAR 33. 155 P. LAM. triquetrus, mac. fincis L. 316. 503 Bursa IAM. vulg imbricata M. 20. 456 Purusculus Campeach strijs crassis M 28 L. 347. 504 Per dicea IAM. vulg. M.21. L. 984. 457 Chama IAM minor compr. rubra M 30. 50g P. IAM. nodosa M. 22 L. 989. 458 C. IAN min radiata fragilis W. Brown 506 Unicornu AMER . mwicatum M. 25. 459 C. LAM. alba splendens, radijs rubris Car. 12. 460 C. alba fasciata tenuis Car. 9 L. 308. Explicatio Liter. & c. CAR Acta Philosoph Reg. Societ. LOND. N.º 299. 461 C.CAR tenuis, cunea costatà. L. Lister Hyt. Conchyl. Tabular. No

M. Memoirs for y Curious A.D. 1708. p.97. &c.

462 C. alba, cuneata, fasciam CAR. x1.

463 C. alba cuneata, radijs rubris Stratton







Sutten Nichells souly




ę






.




Sutton Nicholls je




¢















































1.0













Sutton Nicholle Sculp










the off policy of

HANN SHALL









Papilionum BRITANNIÆ Icones, Nomina, &c.

Containing the Figures, Names, Places, Seafons, &c. of above Eighty English Butter-flies, being all that have hitherto been observed in Great-Britain.

By James Petiver, Fellow of the Royal Society, LONDON.

TAB. I. Papiliones BRITANNIÆ albidæ. ENGLISH White Butterflies.

Fig. 1, & 2. Papilio alba, venis nigitis Mus, nost. 301.

White Butterfly with black Veins. Ray Insector. Hist. 115. 3. Its fine black Veins without any Spots, differs it from all the following. Found in Mea-

Spots, differs it from all the following. Found in Meadows about June.

3, & 4. Papilio alba major, apicibus nigris. Great white Cabbage Butterfly. Differs from the next, in having no Spots on its upper Wings, Appears in May, June, &c.

5, & 6. Papilio alba major, bimaculata. Ray 113, 1. Great Female Cabbage Butterfly, Gaz. Nat, 62, f. 3. Muf. noft. 825. These are very common with the last, in Kitchen Gardens where Cabbages grow, on which they lay their Eggs,

7, & 8. Papilio alba media, maculata, Ray 118. 2. Lester white Cabbage Butterfly. Very common in Fields and Gardens most part of the Summer.

9, & 10. Papilio alba media, bimaculata. Lesser white double sported Butterfly. This and the next are found promiscuously with the rest.

o, & to. Papilio alba media, bimaculata. Leffer white double sported Buttersty. This and the next are sound promissionally with the rest.

11, & 12. Papilio alba, media, trimaculata. Lesser white treble sported Buttersty. Differs from the last in having a third spot on the parting of the Wings.

13, & 14. Papilio alba, media, immaculata, Ray 114. 3. Lesser white unspotted Buttersty. The under Wings of these 6 last are frossed below with a pale yellow and without Veins, which the 3 next have,

15, & 16. Papilio alba, media, trimaculata, venis subtus latis, Mus. nost, 302. Common white, veined Buttersty, with 2 or 3 Spots, GAZ. Nat. 62. st. 4. Common with the next in Fields and Meadows most part of the Summer.

17, & 18. Papilio alba, minor, immaculata, venis subtus latis. Lesser, white, veined Buttersty. This has no Spots.

19, & 20. Papilio alba, media, maculata, venis subtus latis. Lesser, white, veined Buttersty. Differs from 15 in having but one Spot on each upper Wing,

21, & 22. Papilio alba minor, apice nigricante. White sin having but one Spot on each upper Wing,

21, & 24. Papilio alba minor, apice nigricante. White sin hampsted and other Woods in June, &c.

23, & 24. Papilio alba minor, Mas. Small white Wood Buttersty. This is sometimes tipt below, but scarce seen above, N. B. I have observed amongst all these white Buttersties, the most spotted to be the Females.

TAB. II. Papiliones Britanniæ, luteæ, albæ, &c. mixtæ. English, yellow, white, and other mixt Buttersties.

Fig. 1. Dapilio Sulphurea, Mus. nost. No. 1. Ray 112. 4. The Brimstone Buttersty. This and the next appear amongst the first in the Spring and again in Autumn.

2. Papilio sulphureus, pallidus, Mus. nost. 2. The Male Straw Buttersty. This being so near white, it often escapes as common.

3 Papilio crocca, limbo niguicante. The Saffron Buttersty. capes as common.

capes as common.

3 Papilio crocea, limbo nigricante. The Saffron Butterfly. Its Ground Saffron, with broad black Edges, feen about Depiford, Peckbam, &c. from June till September.

4. Papilio crocea, limbo nigricante, maculato, Gaz. Nat. 14. f. 11, Rap 112. 6. Spotted Saffron Butterfly, Found with the laft, and differs from it in having a fpotted Edge.

5. Papilio major caudata, ex nigro & luteo variegata, Ray 110. Royal William. Muf. noft. 328. Its Beauty, Size, and Tail differs it from all others. This has been caught about London and divers Counties in England, yet rarely.

6. Papilio albus, lubius viridi colore marmoreatus, Mas, apicibus cioceis, Muf. noft. 306. Ray 115. 6. White, marbied Male Butterfly. Found with the Female, but more particularly observed by its Saffron Tips, by which its easily known flying.

known flying.

7. Papilio alba, subtus viridi colore marmoreata, Famina, Ray 115.7. White marbled Female Butterfly, Mus. nost. 305. This appears in April and May, but not common.

8. Papilio alba minor, subtus viridescens marmoreata. The slight greenish Half-Mourner This is somewhat less than the next, and has fewer Spots, especially above.

9. Papilio Leucamelanos subtus viridescens marmoreata.

ernouns greenish Half-Mourner Mus. nost. 304.Gaz. Nat. 1. fig. 7. Rap 116. 10. This has also been found about Hamp-fied in July or August.

10. Papilio Leucomelanos. Common Half-Mourner, Mus. noft. 3. fig. Ray 116.9. These have near an equal Mixture of white and black, they appear about Midsummer and after,

in moist Meadows.

11. Papilio major nigrescens tricolor, circulo ferè sanguines ornatus. Muf. nost. 327. Ray 12601. The Admiral. A beautiful Fly, and eminently distinguisht by a red List cross the upper Wing. Often seen in Gardens and Fields from the End of July till Autumn.

12. Papilio fusca, areâ & maculis albis. Gaz. Nat. Tab.
f. 12. White Admiral. Found about Dullidge and Wickham near Croyden. as also at Henly upon Thames. TAB. III. Papiliones Britannia maculata.

Brittish spotted Butterslies.
Fig. 1, & z. Papilio Fritislaria major, crozea, lineis subtùs argenteis. Great Silverstreakt Orange Fritislary. Differs from the next in having Streaks above on an Orange Ground.

3, & 4. Eadem Aurea. Great Silver-streakt Golden Fritillary. Mus. nost. 321. Ray 119. 4. The Spots above more and larger, without any black Streaks, on a pale golden Ground. These appear about the beginning of

5, & 6. Papilio Fritillaria major, maculis subtus argenteus. Great Silver-spotted Fritillary. Mus. nost. 320. Ray 119.5. In this the Spots above are larger than the

next, and below the ground between the Spots pale.
7, & 8. Papilio Fritillaria major, subtus viridior, maculis argenteis. In this the Spots are less, and the Ground below greenish. This and the last appear about the below greenish. midst of July.

9, & 10. Papilio Fritillaria, tessellata, serotina, Mus. nost. 323. subtus albida. White May Fritillary. Ray 120. 8.2. Very like the next, but whiter below.

11, & 12. Eadem subtus straminea. Stram May Fritillation. Roch these pretty company in Comp. Word, and differ

Both these pretty common in Cain-Wood, and differ

from the next in being Checker'd above.

13, & 14. Papilio Fritillaria maculata, precox, Mus.
Pet. 322. Ray, 120. 7. April Fritillary. This has many white and Silver Spots below, by which it chiefly differs

from Fig. 18.

15, & 16. Papilio Fritillaria minor. Muf. nost. 324.

Gaz. Nat. 16. f. 10 Ray 122. 12. Vernons, small Fritillary. It's the least of all the Fritillaries yet known. Found in several Woods round London.

17, & 18. Papilio Fritillaria, maculata, præcox, maculis fubrus argenteis paucis. April Fritillary with few Spets. This has but one oval Silver Spot in the midft of each Wing beneath, whereas Fig. 14. has divers. Perhaps Male and Female. Frequent in Cain-Wood.

19, & 20. Papilio Fritillaria media, meandris nigris.

Dandridge's midling Black Fritillary. Ray 121. 10. Mr. Dandridge has observed this and the next about the end of May and beginning of June in Cain Wood and the Oak

of May and Deginning of June in Cain wood and of Honour Woods near Dullidge.

21, & 22, Papilio Fritillaria minor, meandris nigris.

Small black Fritillary. Ray 121. 11. Differs chiefly in being less than the last.

23, & 24. Papilio Fritillaria minor, maculis argenteis subtus perbelle notata Mus. nost. 520. Lesser Silver-spotted or Riga Fritillary. Ray 120. 6. Observed about Cambridge 25, & 26. Papilio Fritillaria minor, undulata, areis subsection of the White Dullidge Fritillary.

rus albis. White Dullidge Fritillary. Found in the Wood, thereabouts in May.

The self of the se

A STATE OF THE STA

monly the Elm.

3, & 4. Papilio Testudinarius minor, Mus. nost. 316. Lesser or Common Tortoile-shell. Ray 117. 1. Very frequent all Summer, and often found in Houses in the

frequent all Summer, and often found in Monges in the Winter.

5, & 6. Papilio Testudinaria, Comma dicta, subtus susca. The Silver Comma. Ray 118. 3. This and the 3 fol lowing appear about Midsummer.

7, & 8. Eadem subtus pallidior. The pale Comma. This below is of an Oker Marble, and paler than the last.

9, & 10. Eadem, alis magis laceratis. Jagged wing'd Comma. These Wings are deeper cut and more vivid; it's finely marbled underneath, with small geeenish Eyes speckt with black. speckt with black.

11, & 12. Eadem minor, Small Comma. It's very dark

below, and in all Parts lefs.

13, & 14. Papilio minor aureus, ex nigro permaculatus Mus. nost. 317. Ray 125. 20. Small Tortoise shell. In Meadows and about Hedges from July till Autumn.

15, & 16. Papiliunculus fuscus, pundtis paucioribus albicantibus, Gaz. Nat. Tab. 36. f. 9. Ray 132. 20. Brown Marsh Fritillary. The Spots in this are larger and sever than the next, and Found with it.

17, & 18. Papiliunculus suscus, pundtis plutimis albicantibus Mus. nost. 325. Ray 133. 21. Gaz. Nat. 36. f. 6. Small-spotted brown Marsh Fritillary. The Spots in this are smaller and more distinct.

19, & 20. Papilio niger suscus Hampstediensis marmorea-

19, & 20. Papilio niger fuscus Hampstediensis marmoreatus, Gaz. Nat. 36. f. 3. Ray 132. 18. Handley's small brown Buttersty. It's brown above and paler below, and dully marbled.

21, & 22. Papilio eleganter variegata agilis, Bella donna dicta Muf. noft. 326. Ray 122. 13. Painted Lady. It usually settles on Banks and dry Ground.

23. Papilio minor fuscus, subtus striatus, Gaz. Nat. Tab.

11. f. 9. Ray 129. 8. The brown Hair-streak. Mr. Ray has seen this and 26 in Coitu on Nettles, July 8. 1692.

24. Idem agro aureo. Gaz. Nat. 11. f. 11. Ray 130. 10. Golden Hair-streak. Found with the last about Croydon.

25. Pars inferior. This Figure represents the Under-side of the 2 last, which differ not below.

of the 2 last, which differ not below.

26. Popilio minor carulescens, subtùs striatus, Mus. nost.

319. Ray 130. 9. Ray's blew Hair-streak. The Ground above is a dark hair brown, with a blewish Cast

27. Idem fuscus, agro caruleo. Our blew Hair-streak. The Ground the same with the last, but the Inside of its upper Wings has a broad Field of vivid blew.

28. Pars inferior. The 2 last are the same below.

TAB. V. Papiliones Britannia Oculatæ.

British Eye-wing'd Butterslies.

Fig. 1. Papilio Oculus Pavonus dictus Mus. nost. 314.

The Peacock's Eye. Ray 122. 14. This often appears early and continues long, it's pretty common both in Gardens and Fields.

in Gardens and Fields.

in Gardens and Fields.

2. Papilio Oculatus Hampstediensis, ex aureo suscess. Albin's Hampsted Eye. Where it was caught by this Curious Person, and is the only one I have yet seen.

3. Papilio oculis nigris, subtùs marmoreus, Mus. nost. 307.
Ray 128. 6. Gaz. Nat. Tab. 14. f. 9. The Tunbridge Grayling. Because first caught in those Parts. This I take to be the Female, and appears about Midsunger. be the Female, and appears about Midfummer.

4. Idem obscurior. Brown Tunbridge Grayling. Much darker on the upper side and somewhat smaller.

5. Papilio oculatus, è fusco & flavescente mixtus, Gaz. Nat. Tab. 24. fig. 3. opt. Cat. 23. The Enfield Eye. It being the Place I first observed them in. These appear

from April till July.

6. Idem obscurior & serotinus. Brown Ensield Eye. Mr. Dandridge first observed this about Waiford towards the End of July.

7. Papilio Oculatus, ex aureo & fusco marmoreatus, Mus. nost. 312. Ray 123. 5. The London Eye. These I

TAB. IV. Papiliones Britannia, maculatæ & striatæ.

British spotted and streaks Butterssies.

Fig. 1, & 2. Apilio Tessudinarius major. Mus. nost.

315. Great Tortoise-shell Butterssy. Ray

118. 2. A large Fly, I have observed them both in

Spring and Autumn, they often settle on Trees, and commonly the Elm.

have met with from May till August, but not very constitute.

8. Idem area success. London Fye, with a brown Lift.

Found with the last, and may be Male and Female.

9. Papilio pratentis Oculatus aureus, Mus. 1981. 309. Golden Meadow Eye, Ray 124. 17. Common in Meadows before Moving. and perhaps Female to the next. 1001.

8. Idem area fusca. London Fye, with a brown List. Found with the last, and may be Made and Female.

9. Papilio pratentis Oculatus aureur, Mus. wist. 309. Golden Meadow Eye, Ray 124. 17. Common in Meadows before Mowing, and perhaps Female to the next.

10. Papilio prat. Oculatus suscur, Mus. nost. 319. Ray 124. 16. Brown Meadow-Eye. Found frequently with the last in May and June.

last in May and June.

11. Idem fegutenre, area susca. Differs from the next in having a brown List. Perhaps Male and Female.

naving a brown List. Perhaps Male and Female.

12. Papilio aureo-fuscus minor Oculatus, pupillis geminis, Mus. nost. 310. Ray 124. 18. Hedge-Eye with double Specks. Seen about Hedges in August.

13. Papilio med. omnino suscus, plurimis Oculis nigris in circulis luteis subtùs ornatus, Mus. nost. 313. Ray 129.

7. Brown & Eyes. These rarely appear before August, and mostly near Rivers.

14. Idem 7 Oculis. Brown 7 Eyes. This has often underneath one Eye fewer than the last.

15. Golden Heath-Eye. Papiliunculus aureus oculatus in Ericetis frequens, Mus. nost. 311. Ray 125. 19. Very common on Heaths, from May till Autumn.

16. Idem margine susce. Selvedg'd Heath-Eye. Differs

from the last, in having a brown brim or edge and found

with it. TAB. VI. Papiliunculi Britan. carulescentes, aurei, Esc. Small English, Blue, Golden and Brown Butterslies.

Fig. 1. Papiliunculus cærulescens, limbis nigris, Gaz. Nat. 35. s. 2. Pale blue Aigus. I sirst observed this in Thickets near Banstead Downs, about July.

2. Idem, lineâ subtus nigrà. Nicely differs from the last, by a small line underneath.

2. Papiliunculus caruleus guident subscriffinges. Gaz. Nat. 25.

3. Papiliunculus caruleus vulgatissimus, Gaz. Nat. 35. f. 1. Blue Argus, Mus. nost. 318. Ray 131. 11. Very common on Heaths from June till August.

4. Papiliunculus caruleus, limbis nigris. Selvedg'd blew Argus. Found with the last, but more rare.

5. Papiliunculus è cæruleo fuscus. Gaz. Nat. 35. f. 3. Mixt Argus. It's brown above mix'd with blew.

6. Papiliunculus carulescens minor, limbis nigris. Disfers from fig. 4. in being less and with smaller eyes.

7. Papiliunculus è suscendentes. This has sewer Spots underneath than fig. 5.

8. Papiliunculus plumbeus. Lead Argus. Eyed underneath like the server ing. and sound with these

neath like the foregoing, and found with them.
9. Papiliunculus plumbeus parvus. Small Lead Argus.

Less than the last. 10. Papiliunculus fuscus, marginibus aureis, Gaz. Nat. Tab. 35. f. 4. Edged, brown Argus. Its ground darker than the 2 last.

than the 2 last.

11. Papiliunculus carulescens, subtus punctatus. Blue speckt Buttersty. I have observed this on Holly Trees.

12. Papiliunculus casuleus, apicibus nigiis, subtus puncta-

Blue speckt Butterfly, with black tipps. Found with

13. Papilio minor supernè fuscus, infernè viridis. Holly uttersty. Because I sirst observed it on that tree.

Butterfly. Because I first observed it on that tree.

14. Papilio minor aureus, immaculatus, Gaz. Nat. 34. 9.

in Defcript. The fpotlefs Hog. Ray 125, 22.

15. Papilio minor auteus, flria nigrà distinctus, Gaz.

Nat. 34. f. 9. Ray 125, 21. Streakt Golden Hog.

16. Papilio minor ex aureo & fusco mixtus, flria nigrà

Gaz. Nat. 34. f. 8. Streakt cloudy Hog. Differs from the last in being clouded with brown.

17. Papilio minor ex aureo & fusco mixtus, Gaz. Nats 34. f. 7. Cloudy Hog. This has no Slants. All these Varieties are found in the Woods, &c. about Hampstead.

Ab Autore nuper editi.

1. petiveriana 1. 2 & 3. five Collectanea NATURA, Domi Forisque Autori Communicata, Lond. Fol.

2. Hortus Peruvianus Medicinalis: or the South Sea Herball. Illustrated with many curious I'ig res of divers Medicinal Herbs, Gum-Tiees, &c. growing in those Parts with their Virtues and Use. Lond. Fol.


HUS PLANTEN


Papiliones BRITANNI F. albidæ ENGLISH Wnite BUTTERILIES THE . Here , co to per heily Somer, 1715;




Papiliones BRITANNIE Lutea Alba & Mixto.


LNGLISH Yellow, White & other mixt-BUTTERFLIES.








A Tempson, wet or South, Lendon, .



Papiliones BRITAN. maculativ & striate .ENGLISH Spotted & Streakd BUTTERFLIEN H. Tenafson, chin or Side.




Papiliones BRITAN. Oculatee, BRITTISH Eye wing'd Butterflies. TAB. V





Papiliunculi BRITANNIÆ cærulescentes, aurei &c TABN1. Small ENGLISH, Blue, Golden & Brown Butterslies.



Sullen Val. //



MONSPELII Desideratarum Plantarum CATALOGUS.

E R A 10, Magnolo, Chabræo, C. & I. B. &c.

A Jacobo Petiver, Societ. Regal. Soc. LONDINI.

Acanthium Seriphium Gallicum CB. Fl. Aug. R. 370.15.

Acanthium Illyricum Park. 979. f. 2. Ray 373; 21

Acet Monfielfulamm Lugd. 95. f. 3. Ray 1701. 3.

Acet Monfielfulamm Lugd. 95. f. 3. Ray 1701. 3.

Actofu matin, fol. finuatis & dentatis Bot. Monfip. 4. R.175; dettofu matin, fol. finuatis & dentatis Bot. Monfip. 4. R.175; dettofu matin. fol. Chabr. 579. H. 5. R. 82. r.

Aleyonium 12 five Burfa matina CB. Chabr. 570. E. I. R.83. 8.

De longiffimo, lato, tenuique folio B. pr. 154. R. 74. 21

Aleyonium 12 five Burfa matina CB. Chabr. 570. E. I. R.83. 8.

De longiffimo, lato, tenuique folio B. pr. 154. R. 74. 21

Allium f. Moly mont. 2 Clul. 194. 65. Pl. Jun. R. 1118. 6. Var.

Allium f. Moly mont. 5 Clul. 194. 65. Pl. Jun. R. 1118. 6. Var.

Allium f. Moly mont. 5 Clul. 194. 65. Pl. M. 1118. 6. Var.

Allium f. Moly mont. 5 Clul. 194. 65. Pl. M. 120. 5. Fl. I. & Y.

Allium f. Moly mont. 2 Clul. 194. 65. Pl. M. 120. 67. Fl. Maio.

Alia f. f. Moly molecutem capillaceo folio CB. pr.28. Fl. Maio.

Alia f. f. Moly molecutem capillaceo folio CB. pr.28. Fl. Maio.

Alia f. f. Moly molecutem capillaceo folio CB. pr.28. Fl. Maio.

Alia f. f. Moly molecutem apillaceo folio CB. pr.28. Fl. Maio.

Alia f. f. Moly molecutem apillaceo folio CB. pr.28. Fl. Maio.

Alia f. f. Moly molecutem apillaceo folio CB. pr.28. Fl. M. 20.

Alia f. f. Moly molecutem apillaceo folio CB. pr.28. Fl. M. 20.

Alia f. f. Moly molecutem apillaceo folio CB. pr.28. Fl. M. 20.

Alia f. f. maior CB. Thurnigiaca CA. M. H.12. f. 67. 60. 22.

Alia f. f. maior CB. Thurnigiaca CA. M. H.12. f. 67. 60. 22.

Alia f. f. maior CB. Thurnigiaca CA. M. H.12. f. 67. 60. 23.

Antine apilla gubara five. Cheele I. B. Chabr. 108. I. 6.

Anonis vilcola fpinis carens, lutea latif BM. 21. Fl. Aug.

Anonis vilcola fpinis carens, lutea minor BM. 21. Fl. Aug.

Anonis vilcola fpinis carens, lutea minor BM. 21. Fl. Aug.

Anonis vilcola fpinis carens, lutea minor BM. 21. Fl. Aug.

Antirchium maior cutudiore fol. CB. Chi. Fl. M. 20.

Antirchium max. Seriphii folio 24. ft. B. pt. No.

77 Centaurium minimum purp. ramofum BM. 57. Fl. Jul.;
78 Centaurium pufillum luteum CB. R. 1092. 4.
79 Chamamelum leucanthenium magno flore CB. pr. 70. 1.
80 Chamamelum inaritimum Dal. Lugd. 1394: f.1. Ad.; 45.f.
81 Chelidonia rotundif. major B. pr. 137. R.580. 2.
82 Chondrilla vificola, caule foliis obducto CE. Fl. Aug. &c.
83 Chondrilla viminea Monspeliaca B. pr. R. 223. 9. (f. 1.)
84 Chondrilla pufilla marit. lutea, bulbosa Ad. 83. f. Fl. Maio.
85 Chondrillis affinis quædam laciniata BM. 64. R. 227. 2.
87 Eadem minor BM. 65. Fl. Apr. circa Gigean.
88 Cirstum fol. non hirsutis, fl. compactis CB. Chabr. 346: 3.
89 Cistus mas 1. Clus. 68. f. 2. fol. oblongo incano CB. Fl. A.
90 Cistus Salvia folio CB. fem. Clus. 70, f. (& Maio.)
91 Eadem vingis humisparsis BM. 67.
92 Cistus Ladanifera Monspel. CB. fol. latioribus BM. 67.
93 Cistus Ladanifera Monspel. CB. fol. latioribus BM. 67.
94 Chamacistus v. Clus. 75. f. 1. fol. minor. & incanis CB.
95 Chamacistus v. Clus. 74. f. 2. fol. Thymi incanis CB.Fl. Jun.
96 Helianthemum fol. Thymi incano I.B. Chabr. 103. 3. R. 1016.11.
97 Helianth. fol. Sampsuci capit. valde hirsuris I.B. Ch. 101. 14.
98 Chamacistus 2. Clus. 73. f. 2. repens Serpillifolia lutea CB.
99 Chamacistus 2. Clus. 73. f. 1. fol. Myrtiminoris incanis CB.
100 Cistus. ann. fol. Ledi, fl. luteo I.B. Ch. 90. f. 6. R. 1013. 4.
101 Clematis maritima repens B. pr. 135. R. 621:3. Var. (Fl. Oct.
102 Clematis maritima repens B. pr. 135. R. 621:3. Var. (Fl. Oct.
103 Clematis parva, Montibus propè Castelneuf I.B. 1. 15. p. 127.
104 Convolvulus Spicæfolius argenteus, 12d. repente BM. 74.
105 Convolvulus Spicæfolius argenteus, 12d. repente BM. 74.
106 Conyza caul, rubent, tenuioribus fl. luteo nudo BM. 76. f.
107 Conyza tuberofa lutea &c. BM. 77. marina Lugd. 1366. f. 1.
108 Coronopus Serpentina I.B. Chabr. 500. f. 2. R. 880. 8.
110 Coronopus Serpentina I.B. Chabr. 500. f. 2. R. 880. 8. 107 Conyza Inberofa lutea &c. BM.77. marina Lugd.1366.f. 1.
108 Conya Helenitis mellita incana Lob. Ic. 347. R.263.1c.
109 Corallina fruticofa recta alba CB.
110 Coronopus Serpentina I.B. Chabr. 500. f. z. R.880.8.
111 Crocus verms angustif. 2. Clus. 207. s. 2. Fl. M. & Apr.
112 Cyanus peramatus supinus, capit. squarosis &c. 4d.236. Fl.
113 Cyclamen Byzantinum Clus. 263. Fl. Mart.
114 Cynoglosium Cret. angusto argenteo fol. CB.
115 Cytisus fol. augustis quasi complicatis CB. Fl. Maio.
116 Daucus glauco follo, similis Fanienlo tortuoso I.B.
117 Dens Leonis latiore & rotundiore folio. BM. 85.
118 Dens Leonis minor fol. aspero B. pr.63. R. 246: 113
119 Echium mag. & asperius st. albo CB. Fl. Y. & Aug.
120 Elatine fol. acuminato st. cæruleo CB. Fl. Aug.
121 Erica max. alba CB. pnima Clust. 41. st. R. 1714.5.
122 Erica 2 altera Clüs. 42. f. R. 1715. 10.
123 Erica 3 Clus. 42. f. R. 1715. 10.
124 Erica 5 Clus. f. I. R. 1715. 10.
125 Erica 15(1) 42. f. I. R. 1715. 10.
126 Euphrasia purp. minor BM. 94. stg. B. pr. Fl. Maio.
127 Euphrasia purp. minor BM. 94. stg. B. pr. Fl. Maio.
128 Fast sylv. f. 101. atto CB. Chabr. 140. 6:
129 Fesula fam. Plin. CB. R.420. 1.
130 Ficus sylv. Diose. Captificus Plin. CB. R. 1433. 2:
131 Fanumgracum sylv. Dal. Lugd. 481. f. Fl. M. & Jun.
132 Fanumgracum sylv. Dal. Lugd. 481. f. Fl. M. & Jun.
133 Fucus sensus Faniculaceus BM. 100. Chabr. 571. f.
134 Fucus sensus Faniculaceus BM. 100. Chabr. 571. f.
135 Fucus surs santus Faniculaceus BM. 100. Chabr. 571. f.
136 Funaria tenuissims foliis, s. Brigoule vulgo.
138 Gensstella spinosa Monssel. B. pr. 157. R.1728.3.
149 Geranium Columb. tenuius lacinistum B.pr. Fl. Vere.
141 Geranium Petræum Cicuta fol. rad. crassa BM. sp. 103.
142 Gnaphalium Roseum B. pr. 122. f. J. Maio.
143 Gnaphalium minus repens CB. Lob. lc. 481.2. Fl. Maio.
144 Gramen Canimum arvense CB. Lob. lc. 202. Fl. 4. f. 2.
145 Gr. Canimum marit. asperum B. pr. 2. f. 2. Fl. Aug.
147 Gr. Tremulum minus panicula parvà CB. pr. 4. f. 2.
148 Gr. Tremulum minus panicula parvà CB. pr. 4. f.

```
152 Gr. Aispecuros minor spica longiore CB. R. 1266. 13.
153 Gr. spica subrotunda echinatá B. pr. 16. f. R. 1268. c.10
154 Gr. tomentosum spicatum CB. Lugd. 430. f.
155 Gr. Paniceum, spica simplici CB.
     156 Gr. Arundinaceum spica multiplici CB.
     157 Gr. Arundin. marin. maximum Ad. 4
    157 GI. Arunam. maxim maximum Aa. 4.
158 Gr. Arunam. maximum plumofum album CB. R. 1280. 2.
159 Gr. Cyperoides latif. spica spadiceo viridi majus CB.
160 Gr. Cyperoid. panicula sparia minus CB.
161 Gr. Cyperoides maritimum CB. Ad. 44. f. R. 1298. 17.
162 Gr. Cyperinum nemorosum Tab. R. 1298. 16.
163 Gr. Sparraum Manshel Cappillaceum fel. minimum CB.
  162 Gr. Cyperinum nemorosum Tab. R. 1298. 16.
163 Gr. Sparteum Monspel. s. capillaceum fol. minimum CB.
164 Gr. Loliaceum minus, spica simplici CB. R. 1264. 6.
165 Idem minimum BM. 120. Luco Gramuntio.
166 Gr. Loliace. minus supinum, spica multiplici CB.
167 Gr. Loliacea panicula ramola, maritimum BM. 120.
168 Gr. Phanicoides fol. convolutis &c. argenteis BM. 120.
169 Gr. Spica Briza majus & minus CB. R. 1257. c. 2.
170 Gr. Tremulum Monspel. BM. 121. pl. 6. CB. Theatr.
171 Gr. Aven. max. utriculis lanugine alba, arist. longissimis BM. 121. sie. Fl. Maio. R. 1292.12.
    BM. 121. fig. Fl. Maio. R. 1299.12.

172 Gr. f. Festuca longissimis aristis CB. R.1290.13.

173 Gr. Aven. locustis rubris montanum B pr. 20.f.

174 Gr. Aven. mont. lanuginosum CB. R. 1269. 5.

175 Gr. Aven. lanug glumis varioribus R. pr. 20. f.
  174 Gr. Aven. mont. lanuginofum CB. R. 1269. 5.
175 Gr. Aven. lanug. glumis rarioribus B. pr. 20. f. 2.
176 Gr. rubrum Monspeliense I.B.
177 Gr. aliud Monspeliense I.B.
178 Gr. aliud glabrum Monspeliense I.B.
179 Gr. supinum Monspeliense Laugerii Lob. illustr.
180 Gratiola s. Hyssopisoia angustioribus foliis CB. pr.
181 Grossularia non spinosa BM. 123.
182 Stæchas citrina altera Ad. 203. f. R. 283. 11.
183 Stæchas citrina altera inodora Ad. 204. f. R.283. 10.
184 Helleborine sl. albis I.B. Fl. Malo.
   183 Stæchas citrina altera inodora Ad. 204. f. K.283. 10.
184 Helleborine fl. albis I.B. Fl. Malo.
185 Helleborine 2 Clus. 272. mont. angust. purp. CB. Fl. Maio.
186 Helleborine 3 Clus. 273. f. angust. prat. CB.
187 Hesperis sylv. fl. parvo B. pr. 103. 1.
188 Hieracium pulchrum I.B. Fl. æst. tota. R.234. 17.
189 Hieracium minimum salcatum B. pr. 64.13.
190 Hieracium D. Leonis solio bulbosum B. pr. 63. 5.
191 Hieracium minus D.L. fol. subaspero CB.R.229.1. (B. pr. 102 Hieracium D. L. fol. hirsutie asperum minus laciniatum
    192 Hieracium D.L. fol. hirsutie asperum minus laciniatum
193 Hieracium parvum hirsutum, caule aphyllo I.B.R.246.2.
  194 Hieracium 1. latifol. Cluf. 139.f.R.239.7.
195 Hieracium muror. angust. laciniatum minus CB.
196 Hieracium capillaceo folio B. pr. 66. 21.
197 Holosteum Monspeliense Ad. 18. f. R. 880.6. c.7.
198 Hyacinthus comosus minor Ad. Fl. F. & Mart.
 Hypercoum CB. Fl. Apr &c. R.1328. 1.

200 Hypercoum fupinum tomentosum minus vel Monspel. CB.

201 Jacea squammis pennatis s. cap. villoso I.B.

202 Jacea mont. incana capite Pini CB. Fl. Jun. R.329. 29.

203 Jacea mont. Narbonensis Lob. Obs. 253. R.329.3c.

204 Jacea incana Lugd. 1194. f. Fl. Jun. Ad. 235. f.

205 Jacea latis. pinosis capit. pulchris, sl. purp. I.B.

206 Jacea capitulis spinosis Ad. 234.

207 Jacea lutea capite spinoso CB. Fl. Jun. &c. Ad. 234. f.

208 Jacea Melitensis capitulis conglobatis Boc.

209 Jacobea 3. latisolia 1 Clus. 24. f. Fl. Jun. Jul. &c.

210 Jacobea Senecionis solio H. Bles. Fl. Maio.

211 Iberis latiore folio CB.

212 Iberis Nasturtii solio CB. Fl. Jun. R.827.1.
      199 Hypecoum CB. Fl. Apr &c. R.1328. 1.
    212 Iheris Nasturtii folio CB. Fl. Jun. R.827.1.
212 Iheris Nasturtii solio CB. Fl. Jun. R.827.1.
213 Ilex solio Agrisolii BM. 140.
214 Ilex media Coccisera Ilici planè suppar, sol. Aquisolii Ad.
215 Iris maritima Narbonensis Ad. 32. s. maj. & minor.
216 Iris perpusilla saxat acaulis serine Ad. 32. s. R. 1187. 3.
217 Iris minor lutea Dod. 244. Fl. primo l'ere. R.1188. 6.
218 Juncus acumine restexo majore CB. R.1304. 11.
219 Juniperus major baccà sussente CB. R. 1413. 2.
220 Kali minus villosum CB. R. 199. 15.
221 Kali minus fol. lucidis BM. 146.
222 Lathyrus luteus latisolius BM. 150. Fl. Apr. &c.
223 Lathyrus angustisolius humilior BM. 150.
223 Lathyrus angustifolius humilior BM. 150. (150. 224 Lathyrus humilis fol. Gramineis venosis, st. pheniceo BM. 225 Lathyrus angustissimo s. capillaceo fol. B. pr. st. Apr. 226 Lathyrus maj. Narbon, angustifolius I.B. Chahr. 145. f. 3. 227 Leucoium marit, angustifol. CB. Fl. Vere & astate Ad.
                                   141. f.
 228 Libanotis Cachryophoros quibusdam fl. luteo I.B. Fl. Mag.
229 Libanotidi congener BM. 154. Lugd.
230 Limonium supin. reticulatum Bocc. Fl. Sept. R. 396. 9.
231 Limonium parvum Bellid. min. folio CB.R. 395. 4.
232 Limon, marit, minus, fol. cordatis B. pr.99.1.
233 Limonium minus bullatis fol. BM. 156. f. Fl. Jun.
234 Linaria arvens. carulea erecta B. pr. 107. 5. Fl. Apr. 6
235 Linaria &c. fol. imis rotund. BM. 158. f. Fl. Maio.
```

```
236 Linaria lutea, parva annua I.B. Fl. Maio. R. 755. 12, 237 Linaria folioso capitulo CB. Fl. Sept. R. 292. c. 16, 238 Limum maritimum CB. lutenim Fl. Aug. R. 1074.17.
       239 Linum marit. majus I.B.
      240 Linum sylv. minus, luteo flore CB. Fl. Jul.
241 Linum luteum palmare BM. 160. Fl. Jul.
242 Linum angustifolium Leb. Obs.
     243 Linum angustiosium Lov. Ooi.
243 Linum sylv. st. albie Lob. Obf. Fl. Jun. &c. R. 1073.11.
244 Linum sylv. carul. fol. acuto B. pr. 107.2. Fl. Jun.
245 Linum minimum fellatum B. pr. 107.2. BM. Fl. Vere.
246 Linum sylv. luteum fol. subrotundis CB. Fl. Jun.
247 Linhosvermum Linuxis fol. Manshel CR. Fl.
     247 Lithospermum Linaria tol. Monspel. CB, Fl. tota aft. Lob.
                                        Ic. 411. 2.
    1c. 411. 2.
248 Lotus corniculata marit. hirfuta BM. 164. Fl. aft.
249 Lotus 5 phyllos minor hirfut. fol. angustiss. CB. Fl. Aug.
250 Lotus pentaphyllos maj. hirfutis candicans CB. Fl. Maio.
251 Lotus marit. lutea siliquosa, fol. pingui glabro BM. 165.
252 Lychnis sylv. angust. calic. turgidis striatis CB. Chabr.
    446. f. 1.
253 Lychnis fylv. lanuginofa minor CB. Fl. Maio.R.995.11.
254 Lychnis fylv. minima fl. muscoso BM. 171. Fl. Jul. &c.
255 Lychnis minima rigida Cherleri I.B. Fl. Vere. Chabr. 445.
   f. 1.

256 Lychnis Alpina vel Ocymoides repens mont. CB. Fl. Maio.

257 Lycium Gallicum I.B. Fl. Maio. R. 1626. r.

258 Marrubium nigrum longifol. CB. Fl. Ap. M. & Jun.

259 Marrubium album villofum B.pr. 110.2. R.557.6.

260 Marrubium paluft. hirfurum CB. R. 535. 2.

261 Meliloto vulg. affinis, fol. min. & rotundiore I. B.

262 Millefolium odorat. minus Monfpel. H. Blef.

263 Morus finctu albo parvo BM. 178.

164 Morus fr. nigro parvo BM. 178.

165 Muscus terr. Coralloides erectus, cornibus rusescentibus CB.

166 Muscus Gapillaceus longissimus CB.

167 Muscus fugax membranaceus pinguis BM. 180.
   Muscus Capillaceus longissimus CB.

Muscus fugax membranaceus pinguis BM. 180.

Muscus fugax membranaceus pinguis BM. 180.

Narcissus angustif. albus minor B. pr. Fl. Apr. (BM. 182.

Nascussus pincifol. palmaris, parvo st. luteo, odore Mellis

Nassusus pincifol. palmaris, parvo fl. luteo, odore Mellis

Nassusus pincifol. palmaris, parvo st. luteo, odore Mellis

Nassusus pincifol. palmaris, parvo st. luteo, odore Mellis

Nassusus pincifol. palmaris, parvo st. luteo, odore Mellis

Nassusus pincifol. Pil. Jun.

Nymphea alba minor CB. Fl. Jun. 8ec.

Pun. 8ec.

Nassus CB. Lugd.

Nassus CB. 163.

Nassus CB. 163.

Ampilia CB. Lugd.

Amellau.

Oliva media oblonga angulosa Amygdali forma BM. 187.

Oliva media oblonga angulosa Amygdali forma BM. 187.

Oliva media oblonga angulosa BM. 187. Ouliviere.

Oliva minor oblonga id. Pichouline.

Oliva major subrotunda id. Ampoulau.
     280 Oliva major subrotunda id. Ampoulau.
281 Oliva media rotunda viridior BM. 190. Verdale.
282 Oliva media rotunda præcox id. Mourau.
283 Oliva minor rotunda racemosa id, Bourau.
283 Oliva minor rotunda racemosa id, Bouteillau.
284 Oliva minor rotunda rubro-nigricans id. Salierne.
285 Oliva minor rotunda, ex rubro & nigro variegata id. Pi-
286 Onobrychis fructu echinato minor CB. Fl. Jun. (gau.
287 Onobrychis clypeata aspera minor B. pr. 149. 1. Fl. Jun.
288 Polygala minus saxatile Chabr. 153. f. 3.
289 Ophioglossum minus subrotundo solio B. pr. 150. 2.
290 Cynogorchis latifol.spica compacta CB. Fl. Maio. R. 1214.9.
291 Orchis odore Hirci minor CB. Fl. Maio. R. 1213.2.
292 Orchis Muscam referens major & minor CB. Fl. Maio.
293 Orchis Muscam referens lutea CB. Fl. Maio. (Maio.
294 Orchis Palmata angust. minor, sl. odoravistimo B. pr. Fl.
295 Origanum Monsp. pulchrum Cam. 468. s.
296 Ornithogalum majus Monspel. Clus. 187. R. 1151. 2.
297 Ornithopodio assinis hirluta, Scorpioides leguminosa dd.
Fl. Maio.
                                     Fl. Maio.
    298 Ornithopodio affinis hirfuta, stella leguminosa CB. Fl. Maio.
   299 Eadem flor. & filiq. cauliculis adhæiens BM. 195. Chabr.
 156. f. 5.
360 Orobanche inagna purp. Monspel. I.B. Fl. April. &c. Clus. 301 Orobanche subcæruleo flore s, 2. Clus. 271. Fl. Maio. 302 Orobanche minor purpureu flor. s. ramosa I.B. Fl. Apr. 303 Panax Costinum CB. R. 410. c. 3.
304 Panax Heracleum fronde Passinace Ad. 312. R. 400.5.
305 Panicum Dod. 507. f. 1. panicula minore CB. R. 1247. 1.
                                     156. f. 5.
  706 Papaver Cornic. phæniceum hirfutum CB. R.857.13.
307 Papaver Cornic. Violaceum CB. Lob. Ic. R.857.14.
308 Paflinaca tenuif. marin. fol &c. lucidis BM. 199.
308 Paftmaca tenuit. marin. fol &c. lucidis BM. 199.
309 Peplus minor I.B. 3. l. 34. p. 670. R. 869. 28.
310 Phyllyrea latif. spinosa CB. Fl. Vere. Clul. 51. s.
311 Phillica elatior CB. Clus. 50. f. Ad. 421. f. R. 1608. r.
312 Phillica humilis & quasi repens BM. 203.
313 Pilosella maj. erecta CB.R.241.16.
314 Pimpinella Sanguisorba minor, sem. majore & crassiore
315 Pinus sylv. altera maritima Lob. Obs. R. 1400.7.
 316 Pinaster humilis Lob. Obs. R. 140c. 6.
                                                                                                                                                                                                                                      317 Plantage
```

```
117 Plantago max. tota glabra B. pr. 97.1. R.877. 3.
118 Plantago capitulis Lagopi longioribus BM. 206.
119 Polium Lavandula folio CB. Fl. Jun. &c. R. 525. 6.
120 Polium marit. erect. Monfp. CB. R.524.1.
121 Polium mont. album CB. R.524. 2.
122 Polygala acutioribus fol. Monfpel. CB. Fl. Maio.
123 Polygala erecta angustifolia &c. BM. 208. Fl.M. & Jun.
124 Polygonum baccif. marit. minus CB. (Chabr. 457. f. 6.
125 Polygonum Ater. publilo vermic. Serpilli fol. Ad. 80. f.
126 Polygonum angustissimo Gramineo fol. erectum BM. 211.
127 Polygonum angustissimo Gramineo fol. erectum BM. 211.
128 Polygonum muscosum minimum Bocc. Fl. Maio. R. 346
               Fl. Jul. &c.

128 Polygonum angustissimo Gramineo fol. erectum BM. 211.

Fl. Jul. &c.

128 Polygonum muscosum minimum Bocc. Fl. Maio. R. 216.

139 Polytrichum s. Tricomanes minus & tenerius CB.

130 Porum sylv. Vinearum CB. R. 1226. c. 6.

131 Potamogiton angustisol. racemosum B. pr. 101. s.

132 Potamogiton minimum capillaceo fol. B. pr. 101. s.

133 Potamogiton capillac. capitulis ad alas trisidis B. pr. 101. s.

134 Quinquefolinm minus repens luteum CB. 325.

135 Ranunculus magnus valde hirsutus I.B. R. 582. s.

136 Ranunculus magnus valde hirsutus I.B. R. 582. s.

137 Ranunculus palust. echinatus B. pr. 95. s.

138 Ranunculus villossissimus Monspel. B. pr. 69.11.

139 Ranunculus saxat. magno store B. pr. 69. s.

141 Ranunculus Cherophyllos Asphod. radice CB. Fl. Maio.

142 Ranunculus Cherophyllos Asphod. radice CB. Fl. Maio.

143 Ranunculus Coronopisolius, cap. Alopecuri BM. 218. Fl. Vere.

144 Ranunculus memoros. Anemones store minor B. pr.

145 Ranunculus minor, fol. incisis CB. Fl. Apr. &c.

146 Rapunculus minor, fol. incisis CB. Fl. Apr. &c.

147 Refeda fol. Calcitrapa, fl. albo H. Bles. Fl. Maio

148 Phyteuma alter Monsp. fol. minus laciniato Lob. illust.

149 Rbamnus cort. albo Monspel. 1.B. R. 1592. s.

150 Rosa Pomifera major CB. R. 1472. 17.

151 Rosa Pomifera major CB. R. 1472. 17.

152 Rosa foinosissima pumila fl. rubro I.B.

153 Rubia svy. aspera CB. R. 480. 2.
548 Piptenma alter Monp. 101. minus seniato Los., Hait. Ramon Ramano Los., Hait. Ramon Ramon Lac., Hait. Ramon Ramon Lac., Hait. Ramon Lac
```

```
Trifolium globofum repens CB. Chabr. 162. f. 2. R.945.13.
404 Trifolium capit. glomerato rigido I.B. Chabr. 162. f. 4.
405 Trifolium cap. fpumofo aspero min. B. pr. 140.6.R.947.22.
406 Trifolium argentatum fl. luteis IB. Chabr. 159. f. 6.
407 Medica magna turbinata I.B. Chabr. 162. f. 2.
408 Medica fr. compresse RM 268
    408 Medica fr. compresso minore BM. 268.
    409 Medica hirsuta, echinis rigidioribus I.B. Chabr.
   Trifolium Cochleatum polycarpon CB. R.964.21.

411 Trifolium fructu cochleato, ciliare BM. 269.

412 Trifolium folliculis Corona similibus CB. Chabr. 167. f. 1.
  412 Trifolium folliculis Corona fimilibus CB. Chabr. 167. f. 1.
413 Valerianella fem. umbilicato rot. nudo Morif.
414 Valerianella fem. umbilicato nudo oblongo Marif.
415 Verbafeum fol. subrotundo, fl. Blattaria CB. (147.4-
416 Vicia multifolia cum latis siliquis I.B. Fl. Apr. &c. Chabr.
417 Vicia similis, supra infraque terram frustum edens CB.
418 Vitis sylv. Labrusca CB. (Fl. Apr.
419 Ulmus fol. minus rugoso & scabro ferè lucido I. B.:

In Appendice Botanic. Monspeliensi.
420 Alga tinctoria I.B. R. 73. c. 6. pl. 2.
421 Allium petræum umbelliserum I.B.
422 Alsine littoralis Graminea BM. 290. Fl. Maio.
423 Aparine minima BM. 291. Fl. astate.
    423 Aparine minima BM. 291. Fl. astate.
424 Bellis mont. Gramineis foliis BM. 291. Fl. Maio.
                           Carduus stellatus, fol. integris serratis BM. 292.
     426 Carlina caulifera & acaulos I.B. R.288. 1.
                         Caucalis Dauci fylv. fol. echinato magno frudu BM.2928 Chamaciflus repens Serpilli folio luteus CB. Fl. Maio. Chamaciflus Nummularia folio BM. 294. Corallina fiftulofa fragilis lævis I.B. Chabr.578.f.t. Daucus mont. fol. Faniculi longiona RM. 204. Fl. 36.1
429 Coamacypus summularia tolio BM. 294.
430 Corallina fiftulofa fragilis lævis I.B. Chabr.578.f.1.
431 Daucus mont. fol. Faniculi longiore BM. 294. Fl. Jul.
432 Idem multo minor BM. 295. an. Lugd.
433 Dens Leonis fol. radiatis BM. 295.
434 Doronicum integro & craffo Hieracii folio BM.295.Fl.Jun.
435 Fuci Fungiformes I.B. Ray 78. 12.
436 Gramen paniculatum minimum molle BM.297.Fl.Apr.&c.
437 Gramen paniculâ fparfâ luteâ Loliaceâ BM.297.Fl. Maio.
438 Hieracium mont. laciniat. lanuginofo folio CB.
439 Lafluca fylv. angusto laciniato folio BM.298.Fl. Jul.&c.
440 Lapathum prat. capite comosiore BM. 300. Fl. Maio &c.
441 Laferpitium Gallicum CB. Fl. Jul. R. 426. 2.
442 Idem multo minus BM. 300.
443 Lenticula palustris bisolia, frustu tetragono CB.
444 Limonium fruticosum BM. 301.
445 Lupinus sylv. purp. sl. fem. rot. vario majore & minore BM.
446 Orchis store nudi Hominia estigiem representans mas CB.
447 Orobus sylv. angustisol. Asphodeli radice CB.
448 Porus Anguinus I.B. Chabr.575.f.z.R.64.9. (304. Fl. Jun.
449 Potamogiton minus, sol. densis mucronatis nonserratis BM.
450 Spongia ramosa varietas Bocc.
451 Tithymalus Lithospermi maioris solio RM. 206.
```

Figure of the state of the stat The state of the s A The state of the de de mile. La como de la como dela como de la como de the second of th or the constitution of the Sold of the state The many services of the servi 1.1.1.1.1.1.1

Plantæ SILESIACÆ Rariores ac Desideratæ,

E Casparo Schwenckfeltio excerptæ, & Methodo Rayana digestæ.

A Jacobo Petiverio, Societatis Regalis Socio. Londini.

Cetosa Alpina pag. 8. Ad Rivulos altiss. montium.

Acetosa minima 8. Agris incultis & collibus squalidis.

Atriplex palust. polycaulis 27. Fossis, Julio, &c.

Bistorta Alpina maxima 31. Jugis altiss. Suditorum.

Lapathum palustre minus 118. Fossis humid. frequens.

Polyonum fol. rubentibus 163. Vis & Agris.

Chondrilla Viminea 50. Fl. Jun. & Jul.

Hieracium 2. 6. 8. 9 & 10. Schwenckfeld. 106.

Hieracium 11.12. 13. 14 & 15. Schw. 107.

Laduca sylv. purpurea 114. Mont. Suditorum.

Pilosella Alpina 157. Montibus altis & herbidis.

Sonchus lavis latis. & angustisolius 199.

Sonchus Alpina cærulea.200. Fette Henne. Siles. Fl. Jul. &c.

Aster luteus montanus 25. Dumet. & collibus saxosis.

Erygeron Alpinum 67. Mont. altiss. & humidis. Jul. &c.

Gnaphalium tenusolium 87. Agris & Hortis Oleraceis.

Tussilago Alpina minor 212. Mont. umbros. & humidis. Jun.

Virga aurea 1 & 2. Schw. 219. Wundkraut Germ.

Bellis mont. maxima 29.

Chamemelum fatuum 47. Poldisha Agris Guict & and the Tuffilago Alpina minor 212. Mont. umbrof. & humidis. Jun.
18 Virga aurea 1 & 2. Schw. 219. Wundkraut Germ.
19 Bellis mont. maxima 29.
20 Chamamelum fatuum 47. Ruddille. Agris quiet. & cultis.
21 Abfinthium mont. f. vulg. minus 6. Ruderibus & montosis.
22 Carduus palustr. fruticans 38. Mont. & Sylvis udis. Aug. & c.
23 Carduus pratensis 39. Mont. pratis humidis. Jul. & c.
24 Carduus pratensis 39. Mont. pratis humidis. Jul. & c.
25 Carlina major Schw. 39. Eberwurtz. Pl. Jul. & Aug.
26 Cirsum montanum 1. 52. Schardistel Sil. Fl. Jul.
27 Jacea collina minor, leptophyllos 111. Agger. Viarum. Jul.
28 Jacea mont. villoso capite 111. Pratis Mont. Riphæorum.
29 Scabiosa minima aspera 192. Mont.sabulos. & asperis.
30 Scabiosa sinima aspera 192. Mont.sabulos. & asperis.
31 Angelica Alpina 2. Schw. 18. Lustrwurtzel Sil.
32 Angelica Alpina 3. 19. Geistwurtzel Sil. Mont.
33 Apnum montanum 20. Mont. & arenos. pratis. Jul. & c.
34 Cicutaria montana 52. Montosis humidis.
35 Daucus mont. Boienicus 60 ad Poidebradum, Julio.
36 Libanotis Theophrast alba & nigra. Schw. 122.
37 Meum Silesiacum Alpinum & palustre 136.
38 Millesolium palustre 1. 138. Wasser Fenchel.
39 Millesolium palustre 2. 128. Wasser Fenchel.
39 Millesolium palustre 2. 128. Wasser Fenchel.
39 Millesolium palustre 2. 128. Wasser Fenchel.
30 Millesolium palustre 2. 128. Wasser Fenchel.
31 Aprum montanum 20. Mont. Stefaca. Rote Germ.
40 Sesum aquat. odoratum & aquat. 24 Schw. 197.
41 Thalistri omnes species Schw. 204.
42 Mollugo 1. 2. & 3. Schw. 139. Megerkraut Germ.
43 Rubia sylv. major 184. s. smont. Silesiaca. Rote Germ.
44 Cerinthe minor 46. Viis. Agris & Collibus. Fl. Estate.
45 Cannabis sylv. spuria 1. 2. & 3. Schw. 37. Taubnesseln. Germ.
46 Cerinthe minor 46. Viis. Agris & Collibus. Fl. Estate.
47 Cannabis sylv. spuria 1. 2. & 3. Schw. 37. Taubnesseln. Germ.
48 Hedera terrest. montana 101. Opacis Mont. Sylvis.
50 Lamium pictum 116. Milzatella i. e. Lienaris.
51 Serpillum Boiemicum, odore unctuoso suvissimo 195.
52 Sideritis 1 196. Zeissenka. 49 Hedera terrett. montana 101. Opacis Mont. Sylvis.
50 Lamium pictum 116. Milzatella i. e. Lienaris.
51 Serpillum Boiemicum, odore unctuoso suavissimo 195.
52 Sideritis 1 196. Zeissgenkraut Sil. Fl. Jun. & Jul.
53 Sideritis 2. 196. Brauner Andorn. Agris.
54 Filipeudula montana 71. Monte Giganteo. Julio.
55 Aconitum luteum majus. Goel Wolffswurtz. 8. Fl. Jun. & Jul.
56 Aconitum cæruleum 9. Pratis agro Hirsberg. ad Bober.
57 Napellus minor 143. Saltubus Mont. ad Fluv. Bober.
58 Napellus major 143. Fuchswurtz Germ. Aug.
59 Caryophyllata Alpina minor 42. Vallibus umbros. Suditor.
60 Malva sylv. s. Alcea Querna 132. Fairis Germ.
61 Plantago aq. minor s. angustif. humilis 161. Udis.
62 Pulsatilla nivea Alpina 771. Celsis Suditor. jugis. Jun.
63 Quinquefolium Vernum majus 174. Incultis &c. Mart.
64 Quinquefolium Vernum minus 174. Collib. & Viis. Aprill.
65 Quinquefolium Fragiferum erectum 175. Floret Maio.
66 Quinquefolium Fragiferum erectum 175. Floret Maio.
67 Ramunculus Alpinus candidus 179. Ad Rivos & sequente.
68 Ranunculus Alp. luteus maximus 179. Mont. alt. udis, Jul.
69 Ranunculus Mont. hirsuus Narcissi-florus. 189. Junio.
71 Sedi omnes species Schw. 194. & 5.

72 Tormentilla Alpina maxima 208, ad lacus Sudit.&M.Riphæo.
73 Polygonatum angustifolium 162. Mont. sylvosis.
74 Polygonatum lacifol. ramosum 162. Humid.& Umbrof. Sudit.
75 Anagallis aq. 2.18. Palustribus & Rivulis.
76 Anagallis aq. 3.5. minor 18. Prine.
77 Bursa Pastoris minor, fol. persoliato 34. Mont. apric. & arid.
78 Bursa Pastoris Myagrifolia 35. Nagelkraut. Agris aridis.
79 Dentaria Pentaphyllos 61. Flo. Aprili, Umbros. Sudit.
80 Dentaria Heptaphyllos 61. Dod. 162. f. 2.
81 Dentaria Enneaphyllos 61. 81 Dentaria Enneaphyllos 61. 82 Dentaria baccifera 61. Opacis Mont. Maio. 83 Eruca palustris major & minor Schw. 65. 83 Eruca palustris major & minor Schw. 65.
84 Lysimachia siliq. palustr. & Roseo store Schw. 129.
85 Nasturium montanum 145. Pratis copiosè. Maio.
86 Thlaspi album supinum 205. Viis circa Vratitav. Æstate.
87 Tithymalus Cyparistias capit, rubente 207. Mont. andis.
88 Tithymalus strêtophyllos & platyphyllos Schw. 207.
89 Tithymalus frutic. palustris 208. Ad Sabulos, crepid. Flumin.
90 Rapunculus Nerisolius 181. Vallib. sub. M. Gigantao. Jul.
91 Rapunc. sylv. cær. umbellatus 181. Mont. ad Quissum. Aug.
92 Trachelium Alpinum maximum 229. Flor. Julio.
93 Gentiana omnes species Schw. 83.
94 Assine petwa omnium minima 16. Petrosis Jun. &c.
95 Asine sontana 16. Rivulis limpidis pigiè labentibus. Assine fontana 16. Rivulis limpidis pigie labentibus. 95 Asine fontana 16. Rivulis limpidis pigiè labentibus.
96 Asine palustris 16. Aquis residibus & palustribus.
97 Caryophyllus sylv. 1. 2. 4. 5 & 6. Schw. 42. & 3.
98 Cistus humilis 2. Panax Chironium minus 53. Apricis.
99 Primula Veris inodora Alpina 168. Mont. altiss. Muo.
100 Sanicula Alp. s. Auric. Ursi minima st. puip. 190. Maio.
101 Pyrola frutescens 172. Mont. & Sylvis Gorlicens.
102 Lagopus montanus minor 115. Ad Agros & Vias.
103 Pisum sylv. perenne 156. Mont. asperis inter Saxa. Julio.
104 Pistum sylv. 2. s. minus alterum 159.
105 Polygala repens Nuperorum 161. 103 Pisum sylv. perenne 156. Mont. atperis inter Saxa, Julio, 104 Pisum fylv. 2. s. minus alterum 159.
105 Polygala repens Nuperorum 161.
106 Trifolium sylvat. alb, maximum 210. Ericetis. Julio, 107 Vicia sylv. 3. seu lutea 218. Campis fruticosis. Jul. &co. 108 Allium Alpin. mas & sem. 14. Suditor, humid. Julio, 109 Bifolium minus 31. Mont. & Vallibus opac. & udis. 110 Bifolium 3. seu Opbris trifolia 31. Dieyblat. Germ. 111 Damafonii onines species Schw. 59.
112 Orchides omnes species Schw. 147. &c. 113 Porrum agreste 166. Agro. Marginibus. 114 Porrum agreste poliferum 166. Agris inculsis. 115 Por. Alpin. Sestinum maj. 166. Vallib. Oupa & Mont. set. 116 Satyrum abort. s. Dentaria aphyllos minor 199. Jul. &c. 117 Valeriana f. Phu montana 215. Mont. humid. & th. roid. 118 Adiantum alb. crifpum Alpinum 10. Mont. altiss. & petrosic. 119 Filix sontana 73. Muris puteorum & sontium. 120 Filix mas Alpina maxima 72. Mont. humid. & saxosis. 121 Filix muraria seu minor 72. Saxispixiuptis. 122 Filix spinosa seuleata 72. Mont. opacis altiss. 123 Lonchitis alpera major 125. Asperis altiss. Sudito. 124 Lunaria Botygitis minor polyphyllos 126. Inter Grov na 125 Musci omnes species Schw. 739. 126 Graminum omnes species Schw. 739. 126 Graminum omnes species 88, &c. 127 Juncus sylvaticus 112. Sylvis udis. 129 Juncus minimus lanceatus seu spicatus 11. 130 Acer brusum, intortius criftum 7. Urlenne Flader. 38 128 Juneus acutus minimus 112. Pratis uliginosis.
129 Juneus minimus lanceatus seu spicatus 11.
130 Acer bruseum, intortius erispum 7. Urlenne Flader. Sis
131 Fraxinus erispa 76. Ejus materies unduletur.
132 Fraxinus dura s. petixa 76. Mont, petros. & præruptis.
133 Fraxinus Frugisera 75. Montibus & clivosis. Aprili.
134 Ledum Silestacum majus 120. Montibus. Jun. &c.
135 Pinaster Alpin. 1epens 159. Jugis Montium 121.
136 Ribes inspidum 182. M. Ripboo. Fl. Min. Fr. Autum.
137 Rosa Alpina rubra 184. Ad Rivulos Montium. Jun.
138 Rosa Arvina s. Avenaria 184. Ad Viausav. inter Avenas.
139 Rubus Hircinus 185. Saltibus nemotosis Montium.
140 Salix latisolia pratensis 187. Udis Mont. pratis.
141 Salix saxat. s. Alpina repens 188. Jugis Sudit.
142 Ulmus montana 220. Montosis ad atcem Furstenssein.



PLANTARUM ETRURIÆ Rariorum GATALOGUS.

E Clarissimo Casalpino, Boccone, Mentzelio, Raio, &c.

Arnia glande maxima rotundiore, pag. 32.

Parnia maritima, operibus inutilis, p. 32.

Suberella folio latiore modice finuato 32. Maritimu.

Carpinus folliculis membranaccis 38. Alpibus.

Opulus Aceri fimilis, foliis minoribus 42. Planis.

Frazinus humilis 43. Siccis & faxofis.

Cerafus sylv. ff. parvoex viridi flavo, racemulis oblongis 51.

Ilatrum folio Ilieu angustiori 74. Maritimis.

Legro Puzzo & Putine 74. Agro Plumbiensi.

I. Rhammus fol. Olea, sl. ex purp. candicante 74. Sepibus.

II. Sambucus umbellis parvis, baccis subentibus 92. Montibus.

12. Sambucus palustris, baccis depressis 92. Circa Pisam. 11. Sambucus umbellis parvis, baccis subentibus 92. Montibus.
12. Sambucus palustris, baccis depressis 92. Circa Pisam.
13. Rubus humilis, baccis paucis 98. Humidis.
14. Alvis frutex in Alpibus sine spinis, Ribes vulgo. 99.
15. Lycium fol. serrato perpetuo ror. Humectis & incultis.
16. Abutus maritima, fr. asperiore slavo 103.
17. Crissus st. pallido, siliq. turrectis spum vulgo 113. Mont.
18. Crissus fol. albidis acaulos 114. Collibus maritimis.

Crissus fol. nigrioribus, pedic longis 114. Collibus siridis. 18. Cytifus fol. albiers acautos 114. Collibus maritimus.
19. Cytifus fol. nigrioribus, pedic.longis 114. Collibus frigidis.
20. Genista Alpina aphyllos, vincis palmaribus mollibus 115.
21. Cenista humilis, juncis acutis robustis 115. in Ilva.
22. Spina Christi vulgo, Frutex spinosus aphyllos 117. (Alpibus.
23. Populus fol. leviter servatis infernè subhissus, punct. albis 21. 22. Spina Confer vulgo, Protest pinotus aphyllos 117. (Alpibus. 23. Populus fol. leviter ferratis infernè fubbirfut, punct. albis 21. 24. Salix fol. angustiore, viminibus rubris 125. Alpium fluviis. 25. Salix 3, viminibus ad circulos Lagenarum colligandos 125, 26. Salix 4, fragilis, fol. subrot. histuro, Vetricem vulgò 125. 27. Valeriana Sorbi folio subbirsuto 148. Montibus & Riguts. 28. Nardus montana, Asphodeli 12dicibus 149. 29. Spica Celtica, umbellà luteà 149. Montibus. 30. Valeriana inodora, Viola folio 148 Umbros. Mont. jugis. 31. Valeriana sylv. similis, palmi longitudine 150. Ilva. 32. Valeriana fol. oblongis, fiosculis puniceis 150. Pisa mæniis. 33. Laureola st. albo, baccis nigris 150. Montoss. 34. Olivella Alpicolis, fol. molli, st. albo 151. Apennimis. 36. Olivella fol. duro piloso, st. purpureo 151. in Silicia 36. Cneotum Cupressi tolio supernè candic, Miccia Hisp. 621. 37. Cassa lignea similis, st. albo 155. Alpibus. (Marit. 38. Badem tragilissima, stosculis herbaceis 155. Saxosis. 39. Eadem Erica foliis crassioribus 155. Marit. asperis. 40. Bupleuron latifol. Cubitali altitudinis 156. 41. Bupleuron latifol. Cubitali altitudinis 156. 42. Ambrosiana Arthemissa foliis 158. 43. Herba Turca humistrata, Polygoni foliis 159. Asenosis. 44. Pelvagarum. Centinodia vulce Lingua Passerina altic 168. 44 Ead. fol. angustioribus, Palmiaititud. 159. Mont. labulofis.
45 Perscaria fol. aspero, acerbioris saporis 167.
46 Polygonum, Centinodia vulgo, Lingua Passerina aliis 168.
47 Polygonum erect. fol. & semine longiori 168. (ubique 8 Polygonum candid. humistratum marit. 162.
49 Alsne fol. bins Portulaça crassis, of ore Cucumer. 169 Latic & c. Salosa Oleracea fol. vermic. aculeatis, fl. rubente 170. Mar.
51 Salosa Oleracea Horror Vicaria vulgo 150. 50 Salfosa Oleracea fol. vermic. aculeatis, fl. rubente 170. Mar. Salfosa Oleracea Hortor. Isgarum vulgo 170.
52 Salfosa fol. multis tenuioribus rubris 170. Marit.
53 Salfosa fol. gracili insipido 170. Muris & Sterquilinis.
54 Salfosa fruticans lignosum perpetuum 171. Scopulis Ilvæ.
55 Far iem. rusto, spica crassiore quam Speita 173.
56 Far sem. albo crasso, spica mutica 173.
57 Spelta grano lævi, arista subaspera 174.
58 Hordeum spica brevi, Mont. Lucens. Scandella dista. 176.
59 Hordeum mundum in Ptisanis utitur 176.
61 Avena Graca sem. rubro. aristis longis 178. Arvis seguens 60 Hordeum mundum in Ptisanis utitur 176.
61 Avena Graca sem. rubro, aristis longis 178. Arvis sequens.
62 Gamen Cauda Vulpina dicta, spica mollissima 183.
63 Panicastrella spica aspera 185. Vineis & hortis
64 Panicastrella spica è folliculis latis candidis 185.
65 Luciola herba, quia noste lucet 185. Montibus.
66 Phenix pusilla, spicis Lolii puniceis 185. Muris & tectis.
67 Perlara sol. Tritici, juba Milii, rad. rot. candidis 196. Mont.
68 Herba Capillaris, spicis parvis ramosis 186. Alpium pratis.
69 Herba Capillaris, sculmo cubitali, spicis ex solviculis 186.
70 Gramini similis, stose, purp. spicatis, major & minor 186.
71 Avundo crassa ad Vites sustinendos colitur 186. (Muris.
72 Arundo calamis songist. humi serpens 186. Maritimis.
73 Juneus rotund. lanugine vice Seminis 188.

74 Juncus Levis, floccis orbiculatis migris t38.
75 Juncus magnus ad Tegete: contex indas 188. Agris.
76 Phalaris exilis, 3 aut 4 paureulis 193.
77 Juncus folia 3 in cacumine inflexa 195. Minit.
78 Arum parvum, baccis nigris 227. Monte Sancti Juliani.
79 Rubilium fol. binis angustis, st. purp. 233. Seget.
80 Rubilium fl. multis Roleis, filiq. tenuibus nigris 233. Sylvis
81 Cicercula fl. purp. fem. albo nigro & vario 234. Seritur.
82 Mochi vulgo. Alpibus seritur, ad pabulum Brutorum 194.
83 Aracus fl. tuteo, fem. nigris scabris 235. Seget.
84 Cicer sylv. fol. ternis, Theriaca vulgo, ab odore 238. Coll.k.
85 Melilotus sylv. fol. subhirsutis, filiq. incurvis 244. Ædisc. vet.
86 Glaux fol. hirsutis, fl. ex albo purp 245. Asperis marit. 85 Melilotus sylv. fol. subhirsutis, filiq. incurvis 244. Adific. vet.
86 Glaux fol. hirsutis, f. ex albo purp 245. Asperis marit.
87 Polygala Meliloto similis, f. magnosingulari pallid. 245. Mar.
88 Tragos fol. pusillis, flose. in summo albicant. 245. Marit.
89 Tragos fol. Glauci latiore, bicubit. altitudinis 245.
90 Lupinus sylv. dicla, 247. Marit.
91 Lupinus caulic. dodrant. erectis, fl. rubente 247. Asperis.
92 Astragalus st. salbo aut rubenteorbic. filiq, Meliloti 248.
93 Marit.
94 Anthyllis 1 quibus dam, Lentis fol. flose. luteis aduncis 249.
95 Onobrychis st. rubente, filiq. Viciae migris 249. Mont. riguis.
96 Onobrychis sylv. st. purp. fol. latioribus & paucioribus 249.
97 Sena fol. latioribus, filiq. compressis salcatis 249. Agris serit.
98 Herba humistrata fol. hirsutis, fl. luteis, siliq. tumidis 250.
Saxosis. Saxosis.

99 Lychnis fol. molli, fl. ex puniceo purpurascente 253.

100 Lychnis exilis, fl. Centaurii minoris 253. Montibus.

101 Saxifraga aphyllos, ramis junceis, fl. parvo purp. 253. saxosis.

102 Saxifraga Romana, rad. & fl. albidis 255. Pariet, umbtosis.

103 Saxifraga Romana, rad. & fl. albidis 255. Maritimis.

104 Ocymoides marva, humistiata, fl. rubente 256. Ruderibus.

105 Ocymoides parva, humistiata, fl. rubente 256. Saxos. asperis.

106 Nifeago spica racemosa 257. Atidis & maritimis.

107 Pelemonio levi similis, Rapum sylvestre dicta 259. Arvis. & c.

108 Odontides caule dodrantali tecto, fl. purp. 259. Pratis

Udis. (passim) 109 Centunculus, aliis Centone & Mossus Gallina 259. Ruder.
110 Gratia Dei altiof & ramosior, soi. longioribus 265.
111 Fior Capuccio sol. & st. tenuioribus 267. Segetibus. 112 Onagra 3, folio Hedera serratoribus 268. Montibus.
113 Fumaria caule longiore, fol. rarior. fl. candicante 272.
114 Ferula mas fol. tenuius scissis 276.
115 Ferula semina 276. Monte Argentario.
116 Thapfia fol. latioribus subcandidis 277. Ilva spontė. 117 Panax Asclepium, foliis Femiculi 279. Montibus.
118 Meu salso, fol. capillaceis, st. luteo 280. Montibus.
119 Libanotis nigra, st. luteo 282. Montibus & umbross.
120 Meu fol. Anethi, st. luteo, Imperatoria quidam 283. Alpib.
121 Anni sem. minutissimo suavi 285. Ilva. Anmi pusillam 286. Asperis, Apricis inter saxa.

Daucus Mont. solio & umbella Staphylini 286. Alpibus.

Daucus campestris fol. Fæniculi 287. Latis & Agrer. lim.

Caucalis soliis varie divisis 287. Arvis ubique. 125 Caucalis tollis varie divilis 287. Arvis ubique.
126 Gingidium tol. Staphylini, fl. albo 289. Ruder. & Vineis.
127 Gingidium fem. grandi hispido 290. Collibits afferis.
128 Cicuta brevior & odoratior 292. Alpibus.
129 Cicutaria edulis cocta 273. Collibits, humestis.
130 Cicutaria hisfutior, femine grandiore 293. Alpibus.
131 Finocchiella Alpicolis, odore & sapore Famiculi 293. Panedfeolus flore purpureo 293. Montibus. Pyrethrum Clanarum Incolæ Imperatoria 294. Palustribus 133 Pyrethrum Clanarum Incolæ Imperatoria 294. Palustribus 134 Sefeli Massiliente, Fanicuum marin., quibus dam 294. Pisa 135 Sefeli Creticum 295. Sylvis & Collibus maritimis. (marit. 136 Sefeli Elbiopicium, sem. nigro odorato 296. Arenosis. 137 Sefeli peculiare fol. imegro 296. Segetibus. 138 Sefeli parvum fotio Graminis 296. 139 Crescione & Herba Canella vulgo 300. Aq. stagn. Es limosic. 140 Crescione caule resto, umbellis tummis ramulis 500. 141 Sison sem. parvis nigris odoratis 301. Aquis Es udis. 142 Levissicum alterum st. subrubentibus 305. Incuttis, squis-143 Ligusticum, Alpibus Liguria imperatoria 506. 143 Ligusticum majus, umbelia lutea 307. Montibus. 146 Augelica vulgo Aspicolis, sem. molchato 307. 147 Herba Costa vulgo, store & succo luteis 310. Montibus.
148 Panaw Herenleum vulgo Panacea Etruscis 311. Alpibus.
149 Siler Germanicum, solvis Pafrinacea, &c. 314. 150 Saxifragia 4. fem. ciassiore 315. 151 Rubia lylvestris 323. l'epibus, Mont. & aridu. 152 Rubia lylv. caule bieviori, fem. nigro 323. 153 Lappago quidam hevis 325. Ubique. 154 Cincialis longe exilior 325. Montibus. 155 Crucialis in maritimis minima 325. (Montibus. 156 Herla quadam dodiantalis, fol. binis angustis, &c. 326.
157 Fruticulus in Collib. (axosis, &c.fr. rotundo compresso 326. 158 Herba stella edulis cruda, foliis Gramineis 330. Montibus faxofis. 159 Herbassella altera lanuginosa, Sespentina quidam 330. Ædi-160 Antirchinon alterum exclius, Parietibus fréguens 335. 161 Auricula Muris altera, in Parietibus 335. 162 Phyllon alterum humilius, folio exiliore 337. 163 Pyrola fl. albo 338. Alpibus. 164 Pyrola fol. Bata carnolo, caule foliato 338. Aquosis,
165 Circea similis, fol. submittutis crispis 339. Montibus.
166 Euphrasia altera ramosa, fol. maju culis 339. Liguria Alp.
167 Euphrasia fol. purp. ex latitudine cristata 340. Montibus cultis, &c. 168 Crista altera, fol. angustiss. st. sureo 34r. Ericetis. 169 Crista simplici caule dodrantali 341. c. 37. 170 Crista ramis rectis, cubitali altit. 341. Praris.
171 Crista quædam in Alpibus, folio Filicis 342. c. 38. 172 Orobanche vulgo Cauda Leonis, fl. albicante 342. 173 Orobanche ramofa, fl. purpurascentibus 342. 174 Verhafeum 7. tol. latis lubrot. fi. luteo 347. Montibus. 175 Verbascum 9. fol. angusto sinuato virenti, st. aureo 347.
Ruder. passim.
176 Galesbelos & Urtica mortua, Verbasci structu 349. Ruder
177 Scrophularia tol. purpurascente 349. Alpibus riguis.
178 Scrophularia similis minor, fol. Rutæ scribis 350.
179 Lingvis passa, fol bravibus tibbissuis suran aco Vincia (passim. 179 Linaria parva, fol. brevibus lubhirlutis, fl. purp. 350. Vineis. 180 Os Leonis lol. brevibus lubhirlutis, fl. pallido 351. Scopulis pranuptis.
181 Os Leouis fl. brevi caruleo, capit. rotundioribus 351.
182 Brassica sylv. ingrata, fol. è cand. purpureo 352. Ilva scopulis, Ecc. 183 Sinapi 3 Ruca vulgò, fol. Eruca scabro 359. Hortis.
184 Eruca altera fiuticosa, fl. luteo 360.
185 Eruca sol. hirluto, fl. albo 360. Parietariis & maceriis. 186 Herba Sancti Alberti, fol. laciniato, fl. luteo 361. 187 Accipitina alter fol. crassus divisis, fl. candido 361. Incultis. 188 Herba quadam, caule simplici, fol. seriato, fl. albo 361. 189 Sifyn brium caule & fol. subrubentibus 362. Aq. riguis.
189 Sifyn brium caule & fol. subrubentibus 362. Aq. riguis.
190 Sifynbrium fruticosior, fol. Erucæ laciniato 362. Priore.
191 Hobula humilis Burfa Pastor. assinis, Thymi fol. rot. 366.
192 Iblaspi fol. Iberidis, st. albo 366. c. 76.
193 Thlaspi st. luteo fr. lato valde depresso 366. Monte Sancti Juliani. (ore. Thlaspi Cieric. fol. Lini, fl. subrubente 367. Afperis, pri-195 Thlafpi magnum, Rafanus montanus vulgo. 367. Horris & Montibus. Montibus.

196 Dorella vulgo, fol. Thlaspi nigii, sl. candicante. 367.

197 Hesperis sl. albo sl. purp. nocte olens 368. Montibus.

198 Hesperis pusilla, sl. purpureo 369. Gracilibus.

199 Hesperis tol. & capite rotundis, sl. luteo 369. Montibus.

200 Lunaria Graca z. inodora, fol. Urticæ 369.

201 Lunaria Graca 3. fol. Verbasci, sl. luteo, sliiq. sursis 369.

202 Lunaria Graca 4. stuticosior foi. Thymi. sl. luteo 360. Montibus. Lun. Graca 4. fruticosior, foi. Thymi, fi. luteo 369. Mon-203 Ruta sylv. Aimell Offic. sl. albo 371. (tacuto. 204 Ruta 2. humilior, Greuta Athemenfium quidam 371. 204 Ruta 2. humilior, Cicuta Zinemenjam Galaca, 572. Sept. 205 Thalicilion, vulgo l'erdemarcum 372. Agger. agrorum. Ec. 206 Characias fol. Olea subhissuto, fl. subrubente 373. Marit. asteris. (Montibus. Char. fol. Olea, ramis rubent. paucioribus & brevior. 373. 207 Char. 101. Olea, ramis rubent. paucioribus & brevior. 373.
208 Tithymalus palustris 274. Pifa paludibus fiequens.
209 Efula Offic. fol. Lini, comă subrubente 374. Campestribus.
210 Efula niaj. fol. longior. & sarioribus 374. Humidis.
211 Efula 3, s. candida, fol. angustis 374. Montofis.
212 Alypum sol. Thymi, fr. aspero 375. Petrofis marit.
213 Tithymalus Myrsmites angustioribus soliis 376. Montibus.
214 Tithymalus Arborescens 376. Scopulis & prasuptis.
215 Tithymal. latis. alter, sol. & si. hissuis 377. Ad Fluvias vastin pashm pajim

216 Cataputia admodum pufilla, 4 digitorum 378. Afperis.

217 Scamonia major, fol. Hedera angulofis 383. Hortis.

218 Convolvul. humitpartus, ft. purp. 385. Collibus afper. marit.

219 Phyteuma alter, Archangetica foliis minoribus 386. Erinos fol. angulotis, fiore albido f. purp. Rapunculi 386.

Haific. riguis.

221 Sesamoides alter, caule unico 388. Arenosis. (aridis. 222 Sesamoides 3. fol. brevib. paucis appendicibus 388. Glareos. 223 Cyclamen vulgo Pamporcinus 392. Montibus umbrosis. Cyclamen Veinung, rad. Avellanis magnit. 392. Montibus Sancti Juliani. Viola lutea, gracilioribus foliis 393. Montibus. 225 Viola lutea, gracino nous foits 393. Montibus.
216 Coris fol. angustiss. fl. & fr. Hyperici 396. Montibus.
227 Coridi similis, fl. ab uno latere luteis 396. Marie. asperis.
228 Chironia caule simpl. erecto, fol. maj. hissutis 397. Sabu-10/15.
229 Chironia flore candido majori 397. Collibus Urbevetan.
230 Cepa dulcissima, Nucis magnitudinis 400. Agro Pistoriensi.
231 Cepa lanata, habens lanam sub primis tunicis 400. Littoribus.

232 Bulbus fylv. quidam Hydeinthus fylv. fl. cetul. 40. Campis.

233 Cepa Canina, fl. candicante 402. Pratis paluffribus.

234 Hydeinthus alter exiguus, fl. diffecto 403. Sylvis.

235 Seilla radice nigra 403. Montofis marit.

236 Moly Homericum foboliterum, fl. candido 404. Montibus.

237 Ornithogalum luteum alterum breve fl. dodrans 405. Mont.

238 Onnithogalum luteum alterum breve fl. dodrans 405. Mont.

239 Ampelopraffum fobolit. fl. albo pendente 407. Collibus.

240 Alhum fylvestre, fl. rubente 409. Arvis 85 Vineis.

241 Allium Ursimum alterum, angustiore 601. 409. Sylvis.

242 Colchicum candicans, fol. angustiore 410. Campis &c.

243 Colchicum candicans, fol. latiori 410. Montibus.

244 Crocus alter, fl. albo 6. purp. 411. Agro Pisano, legetibus.

245 Lonchitis fl. luteo, Tulipa species 412. Aprin. Bargens.

246 Narcissus aq. fl. candido, sine medio acetabulo 414. Hu
247 Asphodelus cause nudo simplici 415. Aspibus. (medii. 246 Narcissus aq. st. candido, inne medio acetabulo 414. Hi 247 Asphodelus caule nudo simplici 415. Asphur. (medi 248 Asphodelus caule nudo ramoso 413. Maririmis. 249 Asphodelus parvus, caule nudo ramoso 416. Asperis. 250 Asphodelus parvus, caule nudo ramoso 416. Asperis. 251 Iris pusilla purp. rad. odorata 421. sequente 252 Iris pusilla pallida, rad. inodora 421. Saxosis sterilibus. 252 Iris pusilla pallida, rad. inodora 421. Saxosis sterilibus.
253 Satyrium lingua lata deorsum stexa 428.
254 Satyrium bisurcata purp. 429. Urbevetano. (Mons.
255 Sigillum Sansta Maria, st. 3 st. 4 in cacumine candidis 431.
256 Sigillum S. M. alter. store cand. purp. 431.
257 Sigillum S. M. 3. st. & rad. candidis 431. Collib. maris.
258 Buglossa quasi Lingua Bovina 432. Sabulosis.
259 Anchusa altera, sol. angustiore 435. Sabulosis.
260 Echion st. è candido subente 436. Semisis & Aridis.
261 Lithospermum caul. simplic. junceis 437.
262 Cynoglossa latiore fol. speciolo store 440. Montibus.
263 Salvia niger 441. 263 Salvia niger 441.
264 Salvia 2 leu Herba alba 441. Saxor. crepid. apud Ser avallem.
265 Melinum fl. ex candido pallescente 443. Apenhinis.
266 Melinum alter, brevior & fruticosior, 443. Gracilibus.
267 Betonica odore magis gravi 444. Montibus.
268 Horminum sylv. Melissa folio 448. Asperis & saxos.
269 Horminum sativo similis, folliculis Ocymi 449. Valle 269 Horminum sativo similis, folliculis Ocymi 449. Valle prope Pisas.

270 Milzadella macula per medium folii 449. Ruderibus.

271 Leucas angustioribus foliis quasi Mercurialis 450.

272 Leucas exigua 450. Ruderibus & locis asperis.

273 Sideritis Heraclea 451. Incultis & sabulosis.

274 Sideritis altior ac viridior, fl. subluteis 452.

275 Sideritis fol. angustiore, fl. purpureo 452. Vineis.

276 Sideritis fol. angustiore, fl. albicante 452. Vineis.

277 Prunella hirsutior & brevior, fl. candido 453. Asperis & saxosis. faxofis.
278 Prunella alia exilioribus ramulis, fol. angustiore 453. Pe279 Teucrium alterum ramosius, fol. densius serratis 455. Mon-280 Scordium ampliore folio Mentastri 456. Planis.
281 Polium Mont. stor. candidis. 457. Collibus incultis saxosis.
282 Polium ramis paucioribus, fol. subrot. non denticulatis 458.

Montibus.
283 Polium exilius, fol. Rorismarini infernè candidis 458. Ass.
284 Rosmarinum sylv. Olea folio, fl. grandiore 458. Ilva saxosis.
285 Thymum Italicum, fol. Satureia 461. Monte Sancti juliani 820. tibus. 11ani, &c.

286 Origanum capit. magis sparsis tenuioribus & longior. 463.

287 Scansapepe vulgo & Moncepepe 464. Olitoribus notissima.

288 Pepolina vulgo, aliis Piperella 466. Saxosis littoribus, &c.

289 Mentha aq. odorata, fol. rotundo, cibis grata 453. Aquosis.

290 Mentassrum fol. oblongo lanuginoso 473. Incultis passim.

291 Absintbium alterum brevius, paucioribus ramis 476. Agro

Senensi. 292 Achillea fol. Abrotani vulg. fl. albicante 481. Mont. cultis.
293 Millefolium pufillum, fol. digitalibus Anmi fimilibus 482.
294 Chrysocome fol. Heliochrysi rarioribus, cap. squamoso 485.
Pisa moniis. Chryfocome tenui folo & faxolo nafcens 485. 295 Chryfocome pluribus capitulis in unum aggregatis 485. Mont. 296 Chry3

forsis 485. Planis.

(Montibus.

(Montibus Ruderibus.

109 Incensaria vulgo radice odorata 406. Montos. aridis Es incult.

109 Incensaria altera tenuiore fol. rad. alba inodora 496. Priore.

110 Incensaria altera tenuiore fol. rad. alba inodora 496. Priore.

111 Daonicium fol. ab radice Tussilaginis pedic. longis 497. Al
112 Cinetaria vulgo 500. Maritimis asperis frequens. (pibus.

113 Jacobea folio Eruca laciniato 500.

114 Jacobea Rapbani follo, sed minus hirsuta 500.

115 Anthylii quædam, fol. Lini asperis, st. Tripolii 500. Mon
115 Sylme Voga aurea fol. Laurinis serratis, fl. Conyzæ major. 503.

11 Firga aurea caule fingulari, fl. in umbellis 503. Alpestribus.
118 Clehorium Romanium, longiore folio 506. 18 Clehorium Romanum, nongiore folio 500.

19 Aphaea angustiore solio 508.

10 Chondrilla 2 fol. hissuis, rad. rotundis substavis 509.

11 Chondrilla quædam, Collibus, Vineis & Edificiis verustis 510.

12 Eddem ramis tuberculis nigris 510. Priore.

13 Anarago caule unico, cui in medio 3 sunt fol rot. 510.

(Sylvis.

(Sylvis. Montibus. (Sylvis.

Anarago altera, caulibus multis nudis 510. Montibus &
115 Endivid fylv. fl. luteo ex alis foliorum 511. Hortis incul117 Sonebus crispiore folio 512. Collibus umbrosis & fylvis.
117 Sonebus caule bicubitali cavo, fol. majoribus 512. Mont.
118 Hieracium magnum, fol. Sonebi subhirsuto 512. Ædiscuis.
118 Hieracium magnum, fol. Sonebi subhirsuto 512. Ædiscuis. 119 Hieracium quædam in ruderibus, fol. nigricante 512. c. 13.
130 Terraciepolus vulgo, Olus sylvestre 513. Collibus faxosis & Graes.

131 Terracrepolus quædam, rad. Fabaceis 413. c. 15. Asperis la Ginestrella sil. Terracrepola minore 513. Vinear. superciliis.

132 Ginestrella minima, foliis Gramineis 514. Montibus saxos.

133 Pilosella altera caule solioso 514. Collibus gracili solo.

133 Laduca sylv. Scariola & Serralia dicta 516. Agror.limit. & Edificiis. 336 Lactura mont. fol. brevibus non scissis & c. 516. Alpibus.
337 Lacturiola vulgo, folio maculoso subhirsuto 517. Sylvis.
338 Lacturiola alia, Sopragine vulgo, grandier & ramosor 517 Arvis.

(renosis.

(renosis.

) Crithmum sylv. sol. & umbellis spinosis 522. Maritimis a
40 Carduus sativus, Carciosus vulgo 424. Hortis Cibi gratia.

41 Carlina vulgo 527. Alpibus freq. in Pratis Apenninis.

42 Carlina altera, Aipibus Pistor. ac Lucensibus 529.

43 Spina Arabica 530. c. 41: Montibus & lovis gracilibus.

44 Acambion 530. Sylvestribus & torrentium alveis. (Agris.

45 Acambion maj. & nigrior omnibus Eriocephalo similis 531.

46 Cirfum Agro Pisano uliginosis ad rad. Mont. S. Juliani 551.

47 Cirfum alterum sol. latior. & brevioribus 531. Sylvis,

48 Circium 3 Stoppione vulgo aculeis durior. 531. Arvis passim.

49 Acorna st. aureo patulo 532. Collibus asperis & Campis. 149 Acorna fl. aureo patulo 532. Collibus afperis & Campis.
350 Senecio altior & major, fl. aureo 537. Montibus.
351 Papaver Spumeum, tol. Papaver. Rheados 539. Incultis & Marit.
(Saxolis. Papaver Spumeum, caule erecto, fol. tenuissime scissis 540. 313 Centaurea major Serratula folio 440. Monte Gargano. 354 Scabiofa folio integro 541. Arvis.
355 Ranunculus lævi ac molli folio 544. Citca Piscinas.
356 Ranunculus lanuginosus Malva folio 544. Incultis, pratis.
357 Ranunculus hirsuto semine fol. subrotundis 544. Arvis & Cultis. 18 Ranunculus hirl. sem, fol. Apii tenuioris 544. Priore.
19 Ranunculus aq. fol. parum scissis, fl. albo 544. Aquosis,
10 Ranunculus humiserpens, Apii palust. fol. non acris 545.
19 foveis. Ranunculus insectus, fol. Flantag. angustiore 546. Humidis. Ranunculus insectus folio Tritici 546. Humidis. Ranunculus mont. fol. rotundo lanuginolo 546. Montibus. Kanunculus mont. 3 folio minus leisto, fl. candido 546. Montibus. Chamamelum Eranthemon fl. rubicundo 546. Inter Segetes. Trinitas aliis Tora fol. subrot. fl. cærul. seu rubro 457. Se-

quente.
7 Eddem fl. candido maculoso 547. Montibus & Opacis.
8 dnemone trifoliata 548. Monte S. Juliani & Collibus

NA STON

369 Pulfatilla fol. Cicuta, fl. purpureo 550. Montofis & Sylvie. 370 Garrofilata fol. rotundo, fl. grandi, purp. f. luteo 550. Montofis. 371 Oenanthe similis Filipendula, fol. Agrinonia 551. Montibus. 372 Fragaria bis in anno fructicans 554. Alpibus Batgeis. 373 Heptaphyllon caule recto cubitali 555. Afferis.
374 Heptaphyllon alter. humile, palmi altit. fl. caucidis 556. 375 Tormentilla altera fol. subtus candidis 556. 376 Pes Columbinus vulgo & Gruinum rostrum 558. Ruderibus.
377 Pes Columbinus st. amplo Leucon subrubente 559. Sylvis. 378 Pes Columbinus fol. rot. divilain multas partes 5 59. Sylvis. 379 Balfamimim vulgo fl. ex purp. caruleis, roffris pratongis 559. Sylvis. 380 Rubertiana brevior acminus rubens, fol. vicidibus 55. 381 Linum sylv. brevius, fl. luteo, capite exiguo 565, 382 Harmel alia, fol. Nigella, cap. tripartito 573. 383 Rhododendron Herbaceum & pufillum, fl. alvo 576. Alpibus. 384 Cepaa altera fol. brevibus denfis 578. Montibus. 385 Cepea 40r digit. altit. fol. modice angulosis 579. Pariet. opacis. 386 Cepaa ramosior, sl. candicante, fr. stellato 579. Priore.
387 Umbilicus Veneris grandior, sol. latiore hyemalis 579.
388 Napellus alter sol. Platani, sl. pallidis 585. Alpibus. 389 Dentaria maj. fl. albo, siliq. simplicibus 585. Montibus & Sylvis. 390 Dentaria min. fobolifera, fl. purp. siliq. exilibus 586. De-391 Dentaria altera brevior non sobolitera 585. 392 Vulparia fol. Napelli, fl. Ranunculi 587. Alpibus. 393 Vulparia fol. rot. ferrato, fl. minore 587. Humedis Alpium. 394 Vulparia quædam fol. tenuissimis, fl. pallidis 587. Alp. Li-395 Filix in palustribus Pisanis arenosis 593. (guriæ. 396 Filix parva, foliolis nequaquam serratis 593. Montibus umbrosis. Filix parva, fol. tenuissime denticulatis 593. Mont. umbr. 398 Hemionitis 595. Vetustis Ædificiis, umbrosis & humectis. 399 Ceterach. fol. dodiantali ramoso hirsutiori 596. Ilva saxos. 400 Adiantim nigrum paucioribus ramis 597. Montibus. 401 Adiantum alterum fol. Digiti longit. Pollicis latit. 598. Liguria Montibus. (598. Sepibus. 402 Equisetum scandens. longiss. caule, fol. nullis aut paucis 403 Equiseto similis, capillaribus foliis numerosissimis 600.

Fluviis fere stagnantibus. 404 Fegatella f. Lichen parva & inutilis 601. Umbrosis. 405 Museus fol. Cupressi sed compressis 603. Mont. umbrosis. 406 Muscus longissimus ex Robore antiqua aut Abiete pendens 407 Musci marini varia species 603. cap. 24. 408 Rola marina sol. rotundis, crebris foraminibus 608. 409 Bursa marina rotunda viridis 608. c. 29.
410 Spuma Maris, globus lævis ex ampullis constans 608.
411 Spuma Maris, fol. auriculatis mollibus 608. 412 Corallina fruticosa, cortice rubro, ramul. tortuosis 611. 413 Corallina frutic. cort. albo, ramulis erectis punctatis 611. 414 Eadem ramulis veluti denticulatis 611. 413 Eadem 4. lanugine nigric. aut cinerea, capillis in extremis granulotis 611. Ha omnes Teftis f. Scopulis marit. adhærent.
416 Fungus marinus minimus 619. c. 48. Lapillis mare nascitur.
417 Thlaspi saxtile, foliis rubris. Mentzel. pugillo. Saxis.
418 Viola Cruciata marina id. Littoribus.
419 Viola lutea marina id. Littoribus.
420 Thymbra spinosa id. Monte Juliano.
421 Acetosa mont. Mali Limon. sol. slagel. ramosis Boc. Mus. 164.

Ala S. Renediff. adhærent. (Montibus Etruria. Alp. S. Benedict. 422 Acetofa mont. fol. undulato, flag. non ramosis id. 164. 423 Sonchus pyramidalis Boc. Mul. p. 41. T. 31.
424 Tragopogon parvum, ex luteo purpuralcens CB. Etrur. (M f. 18. Etrur. 425 Aster mont. saxat. lut. angusto &c. Plantaginis fol. Boc. 426 Virga aurea mont. minor Bocc. Mus. 169. f. 118. Montibus. 427 Cyanus mont. caule foliofo, cap. oblongo Boc. M. 20. f. 2.
428 Malva Betonica folio Boc. 15. fig. Pratis. (Montibus.
429 Rubeola Alp. faxat. tenuifolia Boc. Mul. 145. T. 101. Mont.
430 Rubia lævis Lini folio, fl. albis M. Virginis B.M. 83. T. 75.

Montibus (tylvofis. 431 Pulmonaria Alp. angusto solio Italica B.M. T.86. Mont.
432 Pulmonaria mitis Fragaria odore B. M. T. 105. Montibus.
433 Digitalis angusto Verbasci solio montana B.M. 108. T. 85. Montibus. Nasturtium mont. Afari toliis Boc. 5. Aquosis M. Baige. 435 Thlaspilinium erect. fl. albo, turbinato utriculo B. M. 78. T. 69. Montibus. Planta Monte Pania Etruria &c. E Boccone, & Rayo 436 Chimaleon Alp. Sonchifpin fol. lucido, rad. nigra R. 3. p. 437 Helleborus minimus Alp. Aftrantiæ flore. R. 475.3. (200.22. 438 Nepeta tuberosa spicata purp. Ray Vol. 3. p. 295.7. 439 Nepeta saxatilis carul, minima id. 9.

```
441 Braffica lylv. albido f. nutante filiqua R. 410.
  442 Turritis tol. caulem ambientibus, polyceration R. id. 7.
443 Lysimachia siliquosa Asp. erecta, Alsines foliis R. 426. 18.
  444 Tubymalus dulcis, tuberofa radice R. 431. 54.
 444 Thoymans duties, thoesia radice IV. 4(1. 54.
445 Plantæ Montibus Appenninis à Mentzelio, Raio,
446 Adonis fl. luteo Mentzel. Pugillo.
447 Aljine mufcofa id. Saxofis Montibus.
448 Anagallis fol. acutis oblongis, fl. luteo id.
 440 Anagadis foi deutis oblongis, fi. luteo id.
449 Anemone trifolia, fol. integris id.
450 Aquilegia fol. Thalictri, fi. minutiffimis albis id. Petrofis.
451 Bifforta minima, fpica prolifera id. Pratis Mont.
452 Cerinthe fol. & flor. tennioribus angustis purp. luteis id.
453 Draba [axatilis minor, id.
454 Helical Conference of the co
  454 Helianthemum faxat. caule & fol. incanis oblongis, fl. al-
                                                                                                                                                                                                               (biff. id.
  455 Orchis fol. & flor. ex toto luteis id.
456 Tragorchis purp. barbulis longioribus id.
  457 Pentaphyllum Fragiferum repens, fl. luteo id. Jugis.
458 Pentaphyllum fl. albo 3 unciar, altit. id. Petris.
459 Pimpinella Agrimonoides id. à Columna diffeit.
   460 Polygala lutea maj. pedis altitudinis id.
 460 Polygala lutea maj. pedis altitudinis 1a.
461 Polygala lutea minor, 3 uncias longa id.
462 Polygala frutesc. fol. pilosis & glabris id.
463 Polygala frut. fol. Myrthi, fl. variis id.
464 Polygala frut. fol. tenuibus, fl. alb. purp. id.
465 Polypodium saxt. maj. fol. parum serratis & asperis id.
466 Ramunculus nem. fl. pleno albo & purp. major. id.
467 Ranunculus nem. fl. cæruleo fol. major. & minoribus id.
468 Ranunculus dulcis fol. latis. Rapistri perfoliatis, fl. albo
   468 Ranunculus dulcis fol.latis, Rapistri perfoliatis, fl. albo id.
 468 Ranunculus dulcis fol.latis, Rapistri perfoliatis, st. alboid.
469 Saxifiagia alba faxatilis, pusilia id.
470 Thlassi repens, fol. Serpylli incanis, st. luteo id.
471 Thlassi repens, fol. Serpylli incanis, st. albo id.
472 Trifolium Lupulinum faxatile unciale id.
473 Trifolium Lupulinum, faxatile pusillum id.
474 Trifolium Vesicarium, faxatile pusillum id.
475 Barba Hirci minor, fol. albo Cam. Hort. 27. Jugis Apen.
476 Plantæ Euganeis Montibus, è Mentzelio &c.
477 Alsne media, fol. Solani acutis, st. umbellatis Mentz.
478 Raminculus nem. Stignutoides quibustam id. (423, 30.
  477 Aline media, fol. Solam actitis, ft. unidellatis Mentz.
478 Ramineulus nem. Stigmatoides quibutdam id. (423, 30,
479 Feronica frut. durior, oblongo Chamediyos folio R. 3, p.
480 Nigella flore magno, Montis Ruah Mentz. pugillo.
481 Caryophyllus arvenfis glaber minimus C.B. pr. 105, xi. fig.
482 Belliz mont. major, folio acuto C.B. pr. 121, 3, fig.
483 Centaurium minus spicatum album C.B. pr. 130, fig.
484 Trifalium Cochlectum fol. minuto obullo, ft. lariore C.B.
    484
                    Trifolium Cochleatum fol. minuto obtufo, fr. latiore C.B.
                             pr. 140.9. (287. ec. Planta circa Florentiam, vide alia Ray Sylloge
    486 Fragaria sterilis major & minor Mentz.
    487
                    Fragaria fl. pleno id.
  489 Convolvulus munimus, anguito auriculato folio R.3. 373. 490 Lychnis fylv. minima, fi. exiguo C.B. pr. 103. Agror. marg. 491 Planta Agris & Littoribus Liburni. 492 Alga spiralis marituma Enec 60 Win
   492 Alga spiralis maritima Bocc. 69. Fig.
                    Museus subersimus marinus Mentz. pug.
                                                                                                                                                                                                         (xi. Seget.
   493
   494 Vicia Lathyroides minor, fol. longis &c. fl. binis R. 3. 449. 495 Herninia liirfuta C.B. Arenofis. 496 Cichoreum prat. Vesicarium Col. Sepibus.
   497 Chryfanthemun tenuitolium Bæticum Boelii.
   498 Sideritis fol. angustissimo, caule rubio R. Sylloge 237. Marit.
  499 Medica doliata spinosa polycarpos Park.
500 Lathyrus sl. ex albo & coccineo Ray. Satis, Flor. Aprili.
501 Trifolium album tectum hirtutuum valde I.B.
  502 Trifolium Fragiscium, fl. nitidis iubellu Ray.
503 Bulbus fol. Grammeo hirsuto &c. Ray Syll. 269. descr.
   504 Hyacinthus paluftr. veinus fi. alb. nimbriatis id. 268. Fossis.
  505 Iris humilis latif. fi. violaceo 1.B. Rupibus. 506 Iris pumila latif. fi. luteo R. Syll. 268.
506 Iris pumila latif. ft. luteo R. Syll. 268.
507 Orchis macrophylla Col.
508 Limonium parvum Narb. ft. majore R. H. Pl. 395. 2.
509 Vide plura in Ray Sylloge p. 268. & 285. 6. & c.
510 Plantæ Mintanæ prope Lucam Ray Sylloge 267.
511 Hisperis sylv, latif. ft. albo parvo Park.
512 Thlassi folio Sonchi lanati Ray Syll.
513 Thlassi ft. albo monospermos Ray Syll.
514 Alfine Myosotis store maximo Ray. Fl. Mart. & Apr.
515 Sedum minus sem. stellato Ray Rupibus & agger. Sepium.
516 Guaicum Patavinum eive similis Frutex. R. Syll.
517 Acetosa Alp. minima, sol. Afari Ambros. Ray 3. p. 115.
518 Hierasium Dentis Leonis sol. acuto, glabrum &c. id. 143.
 67. Muris.
519 Fentophiloides trifol. Alpin. fl. Chameciffi R. 326. 5.
 520 Rola Apart tupina fi inofiffina R. Dendr. 25.

Vide alive Pianta Lucenfes Ray Sylloge 267 &c.

Flanta Pifana.
522 Aconitum faxat, totundirolium Mentzel, pugil.
523 Anchufa paiva, fol. pilof. incanis, fl. parvis Buglossa id.
```

524 Efula minor frut. spinis longis horridis id.

Jacchea mont. max. fol. incan. integris 6 ped. alt. id.
Fluviatilis Pifana fol. dentatis I. B. Ray 121.
Thymus repens Pifana I. B.
Plantæ Infula Elba Nascentes.
Corallina nigra Chama-conchis innata Mentzel.
Echium. max. fol. Buglossa, sl. magnis. id.
Gramen Ægylops umbell. maximum, 5 ulnæ altit. id.
Linaria marit. sl. luteis, rubris, purp. id.
Muscus ruberrimus marinus id.
Kerpillum erect. srut. sl. in spica multislida digestis id.
Thlaspi marit. sol. oblongis strictissimis, sl. albis id.
Tubymalus pediculis storum longissimis id.
Homonitis altera Dalech. Lugd. 1218. Fig.
Lonchitis aspera Ilvensis Lugd. Lugd. 1221. Fig.
Polypodium Invense Lugd. 174. Fig.

Lanta Aquis, Humidis, Ripis, Udis, &c. Nº. 13, 5,

24,7,108, 29, 39, 40, 1,2, 64, 77, 80, 90, 210, 5, 20, 46, 89, 335, 44, 55, 9 & 62, 93, 403. Palustribus 133, 208, 33,395. Maritimis 2, 16, 8, 36, 9, 48, 50, 2, 72, 7, 86, 7, 8, 90, 2, 103, 6, 34, 5, 55, 206, 12, 8, 27, 31, 5, 48, 57, 88, 312, 39, 51, 68, 407, & 16,8,9,92,3,8,529,32,3,5.

Asperis, Petris, Saxosis, Scopulis, &c. 6,38,9,44,54,86,91. 8,101,5,22,58,80,2,8,94,206,12,12,4,6,8,27,50,2,64,8, 72,7,8,81,3,4,8,95,330,1,3,49,52,73,90,9,417,31,3,50,86 Collibus, Montibus, &c. 11, 4, 9, 20, 3, 7, 8, 30, 4, 7, 44, 58, 65, 78, 82, 4, 95, 100, 12, 5, 7, 8, 9, 20, 3, 7, 8, 9, 30, 2, 43, 5 & 8, 51, 4, 7, 63, 5, 6, 7, 71, 4, 7, 93, 5, 7, 9, 202, 7, 11, 3, 8, 23 & 6, 35, & 9, 43, 5, 7, 55, 7, 62, 5, 7, 79, 81, 2, 3, 4, 96, 8, 300, 6, 7, 9, 10, 1, 5, 7, 21, & 4,6,7,30, 3,4,6,41,2,3,6,9,50,3,62,4,7,&72,83,4,8,9, 90,2,3,4,6,7,400,1,5,20,1,2,6,9,30,&44,7.

Maniss, Muris, Parietibus, Ædific. Tectis, &c. 32,53,66, 70,85,102,59,61,85,220,94,9,321,2,8,30,85,6,98. Incultis, Ruderibus, Sterquilinis, &c. 15,53,104,9,26, 42,75,6,87,270,2,2,90,302,3,8,9,10,51,6,76.

Aridis,Siccis,Sterilib. &c. 6,106,51,222,52,60,309,10. Arenosus, Glareosis, Sabulos. &c. 43, 136, 221, 2,8,58,9,73.

Agrorum marginibus, Sepibus, Vepribus, &c. 151,205, Agris, Campis, Planis, &c. 75,107,25,209,30,2,42,80, 97,345,9. Lætis, Pratis, Pascuis &c. 49,68,108,24,70,3411428.

[335,402.

Ericetis, &c. 168.

Arvis, Cultis, Satis, &c. 61,79,81,2,97,107,66,240,381,38,48,54,7,8.
Segetibus, &c. 79,83,111,38,365.

Hortis, Vineis, &c. 51, 62, 126, 79,83,93,217,4,75,6, 321,2,5,32,40. Sylvis, &c. 80,93,6,135,234,316,24, 6,37,44,7,69,77,8,9,89,431.

Opacis, Umbrosis, &c. 30, 102, 19,44,366,7,85,6,96,7,8,404,5. Semitis, Viis, 191,260 Passim, ubiq; &c. 46,153.

Lecis Propriis.

Acuto Monte 202. Apennini Mont. 34,265,341. Argentario Monte 115. Benedicii Monte 421. Gargano Monte 353. Ilva Insula 21,31,54,116,21,82,284,399, Fuliani Monte 78,193, 285,346,68. Agris & Littoribus Liburni 491 & 569. Liguria Mont. 143,66,394.401. Luca 58,342,510 & 210. Pania Monte 346 & 44. Pisa 12,32,134,208,44,69,94,346,95,521 & 7. Pistoriensi Agro, &c. 250, 342. Plombiensi Agro, 9. Senensi Agro 201. Seravalla 264. Urbevetani Collibus 229.

Arbores & Frutices Etruria No. 1&26,33 & 9,516,20. Planta store Cramineo 42 & 54,421,2,51,75,517. Lactescentes Fl. pleno 318 & 6. Seminibus. Fl. radiato 311 & 7,50,425,

Agio 291. Seravalla 264. Urbevetani Collibus 229.

Arbores & Frutices Etruria N°.1&26,33&9,516,20. Planta flore

Stamineo 42 & 54, 421,2,51,75,517. Lactescentes Fl.pleno 318&
38,523,4,75,96,518. Papposis Seminibus, Fl. radiato 311&7,50,425,
6,525. Cosymbifera 301 & 10,482,97. Capitata 340 & 9,51,&
4,427,36. Nodistora & Nudo flore 291 & 300. Umbellifera 114,
& 50,339,437,52. Stellatis foliis 151 & 9, 429,30. Asperisolia
258 & 62,431,2,523,30. Verticillata 110,263 & 90, 420,38,9,40,
98,527,34. Nassiona 111, 66 & 73,7 & 81,225,388,94,418,933,
50,522,32. Nudo semine 355 & 75,92,3,428,46,9,57,8,66,7,8,86,7,8,
66.7,8,86,7,8,515,9. Testapetala 112,60,1,82 & 20,417,345,41,
2,3,53,70,1,9,511,2,3,35. Pentapetala 12,60,1,82 & 20,4417,345,41,
2,3,53,70,1,9,511,2,3,35. Pentapetala 99,& 112,62,3,4,74,5,6,217
& 22 6 & 9,447,54,69,77,80,1,3,9,90,514,9,33. Pissiona 79 & 98,
113,460 & 4,72,3,484,99,592. Bulbosa & affines 223,4,30,50,7,455,6,50,87. Anomala 27 & 32,40,1,78,195,205 & 16,444,57,455,6,50,87. Frumentacca 55 & 77,531 Planta minuspersectar 395,
& 416,465,92,3,526,9,33,7,8,9.

PETIVERIANA seu NATURÆ COLLECTANEA;

Domi Forisque Auctori Communicata.

AJACOBO PETIVER, Regia Societatis Socio, &c. LONDINI.

Ivers Shells, Fossils, Minerals, Marbles and Plants found about Lisbon, and Communicated to me from thence, by the most Curious and Learned Don Pedro d'Almeyda, Son of the Noble Count d'Azumar. 1. Pecten Lusit. minor, friis plurimis. 1. Petter Lajar. minor, juris prarians.
2. P. L. min. compressive, medio extus cavus.
3. Chama Medit max, percrassa Nob. Bon. 106, sig. 59. Lift.
4. Hist. Conchyl. Tab. 414. f. 258.
4. Chama minor tenuis ferè unescens.
5. Edden alba, medio rubro.
6. Chama Lust minima forà Callenga. 6. Chana Lufit, minima, ferè Castanea.
7. C. L. alba, rotunda, perexigua.
8. Purpura rugosa, ore crispo.
9. Aporhais Edinburg. minor, nodosa Gaz, Nat. Tab. 79. f.6.
8. Buccinum tessellatum. cossis pulvinasis 10. Buccinum tessellatum, costis pulvinatis.
11. Rhombus minor fuscus, fascia alba.
12. Rhombus Gibraltar. albus perexiguus.

12. Rhombus Gibrattar. aldus perexigues.
13. Coebleola alba, compressa, parva, ventre pulvinato.
14. • • perexigua ex Anja, 4 mill. à Lisbon.
15. Balanus compressa & squammata Grew 149. Teb. 12. fig. Sect. 2. Fossilia Lustinanica. Or sortugal Fossils.
16. Chama fossilis compression, de Alcantara.
17. Chama fossilis ventricosion, de Alcantara.
18. Rhombus tossilis, claviculà latà altè incisa.
19. Unicornu foss. Luste. orbibus perineiss.
20. • Barcin. orbibus elatis peracutis.

10. • Barcin. Orbibus elatis peracutis.
11. Stalactites Offeocolla facie de Ervedal.

21. Stalagmites Offeocolla tacie de Ervedal.
22. Stalagmites Botryoites è Grotto d'Alcaneda.
23. Chryfiallus alba, è Poco di Lanca, in Alenteio.
24. Chryfiallus pallidè lutescens, id.
25. Chryfiallus anthracinus columnaris, id.
26. Selenites Rhomboides pellucidus, de Alcanede, qr? Lang.
Hill. Lap. Helvetii 27. Tab. 3. Scheutz. p. 49. f.71. 2. & 3.
27. Ibyt-Alumen ruffum de Alcanede.
28. Maymor albescens è Monte Cintra

28. Marmor albelcens è Monte Cintra. 29. Marmor cinereum, micis aspersis, id.

30. Marmor flavescens, lentiginosum, id.
31. Marmor nigrum, Capillis rarisalbis, id.
32. Marmor Alabastrites è Mont. Clares inter Villam Vicosam, &c. Estremos, Provincia de Alenteio.
32 * M.cinereo albedine mixtum. Priore.

33. Marmor carneum, lituris albidis, Lapidicinis de Loures, 3 mill. à Lisbon Copicie.

34 Marmor regelegans, lituris maculisque variis. E Monte de Arrabida, 4 mill, à Lisbon, Sest. 3. Planta Rariores Monte Cintra nascentes, &c.

35. Plantago augustifolia, capit. Lagopi CB. Ray. H. Pl. 878.8. 36. Chamamelum fl. nudo.

37. Abrotanum fæm. fol. teretibus, CB. 136. Ray 359. 1. 38. Cnicus cæruleus asperior CB. Ray 302. Resta Bovis Cintris. 30. Centaurium majus Lusti, fol. integro Nob. Ray 330. 4.

40. Er, ngium pumilum, capitulo cæruleo.

40. Eringium pumilum, capitulo cæruleo.
41. Eringium inarit. Lusit, ampliore sol. R. 3, p. 240. pl. 12.
42. Ocnanche Millefolii sol. Nob. R. 459. 6. Lugd. 718. s.
43. Corallum stavescens, striatum, nodolum Gazophyl. Natur.
Tab. 76. sig. 7. de Ericera, 4 mill. à Lishon.
Sed. 4. African Shells and Drugs from Pernambuc, &c.:
44. Chama Pernamb. maj. trequetra, costà alatà Coch. Pern. f. 35.
45. • tenuis minor pellucida, serè rubelcens.
46. /unis marina vulg. sacie, è Marzagan Africæ.
47. Nautilus prætenuis, dor so bidentato Nob. List. T. 556.
f. 7. opt. q.? Bon. 91. f. 13.
48. Neritulæ Pern. variæ varietates vid. Gaz. Nat. Tab. 67. &8.
Sed. 5. Drugs from St. Jago one of the Cape Verd isles, &c.

48. Nericulæ Pern. variæ varietates vid. Gaz. Nat. 1 av. 07.00. Sell. 5. Drugs from St. Jago one of the Cape Verd isles, &c. 49. Indigo Officinarum seu Anil Ray. 926. c. 4. Jam. Cat. 141. 50. Anacardium Occidentale Offic. R. 1649 c. 15. Jam. Cat. 148. 51. Cassia sistula Offic. Ray 1746. c. 8. Jam. Cat. Plant. 145. 52. (assia sistula Brasiliana CB. 403. 2. R. 1747.2. Jam. C. 146. 53. Cela amargo sa Lusit. Arbor permagna, certica valde amaro, Seminibus Olei plenis.

54. Spinho Beto Lusit. Acacia siliq. falcatis cinereis, velutinis. 55. Mammee. Arbor Indica Mammei dicta C B 417, Ray 1665.

c. 14. Pluk. Tab. 268. f. 4. Jam. Cat 179.

56. Ouri ou Avila. Incolis. Bonduc cincerea Nob. R. 1713. c.3.

Ouri ou Avila. Incolis. Bonduc cincerea Nob. R. 1713. C.3.
 Semente de Purga Luste. Ricinus Amer. major. sem. nigro CB. 432. Ray 166. 2. Jam. Cat. 41.
 Tavacom Incolis. Coral filiq. articulatis, sem. coccineis.
 Tamarindus Offic. Ray 1748. C. 9. Jam. Cat. Plant. 147.
 Tortollo Incolis. Arbor venenata lattifera, sem. catharticis.
 Jacapucaya Maregr. 128. sig Zabucayo Pist. 66. s. edit. 2. st. 155. sig. opt. Ray Hist. Plant. 1677. C. 7.
 Pareira brava Pomet. I. 2. p. 68. sig. Wild Vine, Hist. of Drugs 37. Tab. 15. st. 14. A Specifick in Stone. H. Acad. R. 1710. 8vo. p. 73.
 O M. II. Shells, Fossis and Corals, sent me from the Reverend F. Philip Bonann, S. I. at Rome.

tend F. Philip Bonann, S. I. at Rome.
63. Pecten Mediter. maj. costis altè striatis Nob. Bon. 98.
f. 3. & 4. List. H. C. Tab. 165. f. 2. Cappa Veneta Italis.
64. Spondylus Medit. externe ruber mucricatus, Nob. Bon. 100.

. 20 & 1. Scaraponzelo Tarentinis.

65. Pedunculus max. craffus levis fere radiatus, Muf. noft. 833.

A. Camadie Tarentinis, Bon. 107. f. 64. 5. 66. Pectunculus Neap. minor crassus, maculis runs, G. N. 52.9.

67. P.N. striatus, ex uno latere serratus, ex altero verruculatus.

68. Lunata Neapolitana minor tessellata

69. Ead. fusca radiata.

70. Lunata Neap. lucens crassa, parum cuneata. 71. Auris rugosa è Sicilia.

72. Cochlea Sicula umbilico aurito, mucrone compresso.

73. Trochus Tarent. catenis rubris quadratis.

74. T. Tar. spiralis, caten. eleganter marmor. & maculatus. 75. Neritula Pernambuc, maculata s. liturata varia.

76. • • Mac. s. liturata fasciata varia.

• • Undulata varia.

78. Buccinulum Medit. gibbosum, ventre albo calloso sessili, 1 ist. 971. f. 26. qr? Bon. 134. f. 175. f. 167. f. 383.
79. • cancellatum, lingula cærulea G. N. 52. f. 13. ex An-

80. B. Corficum exiguum, rubescens, marmor. lucidum.

 B. Corneum exiguum, rubeicens, marmor. lucidum.
 Unicornu minus, costis eleganter elatis Mus. nost. 705.
 Unicornu minus, costis eleganter elatis Mus. nost. 705.
 Unicornu minus, costis eleganter elatis Mus. nost. 705.
 Concha Vener, maj. maculata, ventre albo è Mozambique.
 C. V. Media, Castanea, è Pantalera, G. N. 52 10. C. 151.
 C. V. affinis, alba, parva, ex Alexandria Ægypti. Pourcellanes en Coquilage Pomet. pt. 2. p. 104. c. 49. fig. 97.
 H. Drugs. 202. T. 86. fig. 40. H. Drugs, 302. T. 86. fig. 49. 86. Endem minor.

7. Murex Medit. aculeis rigidis brevibus, G. N. 68. 13.

88. Purpura Medit. rugosa, fasciis Castan. & alb. transsucidis, List. 952. f. 1. qr? Bon. 150. f. 23. 89. Iursa Medit. fasciis eleganter nodotis.

90. Unicornu fossilis, fasciis regular. nodosis. E Monte Pelia prope Urbem vetereni Roma.
91. Unicornu fossilis, fasciis regular. verrucosis. Priore, vid. Buccin List. H. Conchyl. Tab. 1019, f. 82.

92. Corallium Album oculatum, ramonifimum. 93. Corallium albescens, ramosum. id. Pet. 153.

94. Epi-Corallum Ericoides album.
95. Retepora, Escara marina Imper. 625. & 30.f. 2.R.65.c.3.1. COM. III. Crustaceous & Testaceous Animals, sent me from

the Curious Mr. John Salvadore Apothecary at Barcelona, of. Cancer Crista Galli dicta Nob. Cancer Ussus dict. Rondel, Murane ou Ours edit. Gallice, 404. f. C. Hera-564, fig. Migrane ou Ours edit. Gallice, 404, f. C. Hera-cleoticus Aldr. Exang. 199, f. ed. Francof. 61, f. 15, Gal-lus marinus dist. à fimilitudine Chelarum.

97. Cancellus chelis eleganter rugosis, Bernard l'Ermite Rond. 553. f. Brancha seu Brancha Lig. bious cambus Provinc. 98. Squilla lata Rond. 545. f. Gall. ed. 392. f. Gefn. aq. p. 215. Urius Marinus Roma Ejusd. 216, Icone Sittardi.

99. Fehinus Catalonicus aculeis longiffimis Nob. Carduus Maris f. Calve Fehinaus & Aurantia maris Bonan. 92. f. 17. This Elegant sea-Uschin, the Curious Calvadore fays, is very frequent on their Coast: ; some of its Needles are near 4 Inches long, very taper, and not thicker than a Common Ruth.

100. Idem nudus, orbiculis majoribus Nob. Bon. 92. f. 18. It's the large Orbicular Veflique of this Shell, that we meet with Fosfill both here and in Switzerland.

101. Uva marina. Salvad. qt? Sepi rrum ova Rond. 504. f. 102. Pecten Barcin. minor, rubetcens.

103. Sphondilus max. tuber, incus albus, qr? Rond. aq. 41. f. 104. Glycymeris Medit. maj. percrassus, leviter striatus. 105. Glycymeris zonis ferè faciatis. The Girdles are most scen

in the largest Shells; for whilst small they are more Mar-bled, and the inner Edges finely notch'd.

106. My tulus Medit. vulg. Nob. Concha Rhomboides Bou. 103. f. 32. Rond. 29. 72. f. Gefn. aq. 267. edit. Ic. 227. f. 107. Peclunculus Barcin. medius triquetrus. The broad Part

of this Coole, especialy the In-lid is purplish.
108. Lunata Medit. fasciis valde rugosis.

109. Purra Medit. major, eleganter serrata. 110. Tellina Medit. Olivacea, interdum radiata.

III. Musculus Liburn. oblongus, teres, qu? Dadylus Bon, 102. f. 28. Solea Venetum Gefn. Icon. aq. 242. f.

112. Musculus Barein. ore hiato.

112. Musculus Barcin. ore hiato.
113. Pinna Medit. permaxima muricata Nob. qr? Pinna aculeata Aldr. Exang p. 533. fig. ed. Francos. 176. Tab. 16. f. 10. Bon. 101. f. 24. List. 372. f. 213.
114. Dentale Anglicum minus G. N. 65 9. Cat. 8z. List. H. C. Tab. 547. f. z. Dentalium Offic. Dale Pharm. 546. 1. I shall be glad to see the Worms in this Pipe-Shell.
115. Cochlea Hort. Angl. similis, sed compression. Very like G. N. 65. f. 4. but much larger.
116. Cochlea terrest. Fibularia Burcinonensis. qr? Trochus 1. Rond. aq. 92. f. but its more like the Shell as Gesner has sovied it in his Icon. Aquat. 240. f. ult. being there much

copied it in his Icon. Aquat. 249. f. ult. being there much rounder and blunter.

117. Cochea calata Bon. 114. f. 11. 2. & 13. Rond. aq. 98. f. Scaragol vel Cagarolo di mar Hispanis & Monspel.

118. Ejus Operculum G. N. 65. f. 5. 6. Umbilicus marinus Langii H. Lapid. Helvet. 116. Tab. 50. fig. Bon. 114. f. 14. 110, Buccinulum Barcin. Pyramid. nigric, ore elato albo.

· pellucidum, mucrone obtufo. 120.

120. • pellucidum, mucrone obtulo.

121. • Medit. læve, ex albo nigroque variegatum G. N. 45.14.

122. • Gibralt. mac. albis carenatis, fafciå pellucidå.

123. Eursa nodosa Pet. 89. Echinophora Bon. 115. s. 19. List.

1003. f. 68 opt Aldr. exang. 399. fig. reverå.

124. Eursa Medit. major, fascis sulcatis. qr? Bon. 115. s. 20. s. 21. Rond. aq. 83. s.

125. Bursoides Maurit. nodosa, ore croceo. qr? Bon. 115. f. 18.

List. 957. f. 9.
126. Trompeta utriusque Indic. qr? Buccinum Rond. aq. Et. f.

reversa Id. Gallice 52. f.

127. Trompeta rugofa minor, ore crispo. qr? Turbo magnue Rond. aq. 88.f.

128. Unicornu Medit, muricatum Lift. 1019. f. 28. qr? Aldr. exang. 354. f. 3. edit. Francof. 120. Tab. 7. f. 5. N. B. Almost if not all the Shells in both the Latine and French Editions of Rondelet have reverted Figures, viz. their Mouths to the Right, which should be to the Left, as they are copied by Aldround and colner. they are copied by Aldrovandand esser.

Barcelona Fossils.

120. Ostrea Barein, hine inde verrucosa.

130. Offrea crassa & prælonga, eardine longissimo. This re-fembles the Virginia Rock-Oyster, List. 201. s. 34. 131. Pecten Medius Fossilis, vid. G. N. 94. f. 1. 132. Peden minor, striis tenuioribus: vid. Mus. nost. 830.

133. Echinites minor, costis 5. verucosis. These are the Size of a Clock-Button, round and somewhat compress like Plot.

2. Hift. Oxon Tab. 5. f. 6. but doubly bigger

134. Vonica fossiiis major. These are smooth and convex on both Sides, the Largest rarely exceed the Dimensions of a

Shilling, or the Thickness of an Inch.

135. Vontica hæmispherica, circulis subtus capillaceis. This feems to be the last split, whose inward Superficies are composed of circular Fibres, like Scheutz. Lith, Helvet. sig. 48. 136. Vomica parva convexs. Thefe exceed not a finall Lupine

Seed.

137. Vomica compressa, tenuis minor. Very thin, and much

factor than the last, but double its Dimensions.

138. Bradeola paiva pileata fossilis. Not broader than a common Spangle; concave on one Side, and convex on the other, rising in a Point.

139. Trompeta major fossilis. Like 126. It's a Span long, and near a Foot about.

140. Veneroides fossilis fasciata. Like G. N. 97. f. 23. but this has furrow'd Girdles.

141. Rhomhus minor, pyramidalis vide List. 729. f. 14.
142. Rhomhus minor patulus, clavicula cavà vide L. 1750. f. 44.
143. Unisornu fossilie Nost. 20. like Mus. Pet. 738. A fair and elegant Fossill.

1.44. Unicornu fossile maj, inthis albo vitreatum. Like List. 521. f. 59. it's somewhat lirger than the last, with many small saleiæ, if not defaced, the Oths blunt or roundish; it Shines within like white glazed Ware or China.

145. Idem, Fluorescens, matrice Argillaca. Seems the same with the last, but composed of Spar, in a hard elayer

Bed.

146. Marmor flavescens Barcin, dendrite. This bas the fame Mos-like Ramifications as are in our Oriental Mo-

147. Ferri minera Catal. rubiginosa & quasi exusta.
148. Plumbi minera Barcin. albo Fluore mixta.
149. Armoniacum Catalon. fossile. Its upper B Bed Striated like Talk, the rest heterogeneous and more glittering. 150. Stalactites Thuroides. Like Franckincense, rugged and

yellow without, but within whitish.

Corals and other Mediterranean Submarines.

151. Corallum albidum Medit. nodulosum. Like our small Coral, but this grows in Clusters and much lar-

152. Corallum implexum, tubulis intus & extremitatibus sellatis.

153. Corallum chelis brevibus obtusis punctatis.

154. Epicorallum Muscosum. An Incrustation over some fine Sea-Moss.

155. Reticulum Tophaceum marinum Chabr. 577. Ic. 5. I. B. 3. l. 30. c. 54. f. Imper. 642. f. opt. Ray. 65. 3.

156. Rihes marina. From the Nature of their growing in Clusters like Currans, perhaps an Ovary.

157. Bursa marina CB. 368. 12. Alga Pomum Monspel. I. B. 3. 1. 39. p. 3. Pulmo marinus Rond. aq. 131. 28. f. Gesn. aq. Icon. 266. f. Ray 83. 8. I. R.& Chabr. have figured this on the lest Sides of the Root of the Alga angustif Vitrariorum, which others have copied without mention-

158. Pila marina Medit. vulg. G. N. 64. f. 4. Cat. 157. Chabr. 570. Ic. 1. I. B. 3. l: 30. c. 4. p. 783. Ray 83. 9. It's like a rough Maw-Ball, made up of the hairy Fluse about the Roots of the Grass-wrack. &c.

159. Manus marina Gefn. aq. 524. fig. edit. Ic. 273. Chahr. 572. Ic. 4. I. B. 3. I. 31. c. 31. f. Ray 77. 7. This is like Geluer's Figure which Cornelius Sittard fent him. N. B. The Fingers in our Specimen are full of final feedy Warts as in our common Wrach or Quercus maritima. This elegant Sea Body Dr. Salvadore fent me growing to the Pectunculus Echinatus.

160. Sertolaria Imper. 653. fig. opt. Chabr. 572. Ic. 3. Opuntial marina Park. 129. 4. f. 12. Ray 77. 5. Sea Band-

ftrings.
COM. IV. Barbadoes Trees, whose Specimens were collected there and brought me by Mr. Jam. Rheed. Arbores Conifera

there and brought me by Mr. Jam. Rheed. Arbores Conifera & Julifera. Trees bearing Cones, Catkins, &c.

161. Button Tree Pluk. Tab. 204. fig. 3. Ray Dendr. xi. 7.

Dendr. Amer. 51. 1. Alni fructu, Lauri folia, Arbor maritima Jam. Cat. 135.

162. Barbadoes Bastard Cedar Pluk. 159. f. 4: Dend. Amer. 51.

3. Cencramidea 3. Alm. Bot. 92. Alni fructu, Loti Arbor. folio Barbadeusis.

163. Barbadoes Cypres Pluk. 197. f. 4. Alm. Bot. 201. pl.

ult. Juniperus Barbad. Cuprest folio, ramulis quadratis.

164. Futick Pluk. 239.f. 3. Alm. Bot. 395. pl. 4. Dendr. Amer. 52. R. Dendr. 14. 3. Morus Barbad, fr. viridi, ligno Sulphureo tinctorio, Jam. Cat. 128.

165. Manchaneel-Apple Jam. Cat. 129. Pluk. 142. f. 4. Dend. Amer. 52.6. Mancinella Barb. arbor Julifera, lactescens, venenuta. P. vi folio.

Amer. 52.6. Mancinella Barb. arbor Julifera, lactescens, venenata, Pyri folio.

Arbores Siliquiseræ. Trees with Pods, &c.

166. Bean-tree Jam. Cat. 142. Coral arbor Clus. 253. s.

Ray. 1736. c. 1.

167. Pidgeon-pea Jam. Cat. 39. Pluck. 213. s. 3. Ray. 1723.
6. Anagyris India utriusque, filiquis catenatis.

168. Indigo Jam. Cat. 116. R. 926. c. 4. Park. 600. s. 3.

Chabr. 82. s. 6.

169. Wild Indigo Jam. Cat. 148. Pajomirioba hexaphylla subrotundis soliis.

subrotundis foliis. 170 PurPurging Cassa Jam. Cat. 145. R. 1746. c. 8. CB. 403. I. 1710. Horse Cassa Jam. Cat. 146. Cassa fishula Brasiliana CB. 1711. 462. 1. Ray. 1747. 2

Horse Callia Jam. Cat. 146. Cassia fishula Brasiliana CB.
463. 1. Ray. 1747. 2.
Spanish Carnations or wild Sena Jam. Cat. 140. Barkadots Flower Fence Ray 281. 10. & Vol. 3. p. 482. 1. Poinciana India utriusque Nob. H. Mal. Vol 6. T. 23. p. 1.
173. Tamarinds Ray 1748. c. 9. H. Mal. V. 1. T. 23. p. 39.
174. Ab Nickars Jam. Cat. 144. Bonduc sem. cæsio Ray 1743. c. 3.

174 c. 3.

175. Fellow Nickers Jam. Cat. 144. Bonduc Iem. cælio Ray 1743.

175. Fellow Nickers Jam. Cat. 144. Bonduc Iem. flavo.

176. Anotto Jam. Cat. 150. Maucaw, or Roucou-tree Ray

1771. c. 21. CB. 410.8. Cluf. Exot. 74. f. Hern. 74. f. opt.

178. Barbadoes Spanish Briar Ray 977. 2. Jam. Cat. 152. pl. 1.

178. Spanish Ash Pluk. 140. f. 3. Ceratia 1. Ray 3. p. 113.

Caroba Barbad. fol. Fraxini.

Caroba Barbad. fol. Fraxini. 179. Spanish Oak Pluk. 141, f. 2.

Caroba Barkad tetraphylla, fl. slam, spicatis.

180. Barbadoes Rot-bean Almag. Bet. 96. pl. 1. qr? Ray 1760.

Jetaiba Barbad. lobis minoribus.

181. Moabite-tree Pluk. Tab. 1. fig. 6. Jam. Cat. 162. Ray 1754. c. 8. & Y. 3. p. 478. pl. 34. id. Dendr. 113. Int-fia Barbad. tetraphylla, filiq. intortis, Jem. nigri luci-

162. Dogs-prick Pluk. 172. f. 4. Alm. Bot. 126. pl. ult. Jam. Cat. 153. pl. 5. Ray D. 102. 3. Dendr. Amer. 52. 9. Caroba Barbad. Laurifolia, filiq. mucosa miniata. 183. Black Sallow Pluk 221. f. 2 & Alm. Bot. 328.

183. Black Sallow Piuk 221. 1. 2 & Alm. Bot. 328.

Breynia Barbaden f. tetraphylla.

184. White Willow Pluck. 221. 1. Jam. Cat. 153. pl. ult. Ray
D. 100. 5. Brennia Barbad. Oleastri folio.

185. Barbadoes Jessemy - tree Pluk. 207. 2. A. B. 109. pl.
3. Jam. Cat. 154. 3. Ray D. 114. 4. Dendr. Am. 52.9.

Nerium Barbad. arbor. fl. incarnato.

Nerium Barbad, ardor, tt. Incarnato.

186. White-wood Fluk, 204, 3, & A. B. 215. Jam. Cat.

154. pl. ult. Ray D. 114. 2. Dendr. A. 52. 12.

Nerium Barbad. Altum, folia Laurinis palmatis.

187. Cabinet or Prince-wood Fluk. 239. 2. & A. B. 395.

Spanish-Elm, or Prince-wood Jam. Cat. 155. Ray Dendr.

115. 6. Dendr. Am. 53. 15. Nerio affinis, Lauri folio,

fl. luteo amplo. fl. luteo amplo.

188. Swamp Mangrove or Oyster-tree Pluk. 204. f. & A. B. 241. Jam. Cat. 152. 2. R. 1772. 1. Mangle Barb. fol.

Pyri rigido.
189. White Mangrove Pluk. 204. 4. & A. B. 241. Mangle Barbad. fol. Caroba advertis.

190. Silk-Cotton Pluk. 188. f. 4. & A. B. 172. H. Mal. 3. T. 49. p. 59. Ray 1064. 2. & 1899. pl. 5. & 6. Gossipium India utriusque, sol. digitatis planis.

191. Barbadees Cotton-bush Pluk. 188. f. 1. & A. B. 172.

Goffipinm Barb. fol. trifido.

192. Musk Mallow Pluk. 127. 1. Jam. Cat. 98. Ray 1067.
Bamia Barbad. histura, fem. Moschato.
193. Ocra Jam. Cat. 98. pl. 4. Alm. Bot. 16. pl. 5.
Bamia fructu efculento, brevi decogeno.

194. Long Ocra Jam. Cat. 98. pl. 3. Alm. Bot. 16. pl. 6. & Mant. 7. Bamia fructu efculento, longo, recurvo.
195. Dogwood-tree Jam. Cat. 143. Mant. 150. pl. 2. Ray D.
198. 9. Bagua frudu membranulis alato, Laferpitii

Arbores frudu maximo. Trees with large Fruit. 196. Coco-tree Jam. Cat. 132. Park. 1596. f. Dendr. Amer. 2. p. 50. Palma Indica Nucifera Cocous dicha Ray 1356. 4. Hort. Malab. Vol. 1. Tab. 1. &c. p. 1. Chabr. 27. f.

4. Hort. Malab. Vol. 1. 1ab. 1. de. p. 1. Chaor. 27. 1. 4. & 28. f. 1. CB, 508. 1.

197. Cashew Fam. Cat. 187. Fark. 1568. f. Anacardium Occidentale Nob. Ray 1649.c. 15. H. Malab. V. 3. T. 54. p. 65. Chabr. 24. f. 6. CB. 510.

198. Banana Fam. Cat. 192. Dend. Amer. 14. p. 51. Banana stuctu maj. recto Nob. Ray. 1375. 2. Piso 76. ed. 154. f. 2. CR 208. CB. 508. 1.

 Plantain-tree Jam. Cat. 189. Ray 1374. Park. 1496. fig. Muja fructu adunco angulofo Nob. Pifo 75. f. ed. 154. f. 1. H. Mal. 1 T. 6. p. 17. Chabr. 9. f. 5 & 6. p. 10. f. 1. CB. 507. XI.

200. Red Guayava Jam. Cat. 198. Park. 1671. Guayava fru-Ru rubro majore Nob. H. Mal. 3. T. 35. p. 33. CB. 347.

Kay 1455.

101. Papaw-tree Jam. Cat. 202. Ray 1370. c 1. & 2. Park. 1049. f. 5 Papaya major mas & femina Nob. H. Mal. 3. T. 15. p. 21. &c. CB. 431. 2. & 434. 12. & 457. 4. 202. Custard-Apple Jam. Cat. 204. Pluk. 134. 4. Anona augulisticia, frudu majore luteo Nob. H. Mal. 3. T. 30. & 1. p. 2. R. Dendt. 77. 2.

203. Sower-sop or Prickly Apples Jam. Cat. 203. Anona fru-Etu spinulis aspero Not. Rav. 1651. 5. & Dendr. 77. 5. Pluk. 135. f. 2. 912 id. 134. f. 2.

204. Sweet-sop Jam. Cat. 205. 1. Ray 1650. 1. Dendr. 77. 1. Pluk 134. f. 3. H. Mal. 3. T. 29. p. 21. Anonis fructu squamoso dulci Nob.

205. Water sop or River-Apple Jam. Cat. 205- 2. Pluk. 240. f. 6. R. Dendr, 78. 1. Anona aq. fol. Laurinis, frudu minore luteo.

206. Garlick-lear Jam. Cat. 205 Ray 1644 c 3. Dend. 79. 7Tapia arborea triphylla Flum. E. B. 22 T. 21. Pet 4 f. 22Pife 69. f. ed. 140. f. H. Afal. 3. T. 42. p. 4.

207. Arbor Cheurbitifera Parbad fol temiibus apiculatis Nob. Cujete Flum F. B. 23 T. 16 Pet 2. f 14. Pluk. 137. 7 208 Arbor Cueurbit fera, fol crassis consertis.

Citron Jam Cat. 208. Ray. 1654 c. r. Citrus vulg. Nob. Park. 1506. f.

Lemmon Jam. Cat. 209. Ray 1656. c. 2. Limo vulg. Nob Park 1507 f

211. Lime Jam. Cat. 211. Limo sylv. sructu parvo subacido Nob

212. Orrange Jam Cat, 210 Aurantia vulg, Ray 1658c, 3. P. 1508 f

213. China Orange Jam. Cat. 211

214 Shaddocks Jam. Cat 212 Pompelmo's Alm. Pot 39. Ray Dend. 80. 1. Aurantium fructu maximo pallescente

Arbores frustu medio. Trees with midling Fruit.

215. Cabbage-tree Jun. Cat. 176. Ray 1361. 12. Dend. Amer.
1. p. 50. Alm. Bot. 275 pl. nlt. Palma Regia Earbadenfis.

216. Palmetto Jam Cat 177. Ray Dendr. pl 2-3. D. Amer. 7. p 50 Palma Barbad, non spineta, fol plicatili feu Flabelliformi, frudu officulo nigro Alm Bot. 277 pl. 8.

217. Balfam-tree Jam Cat. 167. Ray D 51. 4. Dendr. Amer. 35 p. 74. Barbadoes Balfam-Apple Pluk. 157. f. 2. Alm. Bot. 92. pl. 4. Cencramidea Barbad. Balfamifera, fol. roundo crasso. fol. rotundo crasso.

218. Parbadoes All-spice or Mango Pluk. 170 f. 4. Alm Bot. 124 pl. ult. Lycium Cucurbitiferum Barbadense aromaticum

219. Barbadoes Dartwood or Ink-berry Pluk, 97. f. 5. 6. Alm. Bot. 234. pl. 6. & ult. qr? Ink-wood or Berry Pl 205. f. 2. Alm. B. 246. 1. Lycium Barbadense, fol. lucido subro-

220. Barbadoes Cedar Pl 157 1. A. B 92 1 Ligon H Barb 14. 41. & 73 Cedius Barbad fol. Fravini, fr. ficco pentafori

41. & 73 Cectus Barbad. fol. Praxini, fr. beco pentafori Nob. Jam. Cat. 182. pl. ult. R. Dendr. 41. 8. This is known by its peculiar Fruit and Sweet red Wood.

221. Mattick-Tree. Pl. 217. f 5 A. B. 307. i. Jam. Cat. 180 id. Hift 40. Ligon 14. & 73. R. Dendr. 21. p. 42. Prunus Bar-lvad. Lauri folio, Mafficifera.

222. Soap-Lerry. Jam. C. 184. Pluk. 217. f. 7 CB-511. i. Cluf. Exot. 42. f. Lugd. 1877. c. 121. Stapel. 363. f. Quiti Brafil. Maregr. 113. f. Pif. ed. 162. f. Arbor. Saponaria. America-na. Ray. 1548. 2. na Ray 1548, 2.
223. Barbadoes Olive Pl. 209, f. 3 A. B 269, pl. 6, R. D. 47. 6.

Olea Barbadenfis.

224 White Plum, qr? Pl 216 f. 8 Prunus alba Barbad fol. leviter serrato.

125. Barbadoes Plum. Pl. 218 f. 3. A.B. 306. 3. qr? Fam. Cat. 181. Primus Barbad. fol venoso rigido

226 Barbadoes Damozen Fl. 17. f. 6. A. B. 306. pl. ult. Prunus Barbad, fol. tenniore,

227. Barbadoes Black-bud and Black-wood Pl. 205 f. 3 A. B.

227. Barbadoes Black-bud and Black-wood Pl. 205 f. 3 A. B. 246. 2. Jam. Cat 135. Hift 38 D. A. 23. p. 54 gr. Ray 1687. c. 17 Arbor Barbad. Laurocerafifolio.
228. Jack in a Box Pl. 208. f. 1. A. B. 266. 5. Hernandia Flum E. B. Amer 6. T. 40. Pet. T. 3. f. 22. Hernandia fol. cordato amplo umbilicato, fl. hexapetalo.
229. Barbadoes Emeet-wood Pl. 176. f. 1. A. B. 139. 8. Jam. Hift. 38. Evedexylon Barbad. fol. lato, Beretini frumen.

ćtu.

230 Bastard Sweet-wood Pl. 176 f z. A. B 139 pl. ult. qr?
Ray Dendr 23 p. 85. Evedexylum Lauri folio, f. hexapetalo.

231 Fire-Sprig Pl. 175. f. 6. A. B. 138. pl. 7. Ray D. 127. pl. ult. Efcarox/lum Barbad. fol Cacao.
232. Pidgeon-Fig Pl. 178. f. 4. Jam. C. 180 Ray D. 16. 5. Ficus stolonifera, Pyri folio, fr. Cerafi. The Pidgeons much covet this Fig

233 Clammy Cherries Pl. 158. 1. A. B. 94. pl. 15 Jam. C. 169. 2. R. D. 45. 14. Cerafo affinis, fr. coccineo viscoto, fem. rugofo.

234. Bar.

234. Barbadres Cherry Pl. 157. f. 4. A. B. 94. pl. 18. Jam. C. 172. R. D. 74. t. Sinneraka fr. rutro acido sulcato, pelys ermos. Common in Gardens, &c.

235. Chroquer-Grape. Uvifera arbor Americana, fr. atomatico punctato Pl. 237. f. 4. A. B. 394. pl. 7.
236. Mangiove-Grape Pl. 237. f. 5. A. B. 394. pl. 8. Uvifera arbor Baibad. Cacao tol. ftolonibus proliferis,
237. Jungici Mangrove-Grape Jam. C. 183. Ray 1795. I. & Dendr. 40 5. Pl. 236. f. 7. A. B. 394. pl. ult. Guajabara Caribea fol. rotundis rigidis.
Sir Hung Van fays, the Sequinale with an Ison Pin moite.

Sir Hins Sloan, says, the Spaniards with an Iron Pin write, or rather file on these Leaves, when Paper and Ink are

238. Loblolly Pluk, A. B. 38. pl. ult. Gnajakara Barbad, fol. rot. venefis mollibus. This bears its Fruit on a long Spike like the laft.

239. eathercoat Pl. 222. f. 8. A. B 38. pl. 18. Guajabara Barbad, fol. subrotundis venosis ferrugineis.

Arbores frudu minore. Trees with small Fruit. 240. Birch tree Pl. 151. 1. A. B. 67. Jam. C. 167. R. D. 50. 2. D. A. 33. p. 73. Betula Amer. bacca triquetra odore Terebinthini.

241. Box Fl. 139. 6. A. B. 74. 7. Jim. C. 167. 1. R. D. 65. pl. ult. Buxus Barbad. fullu croceo dipyreno.
242. Great yellow Hercules Fl. 239. f. 6. A. B. 8. p. 396.

Fagara Barbad Citri folio. 243. Prickley Tellow wood Pl. 239. f. 4. A. B. 396. pl. 5. qr?

Jam. C. 138. R. D. 70. 15. Fagara Barbad. Arboric
Fusick fol. roundicribus crenis.

244. Wild-Honey-tree Pl. 142. f. 3. A. B. 43. pl. ult. Jam. C. 137. pl. x. R. D. 75. 4. Baccifera arguiè lerrata, fir. mimimo cioceo. In Woods, &cc. 245. Bastard Locust tree. qi? Jam. C. 165. R. D. 49. I. 47? Pl.

200 f. 6. Baccifera Amer. fol. Lauri, f. calyculato monopyreno. Grows in Meadows.

246. Barladoes Sweet-Bay or Pimenta Pl.155. f. 2. A. B. 88.6. Barbad. Lauri folio bacca ovali.

247. Round Barbadoes Pimenta Pl. 155. f. 3. A. B. 88. 7.

Pimenta Barbad, fol. & fr. rotundis. 248. Black-Sage Pl. 221. f. 3. A. B. 329. Jam. C. 163. 2.

Hist. 39. R.D. 31-18. Camara Barbad. Spicata fol. Salviæ.
249. White-Sage Jam. C. 4. p. 164. Hist. 39. R. D. 17. 31.
qr. Pl. 114. f. 5. Camara Barbad. fol. & flor. rotundis.

250. Bitter-wood Pl. 238. f. 4. A. B. 305. pl. ult. Xylopicron Barbad. fr. oblongo calyculato.

251. Fiddle wood Pl. 162. f. 1. A. B. 108. 3 9r? Jam. C. 170, R. D. 61, 1. Citharexylon Barbad Lauri foliis.

5. R. D. 61, 1. Citoarexyton Balbac Lauri Fildle wood Pl. 162. f. 2. A. B. 108. 5. Citharanyon Balbac. Lauri feliis. 2 ferè minoritus.

253. Fiddle wood with dented Leaves Pl. 161 f. 5. Cithar-exylon Barbad. fol. dentaris.

254. Lightwood Lignum Rhodium or Rosum Pl. 201. f. 3. A. B. 228. 2. Jam. C. 137. 2. R. D. 88. 30. D. A. 55. pl. 34. Baccifera Barkad. fol. Betule planis & perfo-

255. Milk-wood, Pl. 185. I. A.B. 164 Jam. C. 136. Hift. 39. R. D. 87. 27. D. A. 54. 31. Galastoxyium Barbad. Laurocerafi folio majore. This yields a Poyfonous Milk from all its Parts, and hoils into Birdlime.

256. Po, fon-wood Pl. 230. f S. A. B. 369. pl. 10. Jam. C. 136. pl. ult. Hist. 39. R. D. 87. 28. D. A. 54. 32. Ligon 68. Tithymalus Arbor Barbad, Citri fol. leviter cienatis, Common in A cadows.

257. Red wood Pl. 175. f. 5. Alm. Bot. 138. pl.6. Erithroxylum Barbad. Hydrolapathi foliis.

258. Rod wood Pluk, 139, fig. 6. Alm. Bot. 46. pl. 6.
Myrtus Barhadenfis vininalibus virgis.

259. Spirit - wood Pluk, Tab. 215. f. 5. Alm. Bot. 300. pl. 3. Pneumatoxylum Barbad. Face calyculata tetrasperni. Arb. Barbad. flor. f. fruct. fere ignotis.

Barbadoes Trees, whose Flowers or Fruit are little known. 260. Barbadoes Dogwood Mautis, Almag, Botan 150, pl. 2 Pif ieula arbor Barbad, fol. rotundis.

261. Black Dogwood Pluk. 172. f. 6. Alm. Bot. 127. pl. 2. Cynoxylum B.n bad, nigrum, Laurocerafi folio.

262. Barbadoes Iron-wood Pl. 224. f. 2. Alm. Bot. 346. pl. 12.
Sideroxylon Babrad. foi. apiculatis.

263. Whip wood Pluk. Tab. 298 fig. 2, Alm. Bot. 395. pl. 8.
Aylonaftix feu Flagellifera arbor Barbadensis

264. Rat bane. A poyfonous Tree Flowers like Honey suckle.
Arbor Barbad. Fagi foliis pulchre venosis.
265. Rusty bough Pink. Tab. 1.44. sig. 4.
Arbuscula Barlad. fol. seringiness digitatis.

266. Barbadoes Fingrigo or Savin Pl. 107. f. 4. A. B. 219. 2. Fingrigo Barbad. fpinofa, parvis fol. & coffic ala-

267. Black-with Pluk. Tab. 145. 4. Alm. Bot. 48. 3. Volubilis Barbad. triphylla, pediculo alato.

268. Broad-leaf Clymer. Pluk. 146. f. 1. Castanea folio Barbad. scandens.

269 Weeping Bower Pluk. 145 f. 3. Volubilis Barbad, aculeis z. ex alis fol. apiculat.

Volubilis Barbad. aculeis z. ex alis tol. apiculat.

N. B. Divers Trees in Barbadoes, Antego, Jamaica, &c. Sprigs of whose Flowers and Fruit, will be very acceptable. Trees with Catkins, viz. * Bastard Cedars * Fusick. Manchineel. Trees with Pods, viz. Bean trees, Purging and Horse Cassa. Spanish As, Elm, * Oak, Sinacod and Moabite Trees, Dogs-prick, Black and white Willow, * Cabinet or Prince wood, White and other Mangroves, * Dogwood, Common and Mountain Ebony, Indian Savin, * Wild Tamarinds. Trees with large Fruit. viz. Banana's and Plan-Tomarinds. Trees with large Fruit, viz. Banana's and Plantains. All Sop Apples and Gourd-trees, * Prickley-Pole, **Thatch and Dragon tree, Anchovy Pear, Mamee, * M. Sapota, * Santa Maria, * Avocado Pear, * Sapotillas, Wild Pippaws, &c. Trees with midling Fruit, viz. Cabbage, * Palm Oil and Palmetto trees, & Great and small Macaw, * Balfom and Mango trees * Cedar, Maslick, Soap-bries and Oilves. All Plums, Jack in a Box, * Fire and Five-Sprig, Rlack and Speer woods Rlack Sour Coubange and other and Olives. All Plums, Jack in a Box, *Fire and Five-Sprig, Black and Sweet woods, Black, Sower, Coubange and other Cherries. All Grapes, Leather-coat, Loblolly, &c. Trees with Berries and other small Fruit, viz. Birch, Box, Herenles and yellow Woods, All Spice and sweet Bays, Honey-tree, Bastard-Locus, Hog Doctor and Boar-tree, * Cloven and Poyson berries, Bichy-tree, * Olive bark and wild Jasmins. All Soits of Sige Bitter-woods, Fiddle-woods, Iron and Light-woods, Milk and Poyson-woods, Red and Rol-woods, * Spirit-wood, * Silver wood, &c. N. B. Those with this * Mark, will be doubly acceptable to J. Petiver.

CO M. V. Trees and Herbs, sent me from the Isle of Providence, by that Curious Gentleman Capt, William Walker, Chief of the Bahama or Lucay Isles.

270. Brasiletto Djing-wood.

Brasiletto arbor spinosa, foliis Buxeis.

271. White Cinamon Pl. 160, f.7. Jam. 165, ult.
Winteranus Offic. cortice albicante.

272. Wild-Grape. Bears a small red Fruit, which makes good Wine. Walker. Vitis Provident. fol.altè dentatis.

273. Lignum Vite. Jam. Cat. 186. Ray 1685. R. D. 85. Pl. 33. f. 3. 4. A. B. 180. CB. 448. 1. 2. Cluf. exot. 332. f.

uaicum Amer. Buxi foliis. 274. Madera Timber. Jetaiba Petiv. Nat. Collect. 180.

275. Moho-tree Pl. 178. f. 3. quoad Fol. A. B. 144. 8. Mant. 175. Fam. C 95. R. D. 319. 16. Ketmia utriulque India, Tiliæ folio subrus albicarte.

276 * weet Bark. A Tree W. gr. Pl. 220. f. 5. Elutheria Provid. fol. cordato lubius argerneo.

276. Saffron-tree. Celtis Amer. fol. Citri subids aureo, frudu rubro Instit. Rej. Herb. 612

277. A small Tree unknown Walker. Lilac Provid. Mytti fol.

flore tubulofo stamineo. 278. Sea fide meed. Bears a red Berry, which will make Ink or dye a Purple, Walk, Solanum racentofum Americ, minus Jam. C. 85. Pl. 112. f. 2. A.B. 353. pl. 3: Ray V. 3. P. 349. pl. 1.
279. A Plant made use of for Burns and Scalds W. Chrysanthem: fourth fronting.

themi forte species.

COM. VI. My Hearty Friend, Mr. John Partridge Apothocanear the North Pole.

280. Coralloides Amerie. lignosa, pereleganter verrucosa. 281. Corallia marina squamea, squamis deorsum spectantibus Lang. H. Lapid Helvet. p. 57. T. 17. f. 2. Planta mariana Refede facie Clus. Exot. 123, f. Chabr. 576. lc. 5. Park 1302. f. 4. Refeda lapidea CB. 100. 5. & 365. 18.

Ray 68, c. 6.
COM. VII. The Three following Animals from Buen Ayres, were presented me by my Neighbour Mr. William Smith,

282. Armadilla valde pilesa, fronte magno scutato. rate Account and Figure of this rare Creature we hope fuddenly to see from that expert Anatomist Dr. Ja. Douglas, F. R. S. &c.

283. Columba marina palmires, Call'd there Pintado.

Captain George Jesson caught with a haited Hook.
284. The Spur-wing d Lapwing, Vanellus humeris calcari armatis. Mt. John raploe shot this 50 Leagues up the Rio Plata, on the South Shore.

PETIVERIANA II, seu NATURA COLLECTANEA; Domi Forisque Auctori Communicata.

AJACOBO PETIVER, Regia Societatis Socio, &c. LONDINI.

64 Festuca Aven. sterilis humilior, 91. CB. 10. 8. R. 1289.
65 Festuca Graminea, estusa jubă, 91. B. pr. 19. 65. R. 1290.10.
66 Festuca Graminea, glumis histutis, 91. CB. 9. 1.
67 Gramen minimum, 87. D.d. Lugd. 425. f. R. 1279. 15.
68 Gr. nemor. panic. albis capillaceo fol. 33. B. pr. 14.45 Th. 97. 1.
69 Gr. prat spică purp. ex utriculo prodeunte, 273. Lugd. 425. f.
70 Linagrosis paniculă minore, Par. 486.
71 Genista ramosa fol. Hyperici, 451. CB. 395. 4. Lugd. 173. f.
72 Pinus marit. altera, 351. Matth. CB. 492. 6. Lob. Ic. 229. f.
73 Salix platyphillos leucophaus 357. Lugd. 257. f.
Plante PARIS, quarum Specimina seca vel semina recentia, valde
graia erunt fac. Petiver.
74 Atriplex angustissimo & longistimo folio xi. H. L. Bat.
75 Blitum tubrum minus, 383. CB. 118. Cam. 235. f. R. 200.2.
76 Herniaria hissura, 96. Chabr. 455. f. z.
77 Farietaria minor Oeymi folio, 295. CB. 121. Bocc. 147. f.
78 Persicaria latif. storis calyce candido, Par. 220.
79 Paramogicon ramosum angustifolium, 352. CB. 193. 7.
80 Dens Leonis tenuistimo folio, 331. B. pr. 62. Col. 2. p. 31. f.
81 Hieracium max. glabrum Eruca folio Par. 470.
82 Hieracium histuum hissurum 99. Chabr. 319. f. z.
83 Hieracium muteum hissurum fol. subassero, 341. CB. 127. 5.
84 Hieracium mutor. laciniat. minus pilosum, 97. CB. 129. 2.
85 Hieracium mutor. laciniat. minus pilosum, 97. CB. 129. 2.
86 Hieracium phlomoides, 470. Tab. Ic. 184. I. H. 608. f. 8.
88 Scor zonera angustifolia prima, 143. CB. 275. R. 249. 4.
89 Scorzonera palustris pulverisora, 301. H. R. Paris.
91 Chrysanthemum Alpin. fol Abrotani multifidis, 474. CB. 134.
92 Conyza incana, 178. CB. 265. Lob. Ic. 347. I.
93 Doronicum Plantaginis folio, 433. CB. 184. R. 277. 9.
94 Jacobea mont. lanug. angustif. non laciniat, 474. CB. 134.
95 Scorzonera palustris pulverisora, 301. H. R. Paris.
91 Chrysanthemum Alpin. fol Abrotani multifidis, 474. CB. 134.
92 Sonda pilus alterum fol. serratis Par. 302.
93 Sphondylium hirtut fol. angustioribus, 8c. 365.
93 Aubia tylv. Monsp. major, 528. L. B. 3. 715. R. 480. 2. comunic. I. PLANTS growing wild about PARIS, fent me from that Accurate Botanift, D. 64 Festuca Aven. sterilis humilior, 91. CB. 10. 8. R. 1289. Sebassian Vaillant.

1 Fluviatilis fol. denticulatis, Hist. Pl. Paris 196. Chabr. 56 3.f. 5

2 Herniaria glabra, 96. Chabr. 455. f. 1. Ray 214. 6.

3 Persiaria angustifolia, 220, B. pr. 43. R. 183.

4 Chondrilla Juncea vilcola arvensis, 188. CB. 130. R. 223. 9.

5 Hierac. fol. Ceronopi, 201. CB. 128. Tab.Ic. 181.2. H.607. f. 3.

6 Aster mont. lut. Salicis fol. glabro, 77. CB. 266. 6. R. 266. 1.

7 Conyzis assinis, 8. CB. 265. 7. Lugd. 1282. Fig.

8 Jacobea vulg. fol. instar Eruca vulg. 277. CB. 131. I. var.

9 Carduus prat. latifolius; 259. CB. 376. 1. R. 305. 1.

10 Cyanus repens latifolius; 259. CB. 376. 1. R. 305. 1.

11 Chamenclum maj. fol. tenuist. caule rubente, 416.

12 Bryngium mim. mitius, capitulo magno, 421. H. R. Paris.

13 Succisa hirluta, 141. CB 269. 2.

14 Apium montanum, 216. Dal, Lugd. 702. f. 1.

15 Bulbocassanum maj. fol. Apii, 403. CB. 161. 1. Dod. 434. f.

16 Bustum stifol. adulterinum, 195. CB. 148. 4. R. 473. 1.

17 Heum latifol. adulterinum, 445. Chabr. 383. f. 5.

19 Seseli palustre lactescens, 367. CB. 162. xi. R. 314. 2.

20 Cruciata palustris supina minima Par. 426.

21 Cynoglossum minus 401. CB. 257. x. Colum. 179. f. R.491.5.

22 Lithospermum Linaria folio German. 537. CB. 259.8.

23 Lycopus palustris villosus, 288.

24 Marrubium album villosum, 291. B. pr. 10. 2. R. 557. 6.

25 Bunella 2. non vulg. albo store, 62. Clus. 43. f. 2. R. 551.2. Sebastian Vaillant. 23 Lycopus palustris villosus, 288.

24 Marrubium album villosum, 291. B. pr. 10. 2. R. 557. 6.

25 Buinella 2. non vulg. albo store, 62. Clus. 43. f. 2. R. 551.2.

26 Sideritis hirsuta procumbens, 534. CB. 233. 2. R. 564. 7.

27 Aristolochia Clematitis resta, 175. CB. 307. 5. R, 762. 5.

28 Linaria min. repens & inodora, 209. H. R. Paris.

29 Linaria pumila supina lutea 110. CB. 213. I. R. 754. 8.

30 Melampyrum purpurascente comă, 288. CB. 234. I. R. 774. I.

31 Pedicularis lutea, 294. Tab. Ic. 791. I. H. 2. p. 470. f. R. 769. 2.

32 Viola Martia arboresc. purpurea, 239. CB. 199. 2. 1052. I.

33 Ranunculus aq. Plantag. fol. angustissimo Par. 525.

34 Ranunculus neunorosus luteus, 522. CB. 198. I. R. 625. 3.

35 Ranunculus Oleraceus major, 522. Tab. Ic. 52. 2. H. 142. f. 3.

36 Sedim min. lato & crasso cause Portland, Pelgic. 533 H. R. Par.

37 Alparagus sylv. tenuissimo solio, 56. CB. 490. 2

38 Thlassi Alyston didum campestre majus, 52. CB. 107, 1.

39 Thlassi arvense perfoliatum majus, 233. CB. 106. 7 R. 832. 7.

40 Thlassi ment. luteum, 383. I. B. 2. l. 22. p. 929. f. R. 830. 4.

41 Alfinastrum Gratiola solio, Par. 380.

42 Betonica s. Tunica min. similis planta, 490. I. B. 3. p. 338. Dis, Ch. 199. 1, var.

103 Pulfatilla fol. crassiore & majore, fl. laciniato Par. 131.

104 Quinquefolio Fragisero assinis, 350. CB. 326. 5. R. 616.2.

105 Sedum minus teretifolium alterum, 147. CB. 283. 2.

106 Consolida Regal. arvensis 432. CB. 142. 4. R. 708. 4.

107 Brassica sylv. fol. circa rad. cichoraceis 160. CB. 112. 8.

108 Chamadrys spuria maj. angustifolia, 169. CB. 249. 14.

109 Thlassi Alisson dist. campestre min. 52. CB. 107.2. R, 829.2.

110 Campanula arvensis minima, 252. Dod. 168. f. 1.

111 Caryophyllus sylv. vulg. latifol. 185. CB. 229, 1.

112 Alsine hirsuta minor, 120. CB. 251. 3.

113 Geranium Columb. tenuius laciniarum, 85. B. pr. 138. 2.

114 Lychnis sylv. 113. Tab Ic. 293.2.H. 2. p. 10. f. 4.G. 382, f. 1.

115 Verbascum seem. st. albo, 164. CB. 239. 4.

116 Cicercila sylv. 259. Tab. Ic. 500. 2. H. 2. p. 22. f. 2.

117 Trisol. Fragiser. nostras purp. sol. oblongo 44. Ox. 144. 14.

118 Cynosorchis Militaris minor, 508 CB. 81. 3.

119 Orchis Anthropophora Oreades altera, 349. Col. 2. p. 9.

120 Orchis palmata palust. maculata, 509. CB. 86. 15.

121 Orohis abortiva Violacea, 486. CB. 86. 2. R. 1229. 3.

122 Orobanche major secidissima, Par. 123.

123 Helleborine mont. angust. purpurascens, 200. CB. 187. 7. 42 Betonica f. Tunica mim. similis planta, 490. I. B. 3. p. 338. 43 Caryophillus arv. umbell. fol. glab. 51.CB. 210. 4. R. 1028.8:
44 Centaurium purp. minimum, 412. H. Blaf. 566. R. 1092. 3.
45 Chamzeiflus Erice fol. lut. humilior, 465.CB. 466. 12. var.
46 Ciflus Ledon fol. Thymi, CB. 467.14. Cluf. 81. f. R. 1011.12.
47 Helianthemum fl. maculofo, 94. Col. 2. p. 77. CB. 465. xi. R. 1012. 2. 48 Hyssopisolia maj. latioribus foliis, 529. CB. 218. 13. var. 49 Lychnis sylv. angust. calyc. turgid. striatis, 211. CB. 2053. R. 997. 22.
50 Lychnie visc. st. muscoso min. 211. Tab 820. f. 2. H. 2.492. f. 1.
51 Verbaseum fem. st. luteo magno. 163. CB. 239. 3. R. 1094 2.
52 Anonis lutea sylv. minima, 52. Col. 301. f. R. 958. 7.
53 Fanum Grac. sylv. alterum polyceration, 195. CB. 348. 4.
54 Ferrum Equin. Gall. filig. in summitate, 191. CB. 340. R. 925. 1.
55 Trifolium Cochleat. polycarp. fol. crenatis, 492. CB. 329. 4.
R. 964. 21. So Irifolium Cochicar. polycarp. jot. cicinatis, 472. 62. 5-7. T. R. 964. 21.

So Trifolium mont. alb. 540. CB. 328. 5. Cluf. 245. f. 1. R. 943. 6.

Trifolium mont. alb. 540. CB. 328. 5. Cluf. 245. f. 1. R. 943. 6.

Trifolium mont. alb. 540. CB. 328. 5. Cluf. 245. f. 1. R. 943. 6.

Milium fylv. tenuifolium, 413. Lob. Ic. 193. 2. R. 1117. 4.

Orchis fem. procerior majori flore Par. 508.

Orchis palmata Batrach. 509. CB. 86. 10. Lob. Ic. 133. 1.

Orch, palm. paluft. 292.111. CB. 86.14. Lob. Ic. 1932. R. 1225. 3.

Orchis palm. paluft. lat. 509. CB. 86.12. Dod 241. f. R. 1225. 1.

Thatifrum, 5. vel prat. 1. Cluf. 234. Par. 537. 122 Orobanchè major fœidiflima, Par. 123.

123 Helleborine mont. angust. purpurascens, 200. CB. 187. 7.

124 Helleborine fl. albo vel Damasonium mont. lat. 340. CB. 187. 5.

125 Tithymalus mont. non acris, 537. CB. 292. 3. R. 870. 34.

120 Tithymalus palust. fruticosus, 368. CB. 292. R. 851. 8.

127 Valeriana sylv. major altera, folio lucido, 237. H. P. Par.

128 Triehomanes sol. elegant. incists, 535. T. Ic. 797. 1. H. 474. f. 3.

129 Genista alba, 430. Tab. Ic. 1100. 2. H. 2. p. 802. t. 2.

130 Quereus sol. modi lanugine pubescentibus, 352. CB. 420. I.

COM.

```
COM. II. Bafil Flants fent me from Dr. Zuinger, Physician in
                                                                                  that City.
    131 Polygonum littor, min. Flosc. spadiceo albicantilus, 83. B.
   pr. 131. 3. Ray 213. 2.

132 Hieracium Chondrillee fol. glabrum 37. CB. 127. 3.

133 Hieracium Chondrillee fol. glabro, rad. Succifa majus, 37.
   CB. 127 4.
134 Hieracium Chondr. fol. glabio, rad. succisa minus 38. CB.
                   128.5.
    135 Hieracium Chondr. fol. hirfutum 37. CB. 127.2.
    136 Hieraeium Dent. Leonis fol. obiulo, minus 37. B.pr. 63.3.
    137 Hieracium minus glabrum, fol. eleganter virentibus, 37.B.
                   Dr. 63. 2.
    138 Hieracium muror, lacin, minus pilosum 38. latifolium.
    139 Hieracium muror. lacin. minus pilosum, 38. angustifol.
   140 Pilofella major erecta, CB. 262. 2. R. 241. 16.
141 Pilofella major erecta, minus hirtuta, CB. 262. 3. R. 243. 3.
142 Carduus in Avena proveniens, CB. 276. 8.
143 Chamadrys spuria minor, angustifolia CB. pr. 117.5.
    144 Melampyrum luteum angustifolium, B. pr. 112.
    145 Alcea vulg. minor, CB. 316. 2.
146 Sedum Alpinum hirtuum luteum, CB. 284. xi.
  147 Polygonatum angustifol. non ramosum, CB. 303. 8.
148 Papaver Erraticum minus, CB. 171. 13. pr. 28. 2.
149 Cynoforchis militaris mont. spiva rubente conglomerata, E.
    150 Cynoforchis milit. prat. elatior fl. variegatis, B. pr. 28. 3.
  151 Orchis palmata prat. angustifol. major, B. pr. 60. 6. Fig. 152 Orchis palmata angustifolia minor, CB. 85. 6. 153 Orchis palmata angustifolia minor, CB. 85. 6. 153 Orchis palmata angustif. minor edoratissima, B. pr. 30. 7. Fig. 154 Orchis palmata mont. maxima, B. pr. 30. 9. Fig. 154 Orchis abouting folia CB. 86.
    155 Orchis arbortiva fusca, CB. 86. 1
 155 Orents attortiva justa, CB. 60. 1.
156 Gramen atvense fanicula erispa, E. pr. 6. Fig. R.1270.9.
157 Gram. mutorum rad. repente, B. pr. 2. 5.
158 Gr spica eristata subhirsuum, B. pr. 8. 19.
159 Gr. pratense spical slavescente, CB. 3.3.
160 Gr. fol. junceis brevibus maj. rad. nigra, B. pr. 34. xi.
 161 Gr. nemorof. paniculis albis, capillaceo folio, B. pr. 14.45.
162 Feftuea pratentis lanuginofa, B. pr. 19. 68.
163 Feftuea dumetorum angustist. & pilosis foliis, B. pr. 19.70.
164 Gr. hirfutum, angustistol. majus, B. pr. 16. 47. Fig.
165 Gr. Carophyl. foliis, fpicâ divulsa, CB. 3. 6.
 166 Gr. Caroph, latum mont. spica varia, B.pr. 9, 23. Fig. 167 Gr. Caryophylatum nemoros. spica multiplici, B. pr. 9.22.
167 Gr. Caryophylatum nemorol. spica multiplici, B. pl. y. 22.
168 Gr. nemorolum, spicis parvis asperis, CB.7.4.
169 Filix fontana minor, CB. 358. 2.
170 Filix ramosa minor, pinnulis dentatis, CB. 358.4.

BASILIE Plantarum Catalogus, quarum Exemplaria sicca vel Semina recentia, valde grata erant, pari Commutatione
Jac, Petiver.

171 Acetofa arvenf. mm. non lanceolata, 33. CB. 114. 15.

172 Atriplex fylvestris altera, 34. CB. 119. 2.

173 Blitum album minus, 34. CB. 118.2. Ray 200. 4.

174 Blitum rubrum minus, 34. CB. 118.5. R. 200. 2.

175 Campborata glabra, 113. CB. 486. 2. R. 211.

176 Lapathum mm. seu minimum, 33. CB. 115. 5.

177 Lapathum aquaticum minus, 33. CB. 116.

178 Perscaria angustifol. 29. B. pr. 43. R. 183.

179 Plantago angustifol. minor, 56. CB. 189.6. R. 878.

180 Plantago trinervia angustissimo folio. B. pr. 98. 3.
 170 Plantago augustioni minor, 300 CV 180 Plantago trinervia angustissimo solio, B. pr. 98. 3.
181 Chondrilla juncea viscosa arvensis, 39. CB. 130.1.R 223.9.
  182 Cichoreum mont. angustifol. hirsurie asperum, CB. 126. 3.
183 Hieracium Dent. Leonis folio, monoclonon, 37. CE. 127. 8.
184 Id. culmis è rad. pluribus, Cat. 37. var.
185 Hieracium montanum Rapifolium, 38. pr. 64. 1.1. R. 237.xi.
186 Hieracium mont. hirsutum ramos. parvis stor. CB. 128. 9.
187 Hieracium mont. latifol. glabrum majus, CB. R. 233.13.
188 Hieracium muror. anzust. non laciniatum, B. pr. 67.22. Fig.
189 Hieracium protunde sinuatum, pubescens, B. pr. 67.23.
190 Hieracium prat. latifol. non sinuatum majus, CB. 129. 2.
191 Id. minus, 39. CB. 129. 3. Pratis palustribus Michelfeld.
192 Lastuca mont. purpuro cerulea major, CB. 123.3. R. 223.8.
193 Senecio minor, latiore solio I. montana, CB. 131.2.
194 Soncho assinis, Lampsana sylvatica, 36. CB. 125.2.
195 Aster mont. luteus, sol. Salicis glabro, CB. 266.6. R. 266.1.
196 Cacalia sol. crassis & hirsutis, 58. CB. 198.2. R. 299.1.
197 Conyza palust. servatisolia, 79. CB. 266.4. Ger. 347. s.
 183 Hieracium Dent. Leonis folio, monoclonon, 37. CE. 127. 8.
 190 Cacasia ioi. Cianis & minutis, 50. CB. 1902. R. 2991.
197 Conyza palufi. ferratifolia, 79. CB. 266.4. Ger. 347. f.
198 Gnaphalium vulg. majus, 78. CB. 263. 8. 295.4.
199 Gnaphalium medium, CB. 263.4. qr? R. 295.5.
200 Gnaphalium minus repens, CB. 263.6. qr? R. 296.6.
  200 Gnaphatium titilus repens, CB. 203.6. qt. R. 290.6.
201 Virga aurea angustifolia, 79. CB. 268.4.
202 Virga aurea larifol. serrata, 79. CB. 268.2. R. 297.6.
203 Bellis sylv. media, caule carens, 77. CB. 261.6. R. 349.1.
204 Tanacetum minus album, odore Camphora, CB. 132.3.R. 346.3.
205 Tanacetum mont. inadorum, minore flore, 40. CB. 132.1.R.
                 352.12.
```

```
206 Cardous lanceatus angustifolius, 104. B.pr. 155.4.
207 Cardous pratensis latifolius, 102. CB. 376.1. R. 305. 1.
208 Cardous prat. Asphod. rad. fol. profunce & tenuiter lacini-
   208 Carduus prat. Afphod. rad. fol. profunce & tenuiter laciniatis, CB. 377. 4. Ger. 589. 6.
209 Carduus prat. polycephalus, 102. B. pr. 156.5. R. 309 5.
210 Carduus lpinofifimus angulifol. vulg. CB. 305. 3.
211 Carduus spinofif. latifol. lphærocephalus, 102. pr. 385. 1.
212 Carlina acaulos, magno flore 103. CB. 380. R. 288. 1.
213 Carlina caulescens magno flore CB. 380. 2.
214 Cirfum angustifolium, 102. B. pr. 155. 2.
215 Cirfum fingulatibus capit. CB. 377. 6. Dod 37. f. R. 307 9.
216 Cirfum fingulatibus capit. alterum, 102. Monte Wasserfall.
217 Lacea nura angustifol. 81. B. pr. 126.1. R. 326.12
   216 Cirfum fingul. capit. alterum, 102. Monte Wassersall.
217 Jacea nigra angustisol. 81. B. pr. 126.1. R. 326.13,
218 Scabiosa capit. gl.bos major, 80. GB. 270. 2.
219 Scabiosa mont. latis. non laciniata rubra, GB. 270.4.R.379.
220 Succisa hirsuta, 80. GB. 269.2. R. 380.21, var.
221 State major, calyculis non splendentibus, 81. GB. 273 3.
222 Bupleurum angustisol. 81. GB. 278.2.
223 Angelica aquatica, 47. GB. 156.6 R. 434. 3.
224 Apium mont. sol. tenunbus divisis, Cat. 46. var.
225 Capuisolis. 47. GB. 158. 3. Ray 415.4.
    224 Apium mont. 101. tenunus divins, cas. 40. vai.
225 Caruifolia, 47. CB. 158. 3. Ray 415.4.
226 Cicutaria larifol. lœtida, 48. CB. 161.3. R. 451. 5.
227 Daucus Alpin. multifido longoque folio, 45. CB. 150.2.
228 Daucus mont. spii fol. major, CB. 150.5. Ges. 858. f. 3.
    229 Daucus mont. Apri fol. major, CB. 150.5. Get. 856. J. 3.
230 Daucus pratenfis. Apri folio, B. pr. 77. 2. R. 449. 7.
231 Millefolium aq. umbell. capillaceo brevique folio, CB. 141.2.
R. 459.5. Dod 535. f. 1.
232 Sefeli mont. Cicuta fol. subhitsutum, 49. B. pr. 85. 1. R.
  232 Sefeti mont. Cicuta foi. Indontfutum, 49. B. pr. 85. 1. Ri 452. 7.
233 Sefeti prat. tenuifolium, 49. B. pr. 85. 2. R. 453. 3.
234 Sphondylium Ali inum glabrum, 47. B pr. 83.3. Fig. R. 409. 4.
235 Mollugo mont. latifol. ramol., 95. CB. 334. 1. R. 481. 2.
236 Pulmonaria angult. fl. cerul. rubente, 77. CB. 260. 2. R. 489. 3.
237 Brunella folio laciniato, 77. CB. 261. 2. R. 551. 2.
238 Brunella min. alb. laciniata, CB. 260. 1. R. 552. 5.
239 Hedera terrefiris montana, E8. CB. 306. 2 R. 567. 2.
240 Lamium alba linea notatum, f. Milzatella, 66. CB. 231. 5.
R. 560. 4.
241 Fleudoslachys Alpina, 68. CB. 236. 2 R. 555. 5.
  R. 560. 4.

241 Ffeudoslachys Alpina, 68. CB. 236. 2. R. 555. 5.

242 Pieudoslachys Alpina inodora, 68 B. pr. 113. 2.

243 Serpyllum angustisol. hirsut. 63. CB. 220. 8. R. 522. 10.

244 Sideritis vulg. hirsuta, 66. CB. 233.4. R. 563. 4.

245 Urtica fasua fol. non serratis, 66. CB. 232. 3. Lob. Ic. 527.

246 Scrophularia Ruta canina dicta vulg. 68 CB. 236. 8. R. 766.9.

247 Aconium Lycott. albis pallidilye faribus, 54. CB. 183. 3.

248 Millefolium 2q. cornusum majus, 43. B. pr. 73.5. Fig. R. 191.24

249 hillefolium 2q. cornusum minus, CB. 516. defer.

250 Pulsacilla fol. tenunus meisto. fl. minore, B. pr. 94 s.

251 Quinquesolium minus repens luteum, 92. CB. 325.3.

252 Quinquesolium mont. eredt. hirtut. luteum, B. pr. 139. 3.

R. 613. xi.
                                 R. 613. xi.
253 Kanunculus nemorolus luseus, CB. 178. 1. Fufch. 152 f.
254 Ranunculus palustris serratus, 54. CB. 180.2. R. 587. 7.
255 Sedum Alpin. Gramin. sol. lacteo store, 83. CB. 284. 4.
256 Sedum arvense st. rubente, CB. 283. 7. B. pr. 132.9.
257 Bryonia lævis, i. nigra baccitera, 86. pr. 135. R., 661. 3.
258 Polygonatum latif. minus, st. majore, 87. B. pr. 129. R.665.
259 Brassica sylv. ramosa glabra, 33. CB. 112. 7. R. 799.5.
260 Bursa Paltoris media, 31. CB. 108. 3.
261 Dentaria pentaphyllos, 91, CB. 322. 1. Ray 785, 3.
262 Dentaria pentaphyllos, CB. 322. 2. R. 784. 2.
263 Eruca cærulea in arenosis creicens, 28. B. pr. 45. Fig. R. 793. 4.
264 Myagrum sætidum, 42. B. pr. 51. I. R. 820. 3.
265 Nasturium aq. crect. fol. longiore, 30. CB. 104. 3.
266 Nasturium monolpermon, 27. B. pr. 37. 2. Fig.
                               Ranunculus nemorolus luteus, CB. 178. 1. Fusch. 152 f.
     267 Rapistrum monotpermon, 27. B. pr. 37. 2. Fig. 268 Thisfpi Alpin. majus, capit. rotundis, CB. 106.6. R. 822.6.
        269 Thlaspi Alpin. minus, cap. rotundo, 31. B. pr. 48. 6. Fig
                                  R. 821. 5.
  Thlaspi mont, incan, lut. Serpylli, fol. majus, CB. 107. 12.
271 Campanula Alpin. Linisolia cerulea, B. pr. 34.4. R. 740.16.
272 Rapunculus nemoi. angustif. migno st. major, CB. 93. xi.
273 Rapunculus nem. angustif. parvo store, 26. CB. 93. 13.
274 Rapunculus spicasus albus, 25. CB. 92. 3. R. 743.1.
275 Gentiana autumnalis ramosa, 55. CB. 188.2. R. 720.8.
276 Gentiana autumnal. angustifolia, 56. CB. 186.6.
277 Gentiana cruciata, 55. CB. 188-1. R. 717.5.
278 Gentiana major lutea, 55. CB. 187.1. R. 716.1.
279 Linum tylv. angustifol. st. minore, 62. CB. 214.3. R. 1073.9.
280 Millesolium, 2q. Equisetisolium caule nudo, CB. 141.5.
281 Alsne major, 73. CB. 259. 10.
282 Alsne major, 73. CB. 259. 10.
        270 Thlafpi mont. incan. lut. Serpylli, fol. majus, CB. 107. 12
         * 282 Alfine major, 73. CB, 259, 10.
                                                                                                                                                                                                                                                                                                                                                            28;
```

283 Cayophyllus, fylv. vulg. 60. CB. 209. I. R. 950. I.
204 Caryophyllus fl. tenuissine distecto, 61. CB. 209. I.
205 Caryophyllus arvent. bis fut. st. majore, CB. 210. I. G. 477. f. xi.
206 Caryophyllus arv umbellatus, fol. glabro, CB. 210. I. G. 477. f. xi.
207 Caryophyllus sylv. fl. minimo, 61. CB. 211. 5. K. 1033. I. 3.
208 Geranium Cicura tol. maj. inodorum 90 CB.
208 Geranium fol. rot. minus. profunde lacuniato. Cat. 90. Geranium Cicuta tol, maj. inodorum 90 CB.

280 Geranium fol, rot. minus, profundè laciniato, Cat 90.

290 Verhascum fam. sl. luteo magno, 69. CB. 237.3. R.1094.2.

201 Nigella arvensis cornuta, 44. CB. 145.1. R. 1070.3.

202 Lotus pentaphyllos corniculata, 94. CB. 332.1. R.967.5.

203 Lotus pentaphyllos fruticosus, 94. CB. 330. 2. R. 968.7.

204 Trisolium mont. pur pureum, 93. CB. 328. 1. R. 944.6.

205 Trisolium capit. spumoso aspero majus, 93. B. pr. 140.8.

206 Trisolium capit. spumoso aspero minus, B. pr. 140.8. R.

207 Latbyrus major sylv. alter. 96. CB. 344.4.

208 Latbyrus sylv. minor, 96. CB. 344.5. Dod. 529. f. 1.

209 Orobus sylvat. purp. vernus, 97. CB. 351. 2. Dod 542 f. 2.

300 Securidica dumet. maj. sl. vario, sil. articulatis, CE.349.4.

301 Securidica dumet. min. pallidè cerulea, 97. CB. 349. 5. 300 Securidica dumet. maj. fl. vario, fil. articulatis, CE. 349. 4.
301 Securidica dumet. min. pallide cerulea, 97. CB. 349. 5.
302 Vicia sepium fol. rotundiore acuto, CB. 345. 4. Dod. 531. f. 1.
303 Vicia sylv spicata, 97. CB. 345. 8. R. 903. 19.
304 Vicia vulg. acutiore folio, 96. CB. 345. 5. Matth. 416. f. 2.
305 Fumaria bulbolarad. cava major, 44. CB. 143. 1. R. 975 4.
306 Fumaria bulbolarad. non cava major, CB. 144. 2. R. 975 5.
307 Fumaria bulborad. non cava minor, CB. 143. 3. Dod. 347. f. 2.
308 Allium sparieeo capite, fol. latiore, 21. CB. 74.6.
309 Allium splv. latif. capitat. purpuralcens, 21. CB. 74. 10.
310 Allium sylv. capitat. purp. majus, 21. CB. 74. 12.
311 Allium sylv. capitat. purp. minus, 21. CB. 74. 12. R. 1119. 2.
312 Allium mont. bicorne, angustifol. fl. purp. 21. CB. 74. 22.
31. R. 1119. 2. R. 1119. 2.
313 Meudo-Afphodelus Alpinus, 18. CB. 29. 10. R. 1194. 1.
314 Iris vulg. German. f. lylvettris, 19. CB. 30. 1. R. 1180. 1.
315 Iris prat. angustif. humilior. 19. CB. 32. 3. Dod. 246. f.
316 Cynosocrabis latifol. hiante encullo major, 21. CB. 80.1. 317 Cynosorchis angustifol. hiante cuculto, 22. B. pr. 28. 1. R. 1216.16. R. 1216.16.

318 Cynoforchis milicaris major, 22. CB. 1.1. R. 1213.13.

319 Cynoforchis milicaris minor, 22. CB. 81. 3. Ger. 165. f.12.

320 Orchis fol. feffilibus non maculatis, 22. CB. 82. 3.

321 Orchis odore Hirci long. angustioreque fol. 23. CB. 82. 2.

322 Triorchis lutea fol. glabro, 23. CB. 84. 3. R. 1217. 24.

323 Triorchis lutea altera, 23. CB. 84. 4. Ger. 167. f. 1.

324 Meudorchis, 24. CB. 84. 8. R. 1226. 1. Cam. H. 111. f.35.

325 Orchis pilmata mont. maxima, 24.B.pr. 30.9 Fig. R. 1224.4.

326 Orchis abortiva susca 24. CB. 86. 1. Dod. 553 f. 2.

327 Ophris trifolia, 25. CB. 87.2. Ger. 326. f. 2.

328 Ophris trifolia, minor Cat. 25.

329 Helleborine angustif. palustr. s. prat. 55. CB. 187. 3. R. 1230. 2.

330 Helleborine fi. alb. 55. CB. 187.5. K. 1231 6.

331 Helleborine mont angustif. purpurateens, 55. CB. 187.7.

332 Nymphea alba minor, 57. CB. 193. 2. G. 663. f. 3.

333 Thalichum prat. angustissimo folio, 95. B. pr. 146. Fig. R. 404.9. R. 404.9.
334 Tithymalus fylv. limato flore, 84. CB. 290.7. R. 871.41.
335-Tithymalus Linaria fol. 84. CB. 299.2. R. 866.15.
336 Tithymalus Cypariflias, 85. CB. 291.2. R. 866.15.
337 Tithymalus Cypar. fol punctis croceis notatis, CB. 291 1.
338 Tithymalus Paluftris fruticolus, 85. CB 292.2. R. 864.8.
339 Tithymalus mont. non acris, 85. CB. 292.3. R. 870.34.
340 Valeriana fylv. maj. montana, 50. CB. 164. 5.
341 Valeriana Paluft. med parum laciniata, 50. B. pr. 86.2.
342 Valeriana Alpina prima, 50. B. pr. 86.3. R. 390. 9.
343 Valeriana Alpina prima, 50. B. pr. 86.3. R. 390. 9.
344 Gramen Caninum vineale, 3. B. pr. 1.1. 343 Valeriana Alp. Serophularia, folio 50. B. pr. 87.4.1.390.10.
344 Gramen Caninum vineale, 3. B. pr. 1.1.
345 Gr. Canin. supinum minus, 3. CB. 1.7. R.1256.9.
346 Gr. Tremulum minus, paniculâ parvâ, 4 B. pr. 4.9.
347 Gr. prat. panicula & glumis longioribus, Cat. 4.
348 Gr. mont. panic. spadicea delicatiore majus, 5. B. pr. 6.12.
349 Id. minus, Cat. 5. B. pr. 6.12. Monte Wasserfall.
350 Gr. palust. paniculatum altissimum, 5. CB. 3.1.
351 Gr. palustre paniculatum altius, Cat. 5.
352 Gr. palustre paniculatum altius, Cat. 5.
353 Gr. spicatum angust. montanum, 6. B. pr. 8. 20.
354 Gr. tylvat, augustifol. spica alba, 6. B. pr. 8. 21.
355 Gr. spicatum, fol. & spicis hirsut. mollibus, 6. B. pr. 9.24.
356 Gr. Typhoides max. spica longissimà, 6. B. pr. 10. 25. Fig.
358 Gr. Typhoides molle, 7. CB. 4. 2. Ger. 9. f. 2.
359 Gr. Typhoides asperum primum, 7. CB. 4. 3. G. 10. f. s. 350 Gr. Typhoides molle, 7. CB. 4. 2. Gr. 9. 1. 2.
359 Gr. Typhoides afperum primum, 7. CB. 4. 3. G. 10. f. 1.
360 Gr. Typhoides afperum alterum, 7. CB. 4. 4. G. 10. f. 2.
361 Gr. Typhoides, fpica angustiore & longiore, 7. CB. 4.5.
362 Gr. Typhoides, culmo reclinato 7. CB. 4.6. G. 9. f. 3.
363 Gr. Typhoides, spica multiplici. 7. CB. 4.7. R. 1298.13. 363 Gr. Typhoides, spica multiplici, 7. CB. 4.7. R. 1298.13.

364 Gr. Sparteum Basil, capillaceo solio 7. B. pr. xi. 29. 365 Gr. Junceum polystachion, 8. CB. 5.7. R. 1309. 16. 366 Gr. Junceum sol. & spica Junci minus, 8. CB. 6. xi. 367 (r. Arundinaceum spica multiplici tenuiore, Car. 10. 368 Gr. Arundin, enode minu, fylvaticum, 10. CB. 7. 7.
369 Gr. mont, pame. Miliaceâ fparsă, 12. B. pr. 17.56.
370 Gr. Sorghi paniculă erectum, 12. CB. 8.1. R. 1283.1.
371 Gr. latif. fpic: Tricica compactă, 12. B. pr. 18. 53. Fig. 372 Id. folio angustiore, Cat. 12.
373 Gr. angustis spica Triticea compasta, 12. B.pr. 18.56. Fig. 374 Gr. spica Brizæ majus, 12. B. pr. 18. 58.Fig. R. 1257.1. 375 Gi- Hordeaceum mont. f. majus, 12. CB. 9.1. 376 Gr. Loliac. latif ipica anguit. 13. B.pr. 19.60. R. 1263.2. 377 Idem fol. longioritus, Car. 13. 378 Festuca Granunea, glumis vacuis, 13. B pr. 19.64. R. 1290 9. 379 Festuca dumetorum, 13. B. pr. 19.59. R. 1262.2. 380 Gr. nemorol. spica ruselcente molli. 10. CB. 7.3. 381 Gr. hissut, satisol, minus, juncea panicula, Cat. 10. 382 Id. majus, Cat. 10. 382 Cyperoides angust. spica spad, viride minus, B. pr. 19.41.
384 Cyperoides angust. spicis longis erestis, 9. CB. 6.4.
385 Cyperns longus inodorus sylvaticus, 15. B. pr. 24.
386 Juncus acumine restexo major, 14. CB. 12.1. R.1304.xi. 387 Secale luxurians, 17. CB. 23. 4. 388 Secale vernum, vel minus, 17. CP. 23. 3, 389 Zea Amylea, vel Zeopyrum Amylaum, 17. CB. 22. 3. 390 Zea Briza dicta, vel monococcos Germanica, CB. 21. 1. 391 Zea dicoccos, vel majoi, 17. CB. 22.2. Ger. 62. f. 392 Equiseum paluttre Linaria scoparia fol. B. pr.24 1. 393 Equiserum palustre minus polystachion, 15. B.pr. 24.2. 394 Equisetum palust. tenuist. & longist. fetis, B. pr. 24 3. 395 Equifetum nudum minus variegat. Basiliense, B. pr. 24.4, 396 Filix aculeata major, 99. CB. 358. I. pr. 151. 2. 397 Filix aculeata minor, CB. 358. 2. pr. 151. 3. 398 Filix fontana major, s. Adianth. alb. Filicis folio. CB. 258. 1. 399 Filix non ramosa pinnul, tenuiss. & tenuissime dentatis, Cb. 358 4.

* 399 Filix Querna 99. CB 358.5, Tab. 1c. 793.2. Ger. 980.f. 1.

400 Muscus vulgatissimus, 100 CB. 360. 1. Dod. 472.f. 1.

401 Muscus vulg. pennatus major, 100. B. pr. 151. 1.

402 Muscus vulg. pennatus minor 100. B. pr. 151. 1. var. 402 Muscus vulg. pennatus minor, 100. B. pr. 151. 1. var. 403 Muscus ramolus repens, spicatus, 100. B. pr. 151. 4. 404 Muscus ramos. repens, apicibus candicautibus, B.pr. 151.4.var. 405 Muleus ramofus major, 100. B. pr. 155. 5. R. 123. 2. 406 Muscus ramos. maj. elegentior & viridior, 100. B. pr. 151. 5. vai. 407 Muscus ramol erect. minor, 100. B. pr. 151. 6. R. 123. 3. 408 Muscus saxatilis Ericoides, 100. B. pr. 151. 7. R. 123. 4. 429 Muscus polytrichoides, 1. fol. capil. rigid. B. pr. 152. 10. 410 Id. alter minor, B. pr. 152. 10. var. 411 Id. 3. minimus & mollis, B. pr. 152. 10. var. 412 Mufeus parvus stellaris, 100. CF. 361. 13. 413 Museus terreturis Coralloides incanus, 101. B. pr. 152.xi. 414 Museus terrett. Coralloides erectus cornib: tubescentibus, B. pr. 152. 11.
Muscus aquaticus, folio expanso, 101. B. pr. 152. 14. 116 Pinus fylvessis, 113. CB. 491. 2. R. 1399. 3.
417 Pinus fylvessis, 113. CB. 491. 2. R. 1399. 3.
417 Pinaster tenustoius, julo purpurascente, CB. 492.10.
418 Salia vulgaris rubens 111. CB. 473. 3. R. 1421. 8.
419 Salia sativa lutea, fol crenato, 111. CB. 473.4. R. 1421.7. 42) Saliw fol. Emygdalino, utrinque virente aurito, CB. 473. 5. 421 Saliw fragilis, 111. CB. 474. 9. Park. 1431. 4. 422 Saliw humilis, capitulo iquamofo, 111. CB. 474. xi. 423 Saliw oblongo incano folio 111. B. pr. 159. 2. R. 1423.2. 424 Salix humilis latifolia eretta 111. B. pr. 159. 2. R. 1423. 5. 425 Buxns folis rotundoribus 111. CB. 471. 2. R. 1694. 2. 426 Alnus folio incano, 107. CB. 428.3. R. 1410.2. 426 Alnus folio incano, 107. CB. 428.3. R. 1410.2.
427 Alnus nigra polycarpos, 107. CB. 428.3.
428 Alni effigie, lanato folio minor, 109. CB. 452.2. R. 1461.c.9.
429 Tilia montana, maximo folio 106 CB. 426.2. R. 1695. 3.
430 Ulmus campeltris, 106. CB. 426. 1. Dod. 837. f.
431 Ulmus montana. 106. CB. 426. 2. Ger. 197. f. 1.
432 Genista spinosa minor Germanica, 104. CB. 395. 3.
433 Chama-genista fagittalis, 105. CB. 395. 2. R. 1725. 5.
434 Ceraso assinis, 108. CB. 451. 15. R. 1540. c. 15.
435 Rosa sylvestris Fomisera major, 112. CB. 484. 15.
436 Rosa arvensis candida, 113. CB. 484. 17. qr? R. 1471. 14.
437 Sambucus racemosa iubra, 110. CB. 456. 3. R. 1610. 4.
438 Vitis Idea, fol. carnosis, 110. CB. 470. 6. R. 1489. 5.

Plantæ Basilia locis Propriis, viz.

Circa Urbem pratis, 134. 6. aridis * 217. 21. 83. humidis * 207.* 23. * 38. collibus * 276. Agris 286. * 7. 436. marginibus 210. 92. 336. * 7. Viis 143. 79. 80. Sepibus 244, 5. 85. * 9. 93. 302. 3. 4. 6. Fimetis 226. & 1. Ruderibus * 240. Mæniis 140.

Muria Muris 221. * 410. 1. Fossis * 419. Alnetis 435. Dumetis 203. 11. 8. * 34. 300. 1. 5. Sylvis cæduis * 340. Frumentaceis 198. 0. 200. Segetibus 291. Graminibus 141. Montibus 430. Binninghen * 417. Birsa arenonis * 383. 421. dumetis * 167. * 379. * 406. herlidis 316. paludosis 384. ad pontem * 263. Birsec arcis colle * 322. Benehen agris uliginosis * 361. Brilingen viis 143. St. Christiana monte 438. Cliben * 377. dumet. 165. 207. 312. 74. Pratis * 193. Crentzachi monte 140. 229. 52. * 72. 94. 9. 301. * 13. 26. * 50. * 69. 74. * 81. 2. * 403. 4. 7. 25. Saxosis * 188. 258. 77. 348. Umbtos & humidis 190. 366. Dornach 270. Fliehen ad Balneum 434. Hiltelingen * 248. Huningen sylvis 168. * 94. 257. 75. 7. 335. 426. sylvis humidis * 366 Dornach 270 Fliehen ad Balneum 434 Hiltelingen * 248. Huningen sylvis 168. * 94 257 75 7. 335. 426 sylvis humidis * 354 agris 145 81. 256 67. Lagenbruch sylvis * 196. Landskron 428. arce 327. montons 437. Michelfeld 350. 81. aquis * 231. 49. * 332. * 423. paludibus 197. * 364. * 86. * 92. * 3.4. udis * 178. 352. 80. * 424. dumetis * 189. 274. Salicetis * 295 * 355 7. * 423. pratis 150. 1. 3. 60. 8. * 280. * 9. * 24. * 30. * 3. * 50. 4. * 333. 50. 81. * 93. 4 paludosis pratis * 164. * 91. 220. 84. * 315. 29. 38. * 41. Monchenstein 156. 306. 81. pratis * 190. muris 314. arenosis ad pontem * 225. sylvis mont. 330. Muteto monte * 144. 54. * 64. * 82. * 6. 92. 5. 201. 2. 5. 22. * 835. * 9. 61. 2. 99. 318. * 9. 319. 82. * 427. * 9. 32. herbossis 412. dumet. humidis * 247. * 367. sylvis 274. pratis 152. * 275. pratis mont. 149. * 317. saxosis 166. 274. pratis 152. • 275. pratis mont. 149. • 317. saxofis 166. • 88. • 258. ad Novum Domum sylvula 306. rivulis. • 415. Otlingen agror. semitis • 486. Reichenstein monte 146. 428. Otlingen agror. semitis * 486. Reichenstein monte 146. 428. Rheni ripis 181. 246. arenosis * 422. saxosis 309. dumetis 259. beh der Baar * 296. 321. 955. Riehen seget. * 389. 90. 1. dumetis 307. sepibus * 167. Wiese ripis 156. 310. I saxis * 408. arenosis 131. 204. 64. 90. 414. 22. 23. ad pontem * 238. pratis * 347. 420. agris 308. dumetis 347. 99. * 413, nemora ad pontem * 206. Alneto 253. * 82. 305. Wilensi sylva 368. 409. Vineis 266. uliginosis * 401. 2. Wassersal monte 135. * 44. 7. * 85. 7. * 92. * 6. 212. * 3. * 4. * 5. * 6. * 27. * 32. * 34. * 41. 2. * 3. * 68. 9. 71. * 323. 4. * 43. * 9. 50. 68. * 75. * 96. 7. * 9. * saxosis 331. * 42. N. B. Ha* in duplo pergratæ erunt, Jac. Petiver. Fac. Petiver.

COM. III. Plants gathered on Mount Serato, lately fent me from Mr. John Salvadore at Barcelona.

439 Hieracium Pyrenaicum Dent, Leonis fol. latissimo Schol.

Bot. 189. Instit. Rei Herbar. 472. Ray Vol. 3. p. 139.14.
440 Elychrysum sylvestre angustifolium, capitulis conglobatis, CB. 264.9 Phytopin. 5. 4.8. Lob. Ic 486. 1. Ad. 203.
f. Belg. 570. f. Ray H. Plant. 283.xi.
441 Globularia montana humillima repens. El. Ros. car. 7-0...

441 Globularia montana humillima repens El. Bot. 571. Instit. 467. Scabiosa X sive repens, Clus. v. 1, 4 f. Pan. 540 f. R. 81.27.

442 Jacea nigra angustifolia, vel Lithospermi arvensis soliis, caule aspero, CB. 271 2 Prodr. 127 1. varietas R 326. 3 var.

443 Laserpitium angustissimo & oblongo folio, El. B. 274. Inst. 324. Laserpitio Lobel. similis, angustifol. I. B. Vol. 3. I. 27. p. 237. f. z. Chabr. 399. Ic. 5.
444 Gallium saxat. minim. supinum & pumilum, E. B. 94. Inst.

445 Polium mont. luteum, CB. 220. 2. Phyt. 417. 2. Tab. Ic. 364. Hist. 2. 80. f. 1. Polium luteum, Lob. 1c. 487. 1. Ad.

207. Belg. 572. f. R. 525. 8.

446 Euphrasia pumila saxatilis, oblongo, angusto, serrato & villoso folio, st. purpurascente Salvadore.

447 Ranunculus Gramineo folio, bulbosus, C.B. 181. 7. pt.

27. 30. Ray 588. 2

448 Veronica Hilpanica fol. tenuissime laciniatis, fl. ceruleo Salv. 449 Convolvulus argenteus umbellatus supinus, EB. 73. Inft. 84. Campanula Lychnidea, I. B. 2. 1. 20. p. 803. f. opt. Chabr.

264. f. 3. Ray 735. 25.
450 Aphyllanthes Monspeliensium, Adv. 190. f. Lob. Ic. 454.2.
B. 5 38. f. Caryophyllus cesul. Monspel. CB. 209. 8. pt. 392. 13. R. 990.1**3.**

431 Heliant bemum fol. Rorifmar. Splendent. Subtus incanis, EB. 216. Ledon 8. Cluf. 80. f. 2. Hifp. 165. f. Ray 1010.

452 Helianthemum fol, Rorifmarini utrinque incano Salvad. 452 Fictianto emum 101, Rorifmarini utrinque incano Salvau.
453 Geranium Pyrenaicum, fol. cineraceis, rad. crafsâ, magno
flore variegato, El. Bot. 233. Ray Vol. 3. p. 515.46.
454 Coronilla minima, EB. 510. Inft. 650. Ray 925. 1.

Lotus enneaphyllos Dalech. Lugd. 510. f. CB. 349. 4.

COM. IV. Planta Ilvenses. Plants sent me from the Reverend
Revero Toxic subject has gathered in the Island Film on Use.

Bruno Tozzi, which he gathered in the Island Elba or Ilva, lying in the Tyrrhene Sea, against Tufcany.

455 Clinopodium Creticum Alp. Exot. 264. f. Ray 554. 4. Calamintha Cret. angusto oblongo folio, Inst. 194.

456 Linaria Ilvensis faxat. st. pallido ridu purpureo Tozzi.

457 Viola Ilvensis montana, fl. albo. 458 Viola Ilvensis mont. fl. ceruleo.

450 Viola Ilvensis mont, st. luteo, 460 Tithymalus Ilvensis fruticosus perennis.

461 Hemionitis altera Dalech. Lugd. 1218. f. 2. 462 Polypodium Ilvense Lugd. 1230. f.2. Chabr. 558.4. R. 137.2. Filicula ramosa Lust. pinnutis ad Ceterach accedentibus El. Bot. 432. Initit. 542. Muf. noftr. 212. Ray Vol. 3. p.

84. & 94. 6.
463 Erinacea Ilvensis st. luteo Tozzi, qr? Pl. Etrur. Cat. 21.
Plantæ Ilvensis desideratæ Jac. Petiver,
464 Genista Ilvensis, Lugd. 174. f.
Vide aliæ desideratæ in Etruriæ Plant. rarior. Catalog. viz.

No. 21. 31. 54. 116. 21. 82. 284. 529, &c.
COM. V. Egyptian Plants sent me from Peter Anthony Michael

Botanist to his Highness the Grand Duke of Tuscany.

465 Parietaria Egyptiaca fol. subrotundo serrato, superne glauco scabro, infernè incano & tomentoso Mich. qu? Gaz, Nat. Tab. 7. fig. 2.

466 Filago Ægypt. pumila spicata, Mich. 467 Cotonaria Ægypt. Chamadryos folio tomentoso Nob. Planta Agypt. incana & tomentola, Uva crifpa folio, frudu peltato Mich.

468 Jacea Egypt. spinola Mich.
469 Polium Egypt. toment. erectum, fol. vermiculatis Mich.

470 Serpylli facie, planta Egyptiaca Mich.
471 Melilotus Egypt. siliquis secundum cauliculorum longitudinem ex foliorum alis exeuntibus Mich. Lotus Agypt fol. eleganter crenato D. Lippia D. Stoschio.

472 Hypericum Ægypt. gummifer elegantissim. fol. subrotuado bullato Mich. Myrto-cissus species Pennei, Chif. 68.
473 Tribulus terrestris Ægyptiacus.
474 Tithymalus Ægypt. exiguus, fol. superioribus crenatis Mich.
475 Kali Ægypt. villosim, ralyce stellato aculeato. D. Vallante.

N. B. Vide plura in Iconibus nostris Plantar. Ægyptiacarum

COM. VI. The following Curiofities were prelented me by my hearty Friend Mr. John Burnet, Surgeon to the English Factory at Porto Bell.

476 An Abortive Negro near full grown.

477 Piscis triangularis Amer. fronte ventroque bicornutis. The male Cunnie fish from the West Indies. Its. Nose and Back Cloud-streakt with Black, somewhat like Tortoisesbell.
478 Piscis triangularis Amer. minor, stellis albis. This is said

to be the Female to the Last. Its Belly wholly compos'd of small Diamond Angles.

Buccinum rubescens, ventre fasciato St. Thoma.

The Inhabitants eat the sower Pulp of this, to quench Thirst

481 Colocynthis Guineensis Pomiformis, sem. parvis. 482 Taggee Guin. radix forte Trass generis Gazoph. Nat. Tab. 17. fig. 10. A fort of Earth Nuts which the Negroes eat plentifully of. I should be glad to see its Leaves and

483 An Oftrich Egg from Buen Agres.

484 Radix Gentiana facie à Buen Ayres. A Specific Emetic in a!l Imposthumes.

485 Radix Tintoria, Rubie forte species. (Raiz roug Hispanis)
With this Root they dye a deep Red and probably may be

an Ingredient in the following. 486 Glastrom rubrum è Buen Ayres. A Composition like Storax but red; with this they dye a very sine bright Scarlet, as may be seen by some Wool, this Curious Gentleman hath lately brought from thence to the Direstors of the South Sea Company, from whom we may now expect greater

Discoveries. COM. VII. Miscellany Communications, from divers Hands. 487 Corallium Balearicum globosum, tubulis brevibus stellaris,

487 Coramum Batearicum giodotum, invitis orevidus itelians, Broke from a Rock in Port Mahon Harbour, by our Curious Friend Mr. James Campbell, Surgeon to that Port, &c. 488 Cambodia Birds-nefts, Hist. of Drugs, 21. p. 265. Nidus gelatinus Cambodiensis Nobis, Dale Suppl. 21. p. 265. Bont. 66. fig. Pomet. pt. 2. l. 1. c. 21. p. 72. Ray Ornith. 7. p. 151. id. Angl. 7. p. 215. Syn. Avium. 6. p. 72. Two of these intire Nests mostly cover'd with Feathers, were generally presented me by Mr. 20 ha Courtes Surgeon. roufly presented me by Mr. John Courtney Surgeon.

489 Brasilian Frog-fish. Rana Piscatrix minor Grew Mus. Reg. Soc. 10). desc. Water Batt, Wafer xi. p. 196. Guacucuia Brasil. Mategr. 143. fig. Ray Icth. 89. Tab. E. 2. f. 3. This strange Fish Dr. Crawford sent me from Jamaica.

490 Stella marina Indica reticulata Lobel in Pharmacop. Rond. 139, fig. opt & defer. This elegant Sea Star. Mr. John Douglass Surgeon, sent me from Antegoa.

PETIVERIANA III, seu NATURÆ

COLLECTANEA;

Domi Forisque Auctori Communicata.

A JACOBO PETIVER, Regia Societatis Socio, &c. Londini.

COMMU- P. L. A. N. T. E. Baleares. Plants found in the NIC. I. Islands Majorca and Minorca, by the Curious Mr. John Salvadore, Apothecary at Barcelona, and Mr. George Bouchere, Surgeon at Port Mahone.

1 Plantago fol. lato Coronopi Nobis. Bouchere.

2 Plantago minima angustifol. villola. alvadore.

3 Perexil Balear. fol. minore Beta. Bouch.

4 Urtica Balear. Pilulifera, fol. alte incuis. Bouch.

5 Chrondrilla Bal. spinosa, Cichor. fol. st. luteo. Salvad.

9 g? Sonchus frut. petræus African. spinosus, Park. &o4. st. 7.

Ray 224. f. 10.

6 Cichoreum caulibus florum Junceis, Nob.

6 Creboreum catilious porum junceis, Nob.
7 Conyza luteus radice odorata.

After luteus rad. odora, CB 269.3 Ray 264.13

Afteris species, an Baccharis, Colum.253. st. opt. c.97.
8 Conyza minima Dod. foliis crispis.
9 Elicbrysum Balea. Gnaphaloides saxatile, flor. in strictiorem umbellam congestis, Salvad.

to Gnaphalium Roseum sylv. polyanthos. This is as much branched as the Gnaphal. Roseum Hort. B. pr. 122, f.

11 Bellis sylv. minor Balear. ramosa & villosa, Salv.
12 Bellis maritima, mm. s. minima Roris Solis fonis Cirnea
Boco. Mus. 149. Tab. 107. Ray Hist. Plant. Vol. 3. p. 228.

38. Calamintha Balear, frutesc. fol. minoribus villosis

Marum Syriacum vulgo, Nobis.

Marum Coreufi I. B.3. 1.28, p.243, fig. Chabr. 418.f.5.

Marum Matth. B.537, f. 1, Cam. Epit. 493, f.

Out first Knowledge of this Noble and Gelebrated Cephalick, I find, is owing to that Judicious Padua Botanift Jacobus Antonius Cortufus, who Communicated it to both
Matthiolus and John Baubine, which they justly own.
Its growing wild to near as Port Mahone I could fearce believe, had I not received Specimens of it from Mr. Salvadore and Mr. James Campbell, who were both there.
Marying Balear marit, fewerfears A allow

Marum Balear, marit. frutescens, fl. albo. Poly-Mountain Lecaan, p.74.30.

Polium mont. 5. purpureo flore, Cluf. 362, f. Hisp. 243, f. Polium candidum tenellum tomentofum, fl. purpureo, 1.B 3.

13.1. 28. p. 300. f. Chabr. 431. f. 4 Ray 524. 4.
17 Serpyllum Balear. faxat. fol. 10t. non crenaro, Salv.
18 Sideritis Hifpan, frutescens seu lignosior, E. B. 160. Inst.

Teucrium, CB.247.1 phyt.471.1 Ray 526.1.

Teucrium multis, I.B.3. 1.28. p.29), f. Chabr.429, f.1.

20 Cymbalaria parva hirluta Nob.

Linaria Hederaceo folio villoso s. Cymbalaria Alpina, E.B.

134. Inst. 165.
21 Digitalis Hispan. purpurea minor. E.B. 134. Inst. 165.
Digitalis Verbasci folio Boc. Mus. Tab. 85. s. Ray 3. p. 396

1 & 97. 5 Digitalis angusto Verbasci solio montana Boc, Mus. p 108. Corruda altera, Clus. 178. f. 1. Hisp. 459. f. Ray 684. 4. Asparagus aculeatus alter 3 vel 4 spinis eundem exort, CB.

490. 2

23 Corruda 3 Clul. 178. f. 2. Hifp. 461. f. Ray 684. 5
Afparagus aculeasus. spinis horridus, CB. 480. 3

24 Corruda Hifp. spinis filiquar. facie, Nob.
Asparagus Hisp. aculeis crassioribus horridus, E.B. 250. Inst.

25 anonis Balear, perennis glutinosa villesa, folio minore serrato, flore luteo Salv.

Anonis Ulissiponensis lævis, calyce paleaceo, Nob.

Melilotus lutea minor, foribus & siliculis minoribus spicatim & dense dispositis, Ox. 161. 5 Tab. 16. ser. 2, f.

Melilotus nova Berardi feu crecta, folliculis rotundis minor, Ray Hift. Plant. p.951. pl.6.

28 Genista Balearica faxantis, pallide virens fore luteo, Salvadore.

29 Tragacantha Bil. fol, incanis, fl. albo firiato, Salv.
30 Tragacantha Balear, rame fillima, fpinis brevibus, Nob.
31 Anthillis Balearica Serpilli folio.

32 Alfine Balear, longius radicata, tenuissimo folio, fl. purp. Sulv.

33 Lychnis Balearica Saponariæ folio molli, Nob.

34 Lychnis Balear. viscosa saxat. tereti crispo folio, Salv. 35 Helianthemum Balear, Satureia fol. angustioni, flore luteo,

36 Myrto Cistus Pennei, Chis, 68. fig. Pan. 67. Ger. 1098. f. 16. Fonst. 1279. f. 16. Park. 665. f. Ry 1011. 13 Bot. Hort, 4.

37 Vinca latifolia, fl. parvo cæruleo, Salv. 38 Kali Crassule mnoris folio, CB. 289. Phyt. 566. 4. Kali Floridum repens Neapolitanum, Colum. pt. 2. p. 73. f-

opi. c. 35. Ray 213. 4.
39 Limonium Balear. Bellidis min. folio, fruticosum, Salv.
40 Limonium minus, Bellidis folio, flagellis Fæniculaceis, Bocc.
Med. 143. Tab.103. Ray Vol 3. p.246. pl.6.
41 Limonium min. flagel. tortuolis, Boc. M. 143. T.103. R.3.

p.245. 42 Limonium minimum, Park.1234.5 R. 396. inter. pl.8 & 9. Limonium maritimum minimum, CB.192.6 pr. 99.2 Bocc, 25.f.3. I.B.3.app.861.

25.7.3. 1.B.3. app. 601.
43 Limonium Lufit. Auriculi Usfi fol. E.B. 284. Inft. 342.
44 Limonium Hifp. multifido folio, E.B. 285. Inft 342.

Limonium Tamarifei folio nonnullis, R.3. p. 247. Pluk. 28.

45 Alaternus Balear. fol. serratis subrus ferrugineis.

46 Mattick tree, Lecaan 68.17 fig.Ray 1579. c. 4.
Lentifcus, Dod.871. fig. Cluf.14 fig. opt. Hifp 85. fig.
Lentifcus vulg.CB. 399.1 Lob. 1c 2. p. 96.B.2. p.113. f.Obs.

47 Lentiscus angustif. Massiliensis, Hort. R. Paris Bot. Monfr.

153.
48 Leutiscus Balear. fol. obtusis, maculis candidis.
49 Sanamunda Balear. 2 Clusii, tomentosior.
50 Stonecrop Sanamund. Lecam 75.34.

50 Stonecrop Sanamund. Lecam 75.34.
Sanamunda 3 Cluf. 89. f. Hisp. 178. f. Jonst. 1596.f. 3. Park.
202. fig. 7. Ray 1589.6.
Sejamoides minus Dalech. Lugd. 1670. f. 1. Park. 203.8.
Thymelaa tomentosa, folio Sedi minoris, CB. 462.7.
N.B. There are divers others, both Herbs and Trees in Mr.
Boucher's Collections, which I here omit, until I receive what the Curious Mr. Campbell a Resident there, has been pleas'd to promise me by the first, and all Oppertunities, which I shall publish as they come to Hand.
COM. II. Lustanie Frumenta, Gramina, Cyperi, junci, &c.
Rariores & Desideratæ, Jac. Petiver.

SECT. 1. Frumenta & Gramina Spicata.
51 Hordeum Nudum. Cevada Santa, Lusit.
52 Gramen Hordeaceum vulgare.

52 Gramen Hordeaceum vulgare. 53 Gramen Hordeaceum marinum pulchrum, ND. 54 Gramen Hordeaceum Lustranum,

55 Grainen Ægylops longius spicata Lusit. ND. 56 Gramen Ægylops prælongis ariftis vallata, ND.
57 Alopecuroides ramofa, ND.
58 Gramen slopecurus lanuginolus.

59 Gramen Alopecuroides aquaticum Toursnense, ND. 60 Gramen Alopecuroides campestre vernum Luste. ND. 61 Gramen Alopecuroides hidauum palustre Luste, ND.

62 Gramen Alopec. palustre bulbofum, ND.

PETIVERIANA III, seu

10 63 Gramen Alopec, Segetum elegans, ND. 64 Gr. Canarium pumilum sopiuum Luste. ND. 65 Cr. Canarium ipica ampla Lusit. ND. 66 Gr. Criftatum campellre Lufit. ND. 67 Gr. Panicetten Spica simplici. 68 Gr. Panic. Spica parva leniter hirsuta. 69 Gr. Paniceum tpica afpera. 70 Gr. Panie, spieh ariftis longis armata, 71 Gr. Panie, plurimis spiers congesta panicula, 72 Gr. Panie, plurimis spiers per culmum sparsis, 73 Gr. Phalaris altera Lusie, ND, 74 Gr. Phalaris bulbofi 3 species, ND. 75 Gr. Phænix zulgare. 70 Gr. I hanix alata venustissima, ND. 77 Gr. Phænix apicibus donata; ND. 78 Gr. Phænix calva spici, ND. 70 Gr. Phænix calva falcata spici minima, ND. 80 Gr. Phanix multiplicibus granis spicata, ND.
81 Gr. Phanix ratioregrano longius spicata, ND.
82 Gr. Phanix multis tenuibus granis, spicata elegans, ND.
83 Gr. Phanix pluribus ten. granis dense spicata, ND.
84 Gr. Phanix puribus ten. granis dense spicata, ND. 84 Gr. Phænix ramofa maritima, ND. 85 Gr. Phænix pilosi spica, ND. Gr. Secal, bullof, spic, longa multis longisque aristatis Lufit, ND. 87 Gr. Spart, argent, caudata spica, rad, longa Luf. & Batic. ND. 88 Ur. Tribinum longa cauda spicata, ND. 89 Gr. Tribuloides majus & minus. 90 Gr Dadylon (picis villofis Lufit. 91 Gr. dupliei seu medio aperta spica, ND. SECT. 2. Frumența & Gram Paniculata. 92 Ægilops barbata, ND. 93 Arundo Cipria seu Donax minor. 94 Avena sterilis. Balanco Lusit. 95 Gr. Avenaceum Alopecuroides, ND. 96 Gr. Avenac. utumnale, ampla panicula, ND. 97 Gr. Avenaceum tult ofum Lufit. ND. S Gr. Aven. capillari tenuissimo folio, ND. 99 Gr. Aven. elegans Salacienfe, ND. 100 Gr. Aven. longa contractiore panicula. 101 Gr. Avenac. pratense Lusitanicum. 102 Gr. Aven. Tremulum montanum elegans, ND. 103 Gr. Bromoides paniculatum molle marinum, ND. 104 Gr. Brom. panic. molle, arift. recurvis obsitum elegans, ND. 105 Gr. Festuca Avenacea maxima & major, ND. 106 Gr. Festuca Avenacea minor & minima, ND. 107 Gr. Xe ampelinum latif. minus, sparsa panicula. 108 Gr. X: tampel, latif. brevi panic. longius radicatum. 109 Gr Xeramp, tenui brevissimo folio, panic, contractiore.
110 Gr. Amabile tremul, max. panicula alba & spadicea. 111 Gr. Amab. trem. maj. panic. copiosis elegantismis. 112 Gr. Amab. trem. minus speciosum. 113 Gr Amab. trem. minimum elegans. Bollebolle Lufit. 114 Gr. marinum Paniculosum. 115 Gr. Miliaceum leget. Lusitanicum. 116 Gr. Miliaceum aftivum vestibus infestum. Lufit, 117 Gr. Milise. pumilum elegantissimum, ND.
118 Gr. palustre Paniculatum.
119 Gr. Plumosum parietinum pulcherrimum, ND.
120 Gr. Pratense brevi sol. rad. nodosa Vernum. 121 Gr. Segetum panicula sericea contracta, ND. 122 Gr. Segetum panicula sericea sparsa, ND. 123 Gr. Sorghi panicula erecta. 124 Gr. Sorghi panicula alterum. 125 Gr. Sparteum Avenacea panic. dumetorum, ND. 126 Gr. Sparteum paniculà comofa.
127 Gr. Sparteum panic. brevi, foliculo inclusa.
SECT. 3. Gr. Miscellanea & incognita.
128 Gr. Cuspidatum odoratum pratense, ND. 129 Gr. Geranites intractabile, ND, 130 Gr. Torofum aquaticum, ND. 131 Gr. Torofum campeftre. ND. 132 (1. Triviale ruder. & muror. Miliacea pendula panicula. SEC 1. 4 Cyperi, C peroides & Juncea. 133 Cyperus radice tong a odorata. Albafor Lusit.
134 Cyperus radice rotundà odorata minor Lusit. ND. 135 Cyperus radice esculenta seu Trasi. Junzas Lusit.
136 Cyperus radice rotunda inodora Lusit. ND. 137 Cyperus tadice socunda inodora marina, ND.
138 Gr Cyperoides caudatis spicis.

139 Gr. Cyperoides caudatis spicis intortis.

140 Gr. Cyp. Vetonicæ Caryophylli fol. spica simplici. 141 Gr. Cyp. Vetonicæ Caryoph. fol. triplici & quadruplic i spica.

142 Gr. Cyperoides echinatum. 143 Gr. Cyp. Cyperi veri umbellis, venustissimum pufillum annuum Autumnale. 144 Gr. Cyp. pumilum Autumnale elegans, ND.
145 Gr Cyp. pumilum Autumnale, globosis capitulis, ND.
146 Gr. Hirsutum nemorosum aliud Luste. ND. 147 Gr. Holosteum monospermon Luste. ND. 148 Gr. Holosteum polyspermon Luste 149 Gr. Holosteum Matth. plur. granis, capitula congestis, ND. 150 Gr. Holosteum idem apice ornatum ND. 151 Gr. Junceum capitatum, ND. 152 Gr. Junceum conatum, ND. 153 Gr. Junceum conatum, apice donatum, ND. 154 Gr. Junceum plur. capitulis in unum congestis. 155 Gr. Juncoides, contractiore panieula, ND. 156 Gr. Juncoides sparsî panicula Lust. 4 species. 557 Gr. Spicatum majus & minus marinum. 158 Juncus seu scirpus medius. 159 Juneus kevis panieula plur, glob. medio eulmo prodiens, 160 Juneus exiguns palustris. COM, III. Fossils Varia à Clar. D. Georg. Anton. Volckman Lignicii Silesiæ. 161 Granati prope Freywalde in Bohæmia. 162 Granati ex monte Granatorum prope oppidum Schmeide-163 Amethystus in radicibus montis Riphai ad pagum Krummem-bubel. 164 Jaspis ruber ex monte Boschelberg dicto, propè Urbens Landeshudt. 165 Smaragdus ex m. Riphao seu Giganteo. 160 Topazius candidus monte Ripheo ad lacum majorem. 167 Acinera Ori in granis rarissima Germ. Goldbohnen ex 3 fodina ad rivulum rubrum propè pagum Schreibers-han. Lavantur etiam in ipso jam dicto rivulo. 168 Æs rude Plumbei, purpurei & aurei coloris ex Metalli-fodinis Kupfferbergi. 169 Marmor Masselense ex Ducatu Olfnensi, cum Turbinulis, Rotulis & aliis marinis. 170 Terra Medica alba cruda Lignicensis. 171 Terra Sigilata alba Lignicensis. 172 Terra Sigilata lutea (potius cinerea) Lignicensis.
173 Jolithos seu Lapis Violaceus, in petrosis inditorum superciliis ut in cacumine monte Giganteis. Calefactus odorem Spirat Viola Martia, & reponitur ad Vestimenta ut iis odorem consiliet & Tineis præservet.

174 Lapilli Ophthalmici è Thermis Lignicensibus in villà Grunthal. Oculi expurgant si quid in eos infiderir.

Lapis cum Coralliis punctulatis, Entrochis sey Rotulis, Petunculis, & aliis marinis refertus, ex monte arenofo prope Cuntzendorff in Ducatu Schwidnicensi. 176 Silices circulares monte Giganteo. 177 Dendrite ex Plumbi & Argenti fodină. Wags mitt Cott dicte, ad opidulum Gotts berg in Ducatu Janraviensi. 178 Dendrite in Sabuletis propè pagum Schonborn & Camerau ad Schwidnicium. 179 Fteri-lithes cum Filice minore, è Carbonariis ad pagum 179 Feeri-lithos cum Filice minore, è Carbonariis ad pagum Hernmisdorff propè Gotts-berg.

180 Pteri-lithos cum Adiantho albo Filicis folio CB. è Carbonariis ad pagum Alt-wasser propè Schwidnicium.

181 Lithoxilum dense striatum & articulatum è Lapidicina Lanssudgens juxta Templum. 182 Semina varia è Carpolithis excisa. 183 Volithus aliàs Hammites, vulgo Pifa Carolina, ex Thermis Carolinis in Bohamia. COM. IV. Trees and Shrubs growing in Carolina, Virginia and New England, with their Vertue, Ufe, &c. SECT. 1. Timber-trees and Shruhs with dry Fruit, as Acorns, Cones, Burrs, Carkins, Nuts, &c.

184 Ash-tree. Is tough like our English, but differs something in the leaf, but more in the Bark, and bears no 185 Water-Ash. Is brittle, the Bark is Food for the Be-186 Afpen tree. It grows in Rapahannock River,
187 Beech. It grows very large; 'tis not a durable Timber,
and therefore little used, except for Fire wood.
188 Buck-beech. Another fort found in Carolina, &c.

pril are much eaten by the l'araketoes.

189 Birch. Grows on the Banks of Rivers; its Buds in A-

Bald Cypress. Because it loses its Leaves. Its Nuts yeilds a most fragrant Balfam, which cures all green Wounds and Cuts; its Body they scoop into Canoes and Pereaugers, some so large as to carry 30 Barrells, of one entire Piece of Timber .

191 Chestnut. Is a large and durable Wood fit for Pallifadoes, House-frames, Thresholds, &c. its Nuts finaller than ours.

but fweeter.

192 Chinkapin. A fort of Chestnut with a small round Nut, and the Kernel much sweeter : It is used to Timber Boats,

Highland Elm. The Indians use its Bark for green

Wounds.

Low-ground Elm. Of this Bark they make Ropes.

Sweet-gum. Because in the Spring it yeilds a fiagrant
Gum, upon cutting its Bark or Wood, of great use in Tetters, Scurfs, Inflammations, &c. it bears a round prickly Bur. The Wood has a fine Grain, fit for Drawers, Scrutores, Tables, &c.

106 Horn beam. It grows plentifully in some Parts.

197 Maple. There are two forts, used to make Trenchers, Spinning wheels, &c.

198 Black-Oak. A durable Wood under Water, and bears a

good Mast for Hogs.

109 Chestnut-Oak. From the largeness and sweetness of its
Acorns: A very losty Tree, chiefly growing in low Land.

200 Live Oak. An Evergreen, its Acorns as pleasant as Chestnuts, and yeilds an Oil, as sweet as the Olive.
201 Red Oak. Is a porous Wood, and used to split in Rails for

Fences, it also makes good Clap-boards.

202 Ring-Oak or White-iron. From the durable Quality of its Wood, and is effected the best for Ship-ufe, its Planks being longer than the Live Oak.

203 Scaly White Oak. From its scaly broken white Bark; it is used in building Ships and Sloops: it bears a large Acorn, but never grows to the Bulk or Height of the Chestnut Oak. 104 Spanish Oak. Splits very well into Clap-boards and Lad-

205 Bastard Spanish Oak. Bears good Acorns, and its chief Use is for Fencing and Clap-boards.
206 Turkey Oak. From a small Acorn it bears which the Wild Turkeys seed on.

107 Water Oak. Is an Evergreen, growing in Swamps, Fresh-

water Ponds, and by River Sides.

208 Willow Oak. From the Likeness of its Leaf, grows in Ponds, and is useful in many Things.
209 Pitch-pine. A large Tree with short Leaves; it yeilds Rosin and Turpentine, by the Heat of the Sun, and Pitch and Tar by that of Fire.

210 White-pine with long Leaves.
211 Yellow-pine. Of this and the last they make Masts, Yards, and other Necessaries.

*211 Almond-pine. From the Sweetness of its Kernels, which are Food for Hogs, It serves also for Masts.

212 Dwarf-pine. An Evergreen as the rest are.
213 Spruce or Hemlock-tree. The Fishermen in New England.
tan their Nets and Sails of a tawney Dye, with the Bark of this Tree.

214 The Sycamore grows in Swamps and by River Sides, its Bark finely mottled with Blue, White, &c. bears a Bur like the Sweet-gum.
215 Tulip-tree or Poplar. From its Grain and Bulk, some be-

ing seven Yards about; they saw it into Planks for Houseshingles and other Uses, it makes also, a very pretty Wain-

fcot.
216 Tulip-tree with a white Flower.
217 Black Walnut. A very big Tree, its Wood firm and durable, and some say, never Worm-eaten; they make Tables.
218 and Chests of Drawers of it.
229 Street Street

*217 White Hiccory. Is a fort of Walnut, its Wood will not bear the Weather long; with these Nuts, the Savages make

divers Dishes and Banquets.

- 218 Red Hiccory, the Heart being very red, firm and durable; they make Mortars, Pestles, and other Turnery Ware of it.
- the Shell foft, Kernel bitter and Leaf fragrant; of its Wood they make Coggs for Mills, &c.

 22) Willow. Grows on tresh Water Banks, but differs from

ours in Bark and Leaf.

SECT. 2. Timber-Trees and Shrubs with Berries, &c. 221 Arrow-wood. Used by the Indians for Arrows and Gunflicks, it grows as strait as if plained, and is of all Sizes

222 Prickly Ash. Of this both English and Natives make Poless it growing up like them; it's very thorny and light, bearing Clusters of purple Berries; the Root purges and vomits, used in Cachexies.

223 Red Cedar.

Red Cedar. An Evergreen, its Wood sweet, very durable, and much used for Tables, Wainscot, Cossins, &c.
White Cedar. Like the last, but a taller Tree, and grows strait as an Arrow; its Wood very light, free to split, and makes the halo. St. 1. makes the best Shingles.

225 Dogwood. Bears a white Flower, and blooms the first in the Woods, has a fine Grain; of Use within Doors, but not laiding. The Bark of its Root is held an infallible Remedy for the Worms.

226 Black Gum. Beers a black Berry, which the Indians mix with their Pulse and Soops, it giving a prettey Flavour

and Scarlet Colour.

227 Bitter Black gum. Its Berries like the last, but bitter; the Wood of both these are used for Cart Naves.

228 White Gum. Bears a long Bunch of Flowers, its Wood

much curled and knotted, yet might be of use.

229 Holly two forts, one with a large, the other with a small Leaf, many are 2 Foot diameter; they make good Tren-

230 Honey-tree. Like the next but more prickly; it bears a Cod; one Side contains the Seed, the other Honey, of which they make very good Metheglin.

231 White and yellow Locals. Are pretty tall Trees; the Indians make Bows of them, being tough and flexible.

232 Salfafras. Bears a white Flower and black Berry, the Bark of the Riot is a Specifick in the Gribes and Heart-burn; it's also of other Savereign View. Of its Wood them

burn; it's also of other Sovereign Use: Of its Wood they make Bowls, &c.

SECT. 3. Dying, Medicinal, Ornamental and other Trees and Shrubs.

233 Lauvel tree as tall as Oaks, the Leaves and Berries dye Tellow.

234 Gall berries. A pretty Evergreen Tree, in Swamps and Ponds; it bears a black Berry, with which the Women dye their Cloaths and Yarn black.

235 Indico. It grows plantifully in Carolina, &c.

230 Person vine. Its Leaf like Hemlock, the Juice stains Linnen, never to wash out it marks a blackish Blue Colour, by breaking a Bit of the Vine and writing with it.

237 Candle Bay. Its Berries yeild a fragrant Wax, which befides its Use in Surgery, they make Candles off,
238 Candle Myrtle. Of the Berries of this, they also make

Candles which will stand the hottest Weather and burn very fweet.

239 Sand-bank or Sea Cassina or Yaupon. Because this Kind particularly grows there only; in some Places 12 Foot high, in others but 3 or 4: It grows like Box, whose Leaves it resembles, but thicker and dented; bears a small red Berry, but in December when tipe, is brown.

240 Swamp Cassian. In rich low Ground and Swamps, its

Leaves greener and larger.

241 Dwarf Cassina, never grows 2 Foot high. Sheep and Deer delight much in this Plant. This is their Indian Tea, used and approved of by all the Savages on the Coast of Carolina, and fold at a good Price to the Westward Indians, it grows plentifully on the Coast of Florida, and is held in great Esteem by the Spaniards in New Spain.

242 Pellitory-tree. From the Hotness of its Bark, which chewest in the Mouth cares the Tootheast.

chewed in the Mouth, cures the Tooth-ach. It grows on

Sand-banks on the Sea Shore.

243 Small Bamboo. It grows in low Land, a fort of Clymer, its Reeds rarely thicker than ones Finger, and very tough; the Indians boyl and eat the Root, which is a round Ball, and when dry, so hard that they make Cane-heads of them, which are carried into pretty Figures.

which are carved into pretty Figures.
244 Oak Vine. It bears a fort of Bur, and runs about Trees; it is so porous that they suck Liquors through a Length

of 2 Foot.

245 Dwarf Palmetto, grows in Carolina, &c. 246 Sugar-tree, grows at the Heads of Rivers, and near Mountains; this by tapping yeilds a Juice which they boyl into a Sugar.

247 Sorrel or Soure-tree. Because its Leaves have that Talle; some are a Foot diameter.

248 Red bud. Bears a purple Blossom, ripe in April and May, it's Flower an excellent Sallad.

240 White Honey-suckle. Grows branched and always in low moist ground: This and the following flower in April and May.

250 Jagged Honey-suckle. Because its Flowers are deeper cut,

This grows in clear dry Ground.

251 Swamp Honey-suckle. Grows in swampy Woods and River Sides about 2 or 3 Foot high and bears a great Bunch

of fine white Flowers.

252 Wood Honey suckle. About a Foot high; its Flowers on small Footilalks, several rogether from the main Stock which is northicker than a Wheat straw.

253 Woodlind. Is here ever green and long in Flower.

254 Jalmin Clymer. From the Shape of its Leaf, but larger; it grows round Trees, and makes a fine Shade and a qui k grower; 'tis an Ornamental Plant with a black Berry.

255 Scarlet Trumpet-Vine. Makes a fine Arbour, bears a

large red Beil-Flower and a long Cod.

- 256 Maycock. A Summer Clymer, with a glorious Flower, it bears an Apple, of an agreeable Sweet mixt with Sowre.
- 257 Larks-heel-tree. From the resemblance of its Flowers. 258 Yellow Jasmin. Of a pleasant Smell, growing in Woods. SECT. 4. Finit Trees, Shrubs and Bushes, viz.

250 Red-Cherry. A large Tree in the Woods, not much unlike

the Cornel berry.
265 Cluster cherry. Grows high and in Bunches like Currans,
where the control of the Relish and yield a crimson they have a bitterish sweet Relish, and yield a crimson

261 Black and white American Damfons. These grow on all the Sand banks along the Coaft.

262 Box Flum. From the Yellowness of its Wood, which is fomewhat porous: The Fruit has a Physical Tatte.

- 263 Early Indian Plum. This makes a fine Shew when in Flower.
- 264 Late Indian Pinm. Its Bloffons white, finall, and very fweet, the Fruit red and palatable.
 205 Wild Fig. Grows up the Mountains in Virginia.

266 Common red Mulberry. Is very sweet and one of our ear-

- liest Fruit, next the Strawberry. 267 White Mulberry. Very common and fit for Silk-worms. 268 Black Mulberry. Bearsa finall Fruit, but very sweet.
- 269 Papau. Bears an Apple of the Size of a Hen's Egg, yellow, foft, and very sweet; of these they make rare Puddings, its Leaves very broad and Stones large,

270 Summer Persimon. Agrees with all Soils.
271 Winter Persimon. Because the Fruit ripens not till the Frost as pears.
272 Black Grape. With a pale Flesh-coloured Juice.

273 Black Bunch-grape. leilds a crimfon Juice, grows common and bears plentifully; they are of a good Relift but small, yet have a large Stone and thick Skin. 274 Black Summer Fox grape. Ripens with the next in July,

frequent in Swamps and low moilt Ground.

275 White Summer Fox-grape. This and the last grow not in Clusters, having but 5 or 6 in a Bunch and ripen in the Shade; they are about the Size of a Damson.
276 Black Winter Fox-grape. Ripens in September in all Soils,

but most plentifully on the Sandy Sea Coasts.

277 White Winter Fox-grape. The Fruit clear and transparent, with smaller Grains: This and the foregoing are pleafant, bur all glutinous.

378 Red Haw. Of an agreeable Taste, and sour times as big

as ours in Europe.

- A flender Tree : These are eaten both by 379 Black Ham. Men and Birds.
- 280 Black-berries. Are bitterish, and not so palatable as in England.

381 Blue Bilberry. Not unlike our common English.

- 282 Great Biberry. Grows in Savana's, Meads, and Woods: This has both a larger Fruit and Seed.
- 283 Small Bilberry. As pleasant as the blue, but larger; grows 3 or 4 Foot high, in low good Ground, and Riverbanks.
- 284 Tree Bilberry. Grows 10 or 12 Foot high, and as thick as a Man's Arm, in low Grounds; the Fruit is pleafant and Lears much.
- 285 April Currans. A very beautiful Tree, on the Banks of Rivers, in Clayey Ground; its a pretty red Fruit, but gone auickly.

280 Bermudas Currans. A Bush, grows in the Woods, like our English, and much eaten by some.

Winter Currans. A pretty bushy Tree, about 7 or 8 Foot high, very spreading; are ripe in October: They make Tarts of them, and dry them instead of Currans.

COM. V. Some American Plants, with their Specifick Vertue and Wonderful Effects, lately fent me by the Reverend an Learned Dr. Cotton Mather, at Boston in New England, and Fellow of the Royal Society, London.
288 Ophiophuga, Cottonis Mather.

Hieracium Novæ Angliæ Succife solio, Flor. parvis ramo-

fis, Nob.

A Poultifs of this bruifed and laid to the Part bitten by the Rattle-Snake, it immediately fetches out the Deadly Poyfon: it's also remarkable, that if put into the Shoes, no Scrpent will dare to come near them. A Tea of it is a good Ophthalmick, C.M.

N. B. I have already 4 or 5 different Sorts of these Rat-tle-Snake Plants from Carolina, Virginia and Maryland,

and this, another, altogether new to me. 289 Caltha Sylvatica or Wood Marygold, C. M.

Flos Solis N. Anglia, fol. feffilib.s fcabris, caule rubente, Nob. This is, perhaps, the greatest Vulnerary in the World; a Balfam of the green Leaves, with Oil Olive, cures the most grievous Punstures of the Nerves, and Cuts of all Sorts, with an admirable Efficacy and Quickness, ordinary Wounds in 4 or 5 Hours, to the great Assonishment of the Patient, and extraordinary Cuts in a very few Days at farthest. The Indians will fight like Milton's Devils, if this Traumatick be near them, C. M.

This fort of Americk Chrysanthemum, Dr. Morison calls Vofaean, and now grows in our Phyfick Garden at

Chelley.

290 Great Bur Comfry. A fort of Caryophyllata Avens, or Herb Bennet.

291 Small Bur Comfry. It does Wonders in internal Wounds, relieves Oppressions in the Stomach, from clotted Blood, not to be parallel'd, C. M. This seems another fort of Avens, but being without Flower or Seed, I cannot determine it, J.P.

202 Formofa, C. M. by the Indians Satanschoo. Two Spoonfuls of the Water in which this Herb is boiled, cures the Malady which we call the Stomach-ach, as also Inflammations of the Eyes, being washed with it, C. M. This with us is called the Cardinal-flower, from its beautiful scarlet Colour. Vid. Ray Hist. Plant. 746. 1.

203 Taututtipoag, so called by the Indians.

The whole Plant with the Poor bailed and the state of the sta

The whole Plant with the Root boiled and applied to the Face, cures what we call the Ague there, and also any Tumour; it also does Wonders in Cancers and other bleers by bathing. A Tea of this inwardly, and a Pultiss to the Part grieved, is the grand Medicine here for the foul Disease, C.M. This is a fort of Geranium

Batrachoides, longius radicatum, Ray 31. p. 1061.

201 Fagiana or Cancer Rape, C. M. Orobanche ramofa N. Anglia, fub Fagis nascens, Nob. A Pultis of this Plant laid to the Part, and a Ten of it inwardly, does Wond: 18 in Cancerous and Scrophulous Cases, as also in Malignant-Fevers; it strengthens and invigorates after Fatigues and Westinels. Grows only under Beech Trees in spady Places, from Midsummer till Autumn.

COM. VI. Divers Miscellang Curiosities sent me from Bombay and Callecut, by Mr. Alex. Christie, Surgeon, residing in

those Parts.

295 Fapilio major è nigro & pallidè cærulescente, alis inferi.

oribus finuosis maculisque atris ornatis, è Callecut.
296 Libella major, apicibus alarum lanceatis, alis inferioribus amplis maculis è Callecut.

amplis macuits e Cauctur.

The following Shells from Bombay.

297 Cochlea auriculata, fafciis castaneis, pulchrèobliquis Gazophyl. Natur. Tab. 101. Fig. 9.

298 Cochlea umbilico magno sinuolo, maculis castaneis. Gaz. Nat. Tab. 101. Fig. 10.

299 Buccinum oblongum minus, costis obtusis parce elatis. Gaz. Nat. Tab. 101. Fig. 14.

300 Buccinum breve minus, costis dense positis. G. N. T. 101.

fig. 15.
301 Murex breve minus, fasciis spinosis. G. N. 101.18.
302 Veneroides exilisalba, fasciis duobus rubelcentibus. G. N. 102, f. 13.

303 Buccinum dentatum album, costis obliquis. G. N. 102.

304 Rhombus fasciis capillaribus suscis, cingulo medio alba. G. N. 101. 17.

305 Operculum album, intus auritum. G. N. 102. f. 21. Rubeola erecta, spica Typhoides carulei. swampy Grounds, the latter End of the Rains. Found in

GRAMINUM, Muscorum, Fungorum, Submarinorum, &c. BRITANNICORUM

NC

A Methodical Concordance of British Grasses, Mosses, &c. in Order to their being more easily known and better Figur'd.

AJACOBO PETIVER, Regia Societatis Socio, &c. Londini.

CAP. I. P. Rumenta & Gram. Spicata,
Spiked Corn and Graffes.
Sed. 1. De Triticis & Gram. Triticeis.

Sett. 1. De Irsticis & Gram. Irstices.
Wheat and its Graffes.

1. Naked Red Wheat. Bobart. Hift. Pl. Oxon. p. 175. I.
Triticum nudum, spica & granis rubentibus, Ray H. Plant.
127. I. Synopf. 177. Pl. 1. edit. 2. p. 244. I.
2. Naked red straw'd Wheat. Bob. H. Ox. 175. 2.
Inicum nudum, cullmo & granis rubris.
White Laumas Wheat. R. Ox. 176. 4. Tab. 1. fig. 1. opt.

Triticum nudum, culmo & granis rubris.

3. White Lammas Wheat. R. Ox. 175.4. Tab. 1. fig.1. opt. Triticum nudum, fpich & granis albis, Ray 1237. 2. Syn. 172.

2. ed. 244.2. C. B. 21. pl. 1. Phyt. 46. 1. Theatr. 352. fig. Mattb. 317. f.2. Lugd. 378. f. Tabern. 258. Ic. 2. Hist. 735. f.1. Ger. 58. fig. Dod. 489. f. Lob. 25. Ic. Belg. 27. I. Obf. 14.1. Adv. 5. Jonf. 65. f. Park. 1120. 1. Chabr. 173. 1. 4. Mixt Lammas Wheat, Bob. Ox. 175. 3.

Triticum nudum, spich albicante granis rusel centibus, seu mixtum. N. B. Most of these have sometimes short, but irregular Reards and Flower about Midsummer.

Beardi and Flower about Midfummer.

5. Cone. Wheat, Boot. Oxon. 176. pl. 13. T. 1. f. 2. f. opt.
Initiam ariftatum, fpied villosa quadrata longiore.

It's a tall Corn, composed of four Rows of Grain, set in grey hairy Glumes or Husks in a large Spike near 6 Inches long,

It's a tall Corn, composed of four Rows of Grain, set in grey hairy Glumes of Husks in a large Spike near 6 Inches long, with very sough and long Beards.

6. Grey Wheat or Duckbill, Grey Pollard, Ray 1238. 8. Spn. 178v 3i; ed.: 245. 3. Triticum aristatum, spica villosa breviore & turgidiore, B. Ox. 176. 14. T. 1. ser. 2. f. opt. Its pikes grosser, shorter and whiter, and rises not so high.

7. Naked Duckbill. Triticum nudum spica villosa quadrata breviore. Differs from the last in being beardless.

8. Red. ear's bearded Wheat. Triticum aristatum, granis & spica subaribus, glumis lævibus & splendentibus, Ray 1238. 6. S. 177 2. ed. 244. 2. B. Ox. 175. 2. T. 1. s. opt. CB 21. 8. pt. 47. 6. Th. 371. f. Tab. 259. 2. H. 735. 3. G. 59. 4. Lob. 27. 2. B. 28. 3. Jonst. 67. f. Park. 1120. 3. Lugd. 377. 1. Tabernemont's Figures and Lobels seem somewhat to differ, but not much. This holds its Beards, tho' sipe.

9. Many-ear'd Wheat, R. 1238. 4. S. 177. 3. d. 245. 5. Tritium aristitum spica multiplici, Bob. 175. 7. T. 1. f. boni. CB. 21.6. pt. 48.8. Th. 371. f. Tab. 260. 1. H. 736. 1. G. 59. 3. Matth. 318. f. Chabr. 173. f. 2. opt. Lob. 26. 2. L. 28. 1. Jonst. 66. 4. P. 1120. 4. This is sometimes sow'd about London.

10. Polonian Wheat, Ray Vol. 3. p. 597. 3. S. 236. ed. 245. 4. It is sometimes fow'd about London.

10. Polonian Wheat, Ray Vol. 3. p. 597. 3. S. 236. ed. 245. 4. It is sometimes sow'd about London.

745. 4. If Truicum Polonieum majus, gluma foliacea, Pluk. Tab. 231. f. 6. opt. B. Ox. 175. 8. Tab. 1. fer. 2. f. bona. Sown in Wortesfermire.

II. Barly Wheat, Ray 1908. S. 236. ed. 245. 6.

Initicum ariff. spic. Hordei polystichi, B. Ox. 176. 11. T. I.

1. 2. f. opt. CB. 22. 4. Th. 423. f. Tab. 262. 1. H-752. f.

Ger. 66. f. I. Chabr. 175. f. 5. J. B. 430. Lob. 30. f. 2 B.

37 2. Ob. 18. 2. J. 72. f. P. 1123. Its Spike like fquare Barley

of fix Rows; bearded, but the Grain of Wheat. Sown about Ishp

and other Places in Oxfordshire.

Gramina Tritical. Wheat. Graffer

Gramina Triticea. Wheat Grasses.

12. Common Dogs-grass. R. 1259.1. S. 179. I. d. 247. I. B. Ox. 178. 8. T. I. s. 3. s. opt.

G. Caninum offic. vulgare. CB. I. pl. 1. pt. 3.1. Th. 7. f. Tab.

201. 2. H. 637. 2. G. 22. I. Lob. 20. 2. B. 21. 2. Ox. xi. 2. Ad. 2. P. 1173.1. Dod. 558. f. mala. 23. I. No good Figure at yet of this Common Grass, except Mr. Robart's Spike of it. Too frequent in Gardens and some Fields in June, &c.

13. Bearded Dogs grass. Gr. Caninum aristatum vulgare, R. 1255. 2. Var. S. 179. I R 247. I. Differs from the Common in being bearded.

4 - 6 - 2 - 2 - 2

14. Bearded Wood Dogs-grass. R. 3. p. 598. 1. S. 235. d. 247. 2. Gramen Canin. sylvaticum aristatum elatius, Bob. Ox. 177. 2. T. 1. s. 2, f. opt. In Stocken-Church Woods plentifully.

July.

15. Sea-Dogs grafs. R. 1255. 6. 5. S. 179. 2. d. 247. 3. Bob. 178. xi T. z. f. Gramen Caninum marit. vulg. CB. 1. 8. pt. 3. 3. Th. 14. f. Tab. 203. 2. H. 637. 4. G. 23. 1. Lob. 21. 2. B. 22 I. I. 25. I. Chabr. 186. 6. Common in Salt Ditches and Sand Banks in June, July. &cc.

15. Bearded Sea-Dogs grafs. Gramen Caninum marit. aristatum. A Variety of the last, and found with it, but more

17. Buddles Sea-Dogs-grafs. Gramen Caninum maritimum fpicis ravioribus crassis Instit. Rei Herb. 516. First Discovered by that accurate Bosanist the late Reverend Mr. Adam Buddle,

on the Suffalk Coast towards Yarmouth in July.

on the Suffilk Coast towards Yarmouth in July.

18. Swaddle headed Sea Dogs grass. R. 1256. 3. S. 179.3.
d. 247. 4. Gr. Caninum marit. spica soliaced. CB. 2. 9. pt.
3. 4. Th. 15. f. opt. Tab. 203. 1. H. 638. 1. G. 23. 2.
Lob. 22. f. B. 22. 2. P. 1277. 3. B Ox. 178. 12. T. 2. se
Found by the Curious Mr. Sam. Dale in Mersey Isle, near Colchester. It Heads in June and July.

SECT. II. De Loliis & Loliaceis. Darnel and its Grass.
19. Corn Darnel. R. 1262. c. 7. pl. 1. S. 181. 1. d. 249.
1. B. Ox. 181. T. 2. st. f. Lolium segetale CB. 9.1. pt. 23.
pl. 102. Th. 121. f. Matth. 331. f. Tab. 239. 2. H. 675. f.
G. 71. 1. Lob. 35. 1. B. 44. 1. O. 21. 2. Ad. 10. Dod. 538.
P. 1145. 1. J. 78. 1. Lugd. 417. f. Found too often amongst
Wheat in July, &c.
20. Bearded Darnel. Lolium segetale aristatum. B. Ox.
T. 2. st. 2 f. opt. Grows with the last, but more rare.
21. Tall Brize Darnel. R. 1257. c. 2. pl. 1. Syn. ed.
248.

248.

248.

Gramen Spica Briza majus CB 9. 1 pr. 18. pl 56. fig. Th. 133. P. 1144. f. 5. B. Ox. 205 4 T. 6. f. 3. f. Pretty common about Oxford in Copfes and Hedges.

22. Common Way Darnel. R. 1263. 3. S. 181. 2. d. 249. 2. B. Ox. 182. 2. T. 2. f. opt. Gramen Loliaceum vulg. CB. 9. 3. pt. 22. 101. Th. 128. f. Lob. 34 2- O. 21. 1. Ad. 10. Dod. 540. f. J. 78. 2. P. 1145. 2. Lugd. 1416. f. Frequent by Path-ways and in Graffy Paftures.

23. Bearded Way Darnel. Gramen Loliaceum vulg. ariftatum.

24. Thin Darnel grafs. Gramen Loliaceum, spicis rarius dispositis. B. Ox. 182. T. 2. s. 2. f. opt. gr? P. 1145. 4. f. 25. Broad Flat Darnel-grafs. Gramen Loliaceum spicis brevioribus compressis. B. Ox. 182. T. 2. s. 2. f. opt. gr?

vioribus compress. B. Ox. 182. T. z. s. f. z. f. opt. qr? P. 1145. 5. f. o.

26. Branched Darnel-grass. Gr. Loliace. multiplici spicata panicula. R. Syn. 235. B. Ox. 182. T. z. s. f. z. f. opt. P. 1146. f. opt. These 3 last grow in Meadows, &c.

27. Sea Darnel-grass. Gramen pumilum Loliaceo simile. Ray S. 182. 4. d. 250 4. Pluk. Tab. 3z. f. 7. opt. qr? Ray 1287. 18 f. 190. 15. d. 259.16. qr? B. Ox. 182. 6. T. z. s. f. z. f. qr? P. 1165. f. & 1278. 6. Found in the Isle of Sheppy by Mr. James Sherrard and my self. It spikes in July, &c.

28. Rush Darnel-grass. Phanix marit. spica aculenta. R. 1263. 5. S.-182. 3. d. 250. 3. Ger. l. 1. c. 2z. pl. 8. Park 1146. f. 6 opt. B. Ox. 182. 8. T. z. s. s. f. 2. f. opt. Frequent in Salt Marshes in July, &c.

29. Small Heath Mattweed. R. 1260. 8. S. 180. z. ed. 248. 2. Spartum minimum Anglicum. Park. 1199. f. 7. opt. qr? B. Ox. 217. 8. T. 7. f. 3. f. qr? CB. Th. 70. f. On Putney-Heath plentifully, Flowring from the End of April, and holds its Spike till Winter.

Sect. III. De Secali & Gram. Secalinis. Rye and its Grass. 30. Rye. R. 1241. I. S. 178. d. 245. B. Ox. 179. I. T.2.

f. 1 f. opt.

Secale Hybernum vel majus, CB. 23. 2. pt. 50. 2. Th. 425. f.

Matth. 325. 1. Tab. 258. 1. H. 720. 1. G. 61. f. Lob.
28. 1. B. 31. 1. O. 15. 1. Ad. 6. Dod. 499. fig. J. 68. f. P. 1128. 1. Lugd. 396. f. 1. Sown in autumn in light fandy Soil. It's in Spike a Month before Wheat.

30. * Burnt Rie. Secales Uftilago. CB. 24. 1. pt. 52. 1.

Lob. 35. 2. B. 45. 1. O. 22. 1: Ad. 11. Jonft. 77. 3.

P. 1128. f. 2. Found amongst Rie.

31. Small Rie. Secale Vernum vel minus, CB. 23. 3. pt. 50. 2. P. 1129. 2. B. Ox. 179. 3. Lugd. 396. f. 2. It's a smaller Corn, with fewer Stalks and a stender Spike. It's fown in the Spring.

32. Wild Rie, or Wall Barley, R. 1258. 1. S. 180. 1. d. 248 1.

Secale Viarum, Nob. CB. 0. 2. pt. 24. 100. Th. 124. fig.

Secale Viarum, Nob. CB. 9. 2. pt. 24 109. Th. 134. fig. Tab. 235. 1. Hist. 670. 1. G. 66. 2. Lob. 30. 1. B. 37. 1. O. 18. 1. Ad. 7. P. 1147. f. B. Ox. 179. 4. T. 6. 53. f. Very common on Walls, Waste Grounds and Path-ways, from

May till Autumn. 33. Great Wild Rie. Secale spontaneum majus.

33. Great Wild Rie. Secale Ipontaneum majus. qr? Pirk. 1147. 2. CB. 9. 1. pt. 24. 110 Thal. 51. Mr. Buddle observed this amongst Corn about Barns in Survey.

34. Meadow Rie, R. 1258. 2. S. 180. 2. d. 248. 2. B. Ox. 179. 6. T. 2. f. 1. f.
Secale Pratense, Jonst. c. 22. pl. 4. qr? majus. Park. 1144 fig. Frequent in Meadows from May till mowed.

35. Wood Rie, R. Vol. 3. p. 599. 11. S. 235. d. 248. 4
G1. Secalinum majus sylvaticum, Bobart. H. Ox. 180. 12
qr? Park. 1144. 7. fig. In Stocken-Church Woods plentiful-In Stocken Church Woods plentifulqr? Park. 1144 7. fig. ly in June, &c.

36. Short Sea Rie, R. 1258. 3. Syn. 180. 3. d. 248. 3.

Piuk. 33. f. 2 opt.

Secale maritimum minus, Nob. B. Oxon. 170 5. T. 6. f. 3. Its Heads flort, with spread Brisles. In Salt Marshes from June till augumn

37. Fnglish Matweed or Helm, Jonst. 42 f. 3. R. 1259. 3

S. 180 1. d 248. 1.

Spartum Anglicanum, spica Secalina, Nob. CB. 5.3. pt. 25. 113. Th. 67. f. Cluf. 121. f. 1. Tab. 230. 1. Hitt. 674. f. G. 38. 3. P. 1198. 3. Chabr. 192. 2. Lob. 89. 1. B. 122.1. O. 45. 3. N. B. The fingle Spike in Bob. Ox. T. 4. under fig. 16. very well represents this Grafs, which I have observed ved on gurown Coasts in divers Places, viz. the Iste of Sheppy,

ved on our own Coasts in divers Places, viz. the Isle of Sheppy, Deal, &c. on the fandy Shores, from July to Autumn.

38. Buddle's Sea Matweed. Spartum Essexianum, spick gemina clausa, Hort, nost, sice. Ray 3. App. 248 The Reverend Mr. Adam Buddle first discovered this Grass, A. D. 1703 in the Marshes on the River Wallsteet near Fambridge. Ferry in Dengey-Hundred in Essex since which that diligent Botanist Mr. James Sherard found it plentifully in the Swail near the Mouth of Feversham Creek about August and September.

SECT. IV. De Hordeis. Of Barley.

24. Common Rayler, R. 1243. C. 5. I. S. 178. d. 246.

3y. Common Bailey, R. 1243. c. 5.1. S. 178. B. Ox. 206. T. 6. fer. 3. f. opt. Hordeum diffichum, CB. 23. 3. pt. 50. 3. Th. 440. f. Tab. 474. 2. H. 764. 2. G. 64. 2. Lob. 29. 2. B. 33. f. ult. Dod. 501. 2. J. 70. 1. P. 1130. 1. Lugd. 398. f. Sown in the Spring, and ripe in Jugust.

40. Battledoor, or Sprat-Barley, R. 1243. 2. Syn. 178. 2.

d. 246. Hordeum distinum, spica brevi lata, B.Ox. 206. 2. T. 6. 1. 3. f. opt. Zeocrithon s. Oryza Germanica, CB. 22. 5. pt. 48. 3. Th. 422. c. 15. Chabr. 175. f. 4. J. B. 2. l. 18. c. 15. p. 430. f Grows lower than the Common, with short flat Spikes and thick Grain. Sown in the Spring.

Spikes and thick Grain. Sown in the Spring.

41. Winter, Square or Bear Barley, R. 1244. 3. Syn. 178.

2. d. 246. 3.

Hordeum Polystichum hybernum, CB. 22. 1. pt. 49. 1. Th. 438. f. Tab. 274 1. H. 764 1. 1 ob. 28. 2. B. 32. 1. O. 15. 2. Ad. 2. Dod. 501. 1. Jonit. 70. 2. P. 1130. 2 Lugd. 399 f. B. Ox. 206. 3. T. 6. f. 3. f. Its Spikes are commonly 4 rowed. Sown in the Autumn, and therefore more frequent in the North, where they call it Big.

42. Burnt Barley. Hordei Ustilago, CB. 24. 2. pt. 52. 2. Lob. 29. f. B. 33. f. J. 70. f. P. 1130. f. It's a smutty Blite, sound a nongst other Burley.

Sect. V. De Paniceis. Panick and its Grasses.

Sect. V. De Paniceis. Panick and its Graffes. Sect. V. De Paniceis. Panick and its Grayer.

43. Naked Logs Panick, R. 1261, 2. S. 181. 2. d. 249.

2. B. Ox. 189 15. T. 4. f. opt.

Gramen Paniceum spica divila, CB. 8. 1. pt 20 92. Th. 136.

f. Tab 279 2. H. 814 2. G. 79 5. Lob. 42. 2. B. 53.3. O 25 2 Ad 15 Dod 559 2 have met with this about Fulham Jonft 85 5. P. 1154 1. 1

1. 1 2.4

44. Single, smooth Panick Gramen Paniceum spica fim-144. Single, thootic ranks Glamen Panetum tpice in-plicitizer, R 1261 r S 180 r d 249 1 B Ox 189 ro T 4 f qr? CB 8 3 pt 20 94 Th 138 f Tab 228 r H 653 1 G 20 8 f qr? Lob 13 2 lt's fometimes found in Stability Fields in Stubble Fields.

in Stubble Fields.

45. Rough Panick, R. 1262. 3 Syn. 181. 3. d. 240. 3. B. Ox. 189. 11. T. 4. f. opt.

Gr. Paniceum spica aspera, CB. 8. 4. pr. 21. 95. Th. 139. f. Tab 200. 2. H. 636. 1. G. 14 1. J. 15. 1. P. 1172.

2. Chabr. 184. 3 Found about the Neat-bouses, Putney and Rowhampton. It spikes in July and August.

46. Rough Fersey Panick, R. 1265. 7. S. 239 d. 251.5.

Gr. Alopecuroides spica aspera, CB. 4. 6. pr. 10. f. 2. Th. 59. f. pr. 13. 51. P. 1168. 6. Chabr. 185. f. 6. Lob. 45.

2. B. 57. . Lugd. 432. f. 2. In all the landy Grounds in Fersey. Dr. Sherard.

47. Cone Panick, R. 1288. 25 S. 190. 19. d. 259. 20. B. Ox. 189. 12. T. 4. f. opt.

H. Cone Panier, R. 1200, 27

B. Ox. 189, 12. T. 4. f. opt.

Panicum ferotinum spica pyramidata, Instit. 515. B.Ox.

189, 12. Pluk. 33. f. 6. Found amongst Corn in July.

48. Soft Sea-Panick, R. 1266 9. S. 183. 5. d. 251. 4.

CB. 44

Panicum maritimesh villolum Nob. Bob. Ox. 191. 3. T. 4. Lz.
f. opt. Adv. 469. defer. 1. B. z. J. 18. p. 474. Park,
1106. 3. f. & qr? P. 68. 3. f. onf. 87. z. f. o. In Sale,
Marsher in July &c.

VI. De Alopecuris & Typhinis. Fox and Cats-tail Graffes, 49. Common steadow Fox-tail, R. 1264. 17 S. 182, 1.

Gr. Alopecuroides prat. vulgare, Nob CB, 4.1 pt. 11. 43.
Th 55. f. Tab. 210. 2. H. 642. f. G. 9. 1. P. 1167. 2.
& qr? 64. f. 1. & 3. Lob. 8. 1. B. 10. 1 Chabr. 199. 3.
Jonft. 10. 1. B. Ox. 191. 8. Very common in Meddows in April. &c. April, &c.

70. Slender Corn Fox-tail. Gr. Alopec, segetale, Nob. R. 1261. 4. S. 181. 3. d. 251. 1. CB. 4. 5. pt. 12. 49. Th. 53. f. Tab. 209 1. H. 641. 1. G. 9. 4. P. 1169 f. Lob. 9. 2. B. 11. 2. Jonst. 11. 4. H. Ox. 192. 12. T. 4. s. 6. opt. qr? Chabr. 185. 5. Frequent in Corn, & from May to

51. Dales Fox tail with crooked Beards. Gr. Myofuroides spice breviore, ariflis recurvis, R. 1269. 8. 184. 1. d. 251. 2. B. Ox. 192. 17. In moist Places and Path-ways. It spikes in Autumn. Spikes in Autumn.

Pinds in way and June.

53. Long Cats tail Grass. Gramen Typhinum maximum, CB. 4. 1. prodr. 10. 25. f. pt. 11. 44. Th. 49. f. Park. 1170. 1. f. H. Ox. 193. i. T. 4. f. 3. f. qr? Chabr. 185. 3 I sometimes meet with a Grass in Meadows which these Fi-

gures resemble.

gures refemble.

54. Common Cats tail Grass. Gramen Typhinum vulgare,
R. 1267. 2. S. 183. 1. d. 251. 1. CB. 4. 1. Th. 51. ft
Tab. 217. 2. H. 655. 1. Ch. 185. 1. G. 10. 1. B. Oz. 1942.
In Path-ways by Hedges and Pastures.

55. Bulb Cats-tail Grass. Gramen Typhinum bulbosum,
R. 1267. 3. S. 183. 2. d. 251. 2. CB. 4.4 pt. 46. & 7.
Th. 52. f. Chabr. 185. 2. & 4. Tab. 218. 1. H. 655. 2.
G. 10. 2. Lob. 10. 1. B. 111. 3. O. 10. 1. Dod. 562. 1.
J. 12. 1. Lugd. 433. 2. Park. 1170. f. 2. & 3. Bob. Ox.
194. 3. T. 4. s. f. It's less, and found with the Common in June, July, &cz.

56. Sea Cats-tail Grass. Gramen Typhinum maritimum minus, R. 1267. 4. S. 183. 3. d. 252. 3. qr? Park. 1170.
f. 4 I have found this on the fandy Shores both here and in Holland. It Heads in June, &cc.

Holland. It Heads in June, &c.:

SECT. VII. Gramina Cristata, &c.:

Crested and other Grasses, S. Smooth Crested Grass.

Gramen Cristatum vulgare, R. 1269. 3. S. 184. d. 252. 2. CB. prodr. p. 8. pl. 18. Adv. 467. s. opt. Park. 1160. f. 3. These two Figures very accurately represent this Grass, so common in our Meadons.

58. Square Crested Grass. Gramen Cristatum quadratum, R. p. 603. 2. Syn. ed. 252, I. A luxuriant Kariety of the Common.

59 Loole Crested Grass. Gramen Cristatum spica laxa, CB. 3.1. pt. 5. 12. pr. 8. 18. f. Th. 43. f. P. 1160. 1. f. Chabr. 184 2. I have met with this loose spiked Grass

in Greenwich- Fark and elsewhere, but not common.

10 Hairy Crefted Grass. Gramen spica cristata subhirsutum,

10 Hairy Crefted Grass. Gramen spica cristata subhirsutum,

10 Hairy Crefted Grass. Gramen spica cristata subhirsutum,

10 Hairy Cowns. I have sound it on Black Heath.

10 Blue Crested Grass. R. 1270.13 Syn. ed. 253. 262 &

Gramen glumis variis, CB. 10. 8 pr. 21. 73 f Th. 158. f Park. Gramen glumis variis, CB. 10. 8 pr. 21. 73 f Th. 158. f Park. 1152 f 0. B. Ox. 192. 16 Its Spikes are first blue, then pale and white. Found in Westmorland.

61 Mr Ray's Small Spring-Grass. Gramen parvum præcox, stiel laxà canescente, Ray Syn. 191. 21 d 260. 22 Pluk. 33. f pr. 192. 24 qr? Ray 1271. 17 S 184. d 253. In and parren Soil, from the midst of April till after Mid-

SECT. VIII. Gramina Miscellanea. Miscellany Graffes. 62 Rough Grafs, R. 1287. 19 S. 190. 16 d 259. 17. B. Ox.

62 Rough Grass, R. 1287. 19 S. 190. 16 d 259. 17. B. Ox. 102. 38 T 6. f opt.

Gramm spicatum aspero solio, CB 3. 5 pt. 5. 11 pr. 9. f 1. opt. Th. 45. f Chabr. 183. f 4. Lugd. 427. f 2. Park. 1160 opt. Th. 45. f Chabr. 183. f 4. Lugd. 427. f 2. Park. 1160 opt. Th. 45. f Chabr. 183. f 4. Lugd. 427. f 2. Park. 1160 of 7. Very common in Passures and by Hedges, from May till ham.

63 Doody's Rough Grass. Gramen aspero simile, sed lævius. of BOx. 203. pl. 40. R 3. p 612. 4 Mr. Doody observed this pretty common about London.

64 Cockssoot Grass, R 1272. 6 S 184. 2 d 253. 2.

65 Gramen Dadylon latiore solio, CB 8. 4 pr. 19. 87 Th. 114 f 7 st. 222. 1 H 658. f 1. G 25. 2 Lob. 24 1 B 24. 2 O 13. 1 Ad. 4 fonst. 27. 2 P 1178. 1 B Ox. 184. 2 T 3. f opt. Found plentifully in the plough'd Fields about Elden in Suffolk, &c. 65 Small Sea Gockssoot. Dadylon marit. minus, R. S. 184. 1 d 253. 1 qr? Lugd. 421 f 2. CB 8 col. 1. 3 Th. 113. f B Ox. 184. 4 T 3. s. 2 f opt. Grows plentifully on the Sandy Shores between Penzance and Markerjeu in Cornwall. 66 Floating Cockssoot, or Flote Grass, R. 1281. 7 S 255. 4

66 Floating Cocksfoot, or Flote Grafs, R. 1281. 7 S 255. 4

Gramen aquaticum fluitans, multiplici spica, CB 3. 1 c 2. pt. 7 23 Th. 41. f. Tab. 216. 2. H 654. 1. G 13. 1. P 1275. 8 f Lob. 12. 2 B 14. 1 Ch. 187. 5 Jonst. 14. 1 B Ox. 183. 16

13. f Common in Ponds and slow Waters from May 11! Sept. 67 Common Rheed. R 1275. 1 S 185. 1 d 254. 1 B Ox.

13. f Common in Ponds and flow Waters from May till Sept.
67 Common Rheed, R 1275. I S 185. I d 254. I B Ox.
118. I R. f
Aundo vulgaris, CB 17. I pt. 39. 2 Th. 269. f Tab. 254. I
H 703. 2 G 32. I Lob 51. I B 70. I O 28. I Ad. 27. Dod 602. I Ch.
193. I John. 36. I P 1209. f 1. Well known on River fides, &c.
100 May till Autumn.
68. Wood Rheed, R 1282. I 2 f 185. 2 d 254. 2
Gramen Arundinaceum particula molli spadicea majus, CB 7. 3
pt. 8. 30 Th. 95. f Tab. 211. 2 H 648. 2 G 7. 2 Lob. 6. 2
B8. 2 Chabr. 186. 3 J 9. I Park 1181. f 3 opt. & 4. B Ox.
118. 2 T 8 f opt. In brambly Woods, as in that against the
biarded River plentifully from June till Autumn.
109 Small Rheed Grafs. Calamografiis minor glumis ruffis
4 windibus: gr? Gramen Arund. panicula molli spadicea minus,
CB 7. 4 Its Panicles some green, others brown and yellowish,
12 maller and more socie than the Wood Rheed. The first discovery of this Grass is owing to Mr. John Scampton a Curious

overy of this Grass is owing to Mr. John Scampton a Curious Borahiff, who sent it me from Leieestersbire.

70 Chassy Rheed, R 1280. 3 & 4 Var. Syn. 186. 3 d

bramen Arundin. acerosa gluma nostras, Park. 1373. f 2. opt. B 02. 203. 41 & 2 T 6. f 2. f opt. Gramen Arundinaceum spicatum (ex sent. D. Vaillant) GB 6. 2 pt. 8. 20 Th. 94. f Id. 212. 1 H 648 3 G 21. 12 f P 1181. 1 f 2. On the Banks of the Thames, and other Rivers, from Mid-May to Autumn

71 Jersey Chaffey Rheed. Gramen Arund. acerosa gluma seleianum, RS 239. d 255. Dr. William Sherard found this in the Ise of Jersey, on the Sea Coast over against Nor-

CAP. II. Frumenta & Gramina Paniculata.

CAP. II. Frumenta & Gramina Paniculata.

Paniculated Corn and Graffes.

SECT. IX. De Avenis & Gr. Avenaceis Oats and its Grafs.

72 White Oats. R 1253. I S 178. I d 246, I

Avena vulg. S. alba CB 23. I pt. 51. I Th. 469. Matth. 326.

1. Tab. 275. 2 H 791. I G 68. I Lob. 31. 2 B 40 2 O 19. 2

Ad. 9. Dod. 511. I J 75. I P 1134. I Lugd. 403. f Chabr.

176. 4 B Ox. 209. I T 7. f. It's fown towards the End of February and Beginning of March, and ripe in August.

73 Black Oats. R 1253. 2 f 179. 2 d 246 2 B Ox. 209. 3

Avena nigra, CB 23. 2 pt. 51. 2 Th. 472. 2

14 Blown Oats. Avena suscent fuse a fuse B Ox. 209.

This is a fuller Oat and commonly sown in Staffordshire.

75 Naked Oat. In Cornwal, Pillis and Pill-corn, R 1254. 3 S 179. 2 d 246. 3 Avena mida CB 23. 3 pt. 51. 3 Th. 47t. f. Tab. 276. 1 H. 791. 2 G 68 2 Lob. 32. 1 B 4t. 1 Dod. 511. 2 J 75. 2 P 1134 2 Chabr. 176. 5 B Ox. 209. 4 T. 7. f. opt. It's smaller than the White, with a closer and more upright

opt. It's imalier than the White, with a closer and more upright Tuft, and florter Pedicles; it easily cass its Husks.

76 Hairy Onts or Haver, R 1254, 4 S 170, 4 d 276, 4

Festuca utriculis lanugine flavescentibus, CB 10, 13 pl. 24, 106

Th. 149, f opt. 10b, 33, 2 B42, 2 O 20 3 Ad, 21, Chab, 176, 6

P 1148, 4 Lugd 406, f It's taller than Onts, with crooked Beards and hairy yellowish Husks. Grows amough Barly and other Corn, and ripens earlier.

other Corn, and ripens earlier.

77 Burnt Oats. Avena Ustilago, CB 24. 3 pt. 52. 3 Th. 476. fTab. 70 3 H. 684. f G 70. 3 Lob. 36. 2 B 45. f 3. Jonst. 77. 2 Found too often with the other Oats.

78 Dale's Silver Oat. Gramen Avenaceum Mont. spica simplici aristis recurvis, Ra, 1290. 11 S 184. 2 d 252. 2 B Ox. 213. 21 T 7. f 2 f opt. Its shining Spike 2 or 3 Inches long, with few alternate unbranched Glumes. Grows on hilly Ground in Flora Kent 850. in Essex, Kent, &c.
79 Doody's Silver Oat. Gramen Aven. hirsutum, panicula

purpurea-argenteo splendente, Ray 1909. S 192. 7 d 252. 3 & 262. 10 B Ox. 213.22 Differs from the last in growing taller and panicle branched. Found in the Pastures about the Earl of Cardi-

gan's House at Twittenham in Middlesex.

80 Bob-rooted Oat, R 128. 4 c 8. 1 S 191. 1 d 260. 1 B Ox.

214 38 T 7. fopt.

Gramen nodosum Avenacea panicula CB 2. 1 pt. 21. 97 pr. 3. f
2. Th. 18. f opt. Lob. 23. 2 B 23. 2 Ad. 2. G 22. 2 J 23. 2 f
Chabr. 180. 4 Lugd 429. f 1. Park. 1176. f 2 Grows in Corn and Hedges.

81 Merret's Dog Oat. Gramen Canin. Aventee panicula non nodosum Mer. Pin. B Ox. 214. 37 T 7. 12. fopt. Differs from the last in having a threddy Root and a longer

Tuft, Common in Heages and Paftures.

82 Bobart's Fine Oat. Gramen Avenae. sparsa panicula speci-osa, locustis minimis, B Ox. 215.41 R 3. p 611.4 Grows 2 Foot and half high, with a handsome Tust a Span long, in graffy Places.

83 Ray's Golden Oat. Gramen Aven. prat. elatius, panicula flavescente, locustis parvis, R 1284. 1 S 187. 2 d 256. 2 B Ox. 215. 42 T 7. 1 3. f opt. Frequent in Meadows about Mid-

84 Doody's great fandy Oat grass. Gramen paniculatum, lo-custis parvis purpuro-argenteis majus & perenne Doody, R. S. ed. 258. 12

Gramen nemorosum panie, albis capillaceo folio CB7.1 pr.14. 45 Th. 97. 1 B Ov. 200.9 T 7. f 3. f ult. In fandy Woods and Heaths, from June till Aucumn.

85 Kay's small sandy Oat grafs. Gramen paniculatum locufis parvis purpuro-argenteis aunuum, R 1286.14 S 189.11 d 258.

11 Gaz. Nat. Tab. 66. f 5. B Ox. 200. 11 T 5. f 3. f opt. In fandy barren Places, from Mid Ma till Jugust.

Jandy Darren Places, from Mid Ma till Zugust.

86 Doody's great Water Oat. Gramen Arund. aquat, panicula Avenacea, R 1909. S 186, 2d 254, pl. ult. B Ox. 215. 46

First found by Mr. Doody on the Banks of the Thames, between London and Chelsey.

87 Ray's nak d Meadow-Oat. Gramen paniculatum elatius, spicis longis muticis squamosis, R 1286. 10 S 188. 9 d 258. 9

B Ox. 183. 15 T 2 f 3. sqr? Park. 1146. 3 f Frequent in moss Meadows from May till Mowing.

88 Naked Wood Oat. R 1280. 4 S 102. 0 d 262. 12

88 Naked Wood Oat, R 1289. 48 192. 9 d 262. 12 Gramen Avenae. locustis rarioribus CB 19. 3 P 1151. f 3. Chab. 177. f 6 B Ox. 215. 49 T 7. f 2. f opt. In shady Woods and Hedges.

89 Naked Mountain Oat, R 1289. 3 Syn. ed. 262 13.

Gramen Mont. Avenac. locustis subris, CB 10. 2 pt. 10. 41 pt. 20.

I f 1 Th. 155. f Ch. 177 f 5. P 1151. f 5 B Ox. 215. 48 T 7. f
2. f Mr. Fitz Roberts tirst observed this, about Kendale in Westmoreland.

90 Ray's Naked Ost. Gramen Avenac. parvum procumbens, paniculis non aristatis, R 1288. I S 192. 8 d 262. 11 Pluk. 34. f I. B Ox. 177 6 T 1. f 3. f opt. On Putney Heath, and in di-

vers Pastures.

vers Pastures.

91 Common Drank, R 1289. 5 S 191 2 d 260. 2
Festuca Avenacea sterilis elatior, CB 9. 7 pt. 23. 104 Tb. 146. f
Tab. 234. 1 H 666. f 1 G 69. 1 Chab. 177. 1 Lob. 32. 2 B 41.
2 O 20. J 76.1 P 1148. 1 H Ox. 212. 11 T 7. s 2. f opt.
Frequent in Hedges and Field-sides, from May till August.
92 Small Drank, Ray 1289. 6 Var.
Festuca avenacea sterilis humilior, CB 10. 8 pt. 23. 105 Th. 148.
8 Found in dry Places in May, about a Span high.
93 Bobart's upright Drank. Festuca avenacea sterilis, spicis erettis, R 3. p 611. 3 S 237. d 261. 3 Pluk. 234. f qr. Ger.

Ger. 69. f. 2. Its lower Leaves very hairy. In the Hedges beyond Botley near Oxford.

Festuca Avenacea sterilis, pani-94 Jerfey upright Drank. B. Ox. 212. 13. T. 7. S. 2. f. opt. On fandy Sea Bores, as about Sand-down Castle, where Mr. James Sherard also found it. Spikes in June.

95 Doody's fine awned Drank. Gramen Avenaceum glabrum, panicula è spicis ratis strigosis composita, aristis tenuissimis, R. 1909. Syn. 192. 6 d. 262. 9 B. Ox. 214. 13. By Rivulets and

Ditches in divers Places.

96 Darnel Hedge-Drank. Gramen Avenaceum dumetorum spicatum, R. 1262. c. 6. 1. S. 181. d. 249. B. Ox. 214. 33. It spikes after Midsummer, in Wood-Hedges, and Thickets.

97 Ray's Hedge Drank. Gramen Avenac, dumet, panicula sparia, R. 1289, 8 S. 192.5, d. 261.8. B. Ox. 213, 26. It tusts in June and July, &c. in Thickets and Wood sides.

98 Bobart's hairy Hedge Drank. Gramen Avenacea dumetorum paniculatum majus hirsutum, B. Ox. 213, 27. T. 7. S. 2 Found with the last, but is more hairy and longer

. Capons - tail Grafs, Ray 1286, 15. S. 189, 12, d. 258. 13.

Gramen Murorum spica longissima, Jonst. c. 22. p. 29. f. 2. P. 1162, f. 8. B. Ox. 215. 43. T. 7. S. 3. f. opt. Lob. illust. 7. On Walls and dry barren Places from May to August.

100 Ray's Cocks tail. Gramen paniculatum Bromoides minus, paniculis aristatis unam partem spectantibus, Ray 1287. 16. S. 189. 13. d. 258. 14. Pluk, 33. f. 10. B, Ox. 215. 44. In

Stubble and dry Ground.

101. Merfey Drank. Festuca sterilis, humilima spica mam partem spectante. Its Spikes short and broad with larger Husks.

Found by Mr. Dale in Merfey-Ifle near Colchefter.

Poind by Mr. Date in Merjey-life near Colchester.

102. Common hoary Gross-Oat. Festuca Avenacea histuta, paniculis minus sparsis, Ray 1289. 6. S. 191. 3. d. 261 5. B. Ox. 213.18. T 7. On Walls, Field-sides, Meadows, &c. nothing more common from the End of April till August.

103. Smooth Meadow Gross Oat. Gr. Avenac. pratense, gluma tenuiore glabra, B. Ox. 213. 19. T. 7. s. opt. In Meadows with the last, but more rare.

Festuca Avenacea, spicis 104. Dale's smooth Grafs Oat. strigosioribus, glumis glabris compactis, R. 1909. S. 191. pl. ult. d. 261. B. Ox. 215. 47. Its Leaves hoary and Spikes mostly purplish.

mostly purplish.

105. Smooth German Gross Oat. Festuca Avenacea, spicis habitioribus, glumis glabris, R. 1289. 7. S. 192. 4. d. 261. 5. B. Ox. 212. 16. T. 7. f. Chabr. 177. f. 2. Grows in Corn.

106. Hoary German Gross-Oat. Festuca spicis habitiorilus, glumis inçanis. q1? CB. 9.1. pt. 23. 103. Th. 143. f. opt. Grows with the last, but more rare,

107. Sherard's tall Gross Oat. Festuca elatior, paniculis minus frastis location, though trivotis acidatis purpures splen.

nus sparsis, locustis chlongis, strigotis, aristatis, purpureis splendentibus, Doody, Ray Syn. ed. 201. Dr. Sherard found this amongst the Corn near Southampton. Spikes in June, &c. SECT. X. De Tremulis, Miliaceis &c. Quaking Grass, Miller's, &c. 108. Cow quakes, Quaking-Grass. Ladies bair, R. 1274. 3.

S. 185. d. 253. ult.

Gramen Tremulum majus, CB. 2.2. pt. 9.37. Th. 22. f. opt. Tab. 231. I. H. 665. 2. G. 80. 2. J. 86. 2. B. Ox. 203. 45. T. 9. 1. 2. f. opt. P. 1165. f. 2. var. Chabr. 184. 4. f. 1. q1? Cluf 218, f. 2. Well known in Meadows from May till Michaelmas.

109. Jerley Quaking-grafs. Gramen Tremulum minus, pa-

109. Jersey Quaking-grass. Gramen Tremulum minus, panicula ampla, locustis jarvis triangulis, Ray Syn. ed. 254. 2. B. Ox. 203.47. T. 6. s. 2. s. opt. Pluk. Amalth. 110. pl. 7. qi? Trag. 670. f. Dr. Sherard sirst found this in Jersey.
110. Sweet Water-Millet. Gramen Miliaceum aquaticum, R. 1281. 8. S. 187. d. 255. 1. & 325. Merretr. pin. B. Ox. 202. 29. CB. 1. 6. pr. 2. p. 1. Th. 6. p. 13. Lob. illustr. 10. qi? C.B. pr. 15. p. 7. B. Ox. 201.27. Frequent about London in slow or standing Waters from May to August.
111. Wood-Millet, R. 1282. 1. S. 187. d. 256. 2.
Gramen Islanding Waters standard sparsa, CB. 8. 1. pt. 7.

Gramen filvaticum panicula Miliacea sparsa, CB. 8. 1. pt. 7. 21. Th. 141. f. Tab. 205. 2. H. 639. 2: G. 6. 1. P. 1153. 1. Lob. 3. 2. B. 6. 1. Pod 561. f. 2. J. 6. 1. B. Ox. 197. 10. T. 5. S. 2. f. opt. In Woods from Mid May till Autumn.

112. Irish Miller. Gramen Montanum Miliaceum minus, radice repente, Ray Syn. ed. 256. B. Ox. 197. 12. descr. Dr. Sherard gathered this on the Mountains of Mourn in Ire-

land.

113. Great Water Millet, R. 1280, I. S. 186, I. d. 254. I. Milium aquat, majus, Nob. Gramen aquat, latifol. CB. 3. col.

1. pl. 4. pt. 1. 22. Ic. 7. Th. 40, f. Ger 6. f. 2. Tab. 216. Ic. 1. H. 653. f. 2. Lob. 4. Ic. 1. B. 6. f. 2. Jonff. 6. f. 2. P. 1274. f. 2. B. Ox. 201. 25. T. 6. f. 1. f. Very common by

1274. f. 2. B. Ox. 201. 25. T. 6. I. I. f. Very common by River Sides in June, July, &c.

114. Bent Millet, R. 1283. c. 7. S. 187. I. d. 256. 1.

Gramen Miliaceum fegetale majus Nob. Gramen fegetum altiff. paniculā fparsā, (B. 3. pt. 18. p. 7. Th. 34. f. 3. G. 5. f. I. Tab. 203. Ic. 2. H. 638. f. 2. P. 1158. f. 3. B. OX. 199. I. T. 5. f. 2. f. opt. Lob. 2. Ic. 2. B. 5. f. 2 G. 5. f. I. Chabr. 181. fig. mal. 5. I. B. 464. f. d. The Country-People call the Bents, and adoratheir Chimneys with it. Its Glumes are shining and without Beards. It sufis in May and June. in west

ning and without Beards. It tufts in May and June, in wet

Ruthy Pastures and amongst Corn.

115 Small red Bent Miller, R. Syn. 187. t. d. 256. t. Var.
Gr man Miliac. minus, pulchrè rubens.
Found in divers Forests in Westmorland.

110. Narrow Millet-Grass. Gramen. Miliac. fol. angustissimo.

Ito. Narrow Millet-Grals, Gramen. Miliae, fol. angustissimo. Its Leaves long and very narrow, the Glumes plain and not thick set. Tusts in June and July in shady Thickets, &c.

117. Narrow Fine Millet Grass. Gramen Miliae, angustisol. glumis perexiguis. Its Leaves shorter than the last, but narrow, the Panicles much spread with very small Glumes. Tusts in Juli in the Grove at the upper End of Greenwich-Park.

118. Great brown Meadow Millet-Grafs. Its Creat brown measure muer-Grais. Gramen Miliae, majus panicula speadicea, q1? B. pr. 12. p. 6. Its Tusts large and brown, composed of fine forked or gaping beardless Glumes. In dry hilly Pastures in June and July.

119. Great green Meadow Millet-Grais. Gramen Miliae. maj. panic. viridi. In Meadows by River-fides in June and

Fuly.

120. Small red Meadow Millet-Grais, R. 1286. 13. S. 189. 10. d, 258. 10.

Gramen Miliac. minus, panienla subente. qr? B. Ox. 199. 4. Common in Meadows in May and June.

121. Euddle's Wood Millet-Grass. Gramen Milias. sylvestr. glumis oblongis. First discovered by him in Bifbops-Wood,

Hampsted. Hampited.

122. Great brown Bearded Millet-Grass. Gramen Miliac.
majus, glumis aristatis spaciceis Nob. R. 1284. c. 9. pl. 2. S.
187. 3. d. 256. 3. B. Ox. 200. 17. T. 5. S. 3. f. CB. 3. 4. pt.
19. p. 7. Th. 35. f. G. 5. f. 2. Tab. 212. f. 2. H. 648. f. 4. Chabr.
181. f. 4.1. I.B. 461. f. Lob. 3. Ic. 1. B. 5. f. 3. Jonft. 5. f.
2. P. 1158. f. 3. Divers Authors have transposed the Synonyms of this Grass to the Bent-Millet; but Lobell's Figure gives its fine Beards, by which it is early distinguished from Gr. C. 114.

123. Great pale bearded Millet-Grass. Gramen Miliae, ma-jus, glumis aristatis pallidis. Grows with the left in moist Mea-

dows, and amongst Corn.

124 Soft Meadow Millet Grais, R. 1285. 7. S. 188. 6. d. 257. 6.

Gramen Miliae, prat. molle, Nob. B. Ox. 202. 34. T. 6. S. 1.

f. CB. 2. 1. pt. 17. p. 6. pr. 5. f. opr. 2. Th. 27. f. d. Chabr.
183. 3. J. B. 465. f. d. 3. Lugd. 425. f. 1. Fonf. l. 1.c.22.

pl. 6. it tufts in June, &c., in Meadows very common.

125. Soft bearded Corn-Millet, Ray 1285. 8. S. 188. 7. d.

125. Sott Dearded Corn-Datuer, Ray 1205. 8. D. 180. 7. U.
257. 7.

Gramen Miliac. ariflatum molle Nob. B.Ox. 202. 35, qr. 10b. pt. alt. 467. f.2. Chabr. 180. f.3. J. B. 457. f. Different from Gr. C. 124. in having a long creeping Root and short Beards. In Clayey and Corn-Fields. Tufts in June, &c.
126. Small, fost, Sea Millet-Grass, Ray H. pl. 1273. pl. 8.
Gramen Miliaceum marit. molle, Nob. CB. 5. p. 5. Th. 5. p.
74. f. Lob. pt. alt. 466. f. opt. Chabr. 182. f. 1. J. B. 463. f. 2. B. Ox. 202. 30. Lob. illust. 8. pl. 1. Lobel first discovered this on the Coast of Kent, and Mr. Buddle since him, in Suffolk. in Suffolk.

SECT. XI. Gramina Paniculata Pratorum, &c. 127. Great Meadow Grafs, R. 1285. 4. descr. opt. S. 188.

5. d. 275. 5.
Gramen Prat. panicul. majus, latiere folio. CB. 2. pt, 7. p. 4.
Th. 28. f. 2. Ger. 2. f. 1. Tab. 200. f. 1. H. 636. f. 2. Lob.
I. f. 1. B. 4. f. 1. O. 9. f. 1 Jonfl. 2. f. 1. P. 1156. f. 4. Very common in Meadows, Pastures, &c. 128. Great, narrow Meadow-Grass, R. 1285

Gramen prds. panic. majus, angustiore folio, CB. 2. pr. 10. p. 5. Th. 3. p. 29. B. Ox. 201. 19. T. 5. f. 3. f. opt. Its Panicles stender, with small Glumes. In Thickets, Pastures, &c. 129, Midding Meadow. Grass, R. 1285. 6. S. d. 257. I. B.

Ox. 201. 20. Gramen Prat. paniculatum medium, CB. 2 pt. 8. p. 4. Th. 5. p. 30. f. G. 2. f. 2 Tab. 231. Ic. 1. H. 637. f. 1. Lob. 1. Ic. 2. B. 4. f. 2. J. 2. f. 2. P. 1156. f. 2. Lob. illustr. r.

Leaves and upper Stalks smoother than 127. with Tufts is spread. In Passures and Hedges.
130. Flat-straw'd Meadow-Grafs. Gr. Prat. medium, culmo compresso. Mr. Buddle, that accurate Graminist, observed this on the Walls of Malden in Essex, and amongst the Corn in Sussell. It differs from the last in having statks with sirer Glumes, and Flowers later.

131 Common small Meadow-Gross.

Gramen Prat. minus, seu vulgatissimum, R. 1284. 3. descr. opt. S. 187. 4. d. 256. 4. B. Ox. 201. 21. T. 5. s. 3. s. opt. CB. 6. c. 2. p. 2. pt. 14. p. 6. Th. 31. f. Chabr. 182. f. 3. I. R. 465. f. I. G. 3. f. 1. & 2. J. 3. f. 2. P. 1156. f. 3. & 4. Inh. 207. f. 1. & 2. H. 639. S. 4. & 40. f. I. Well known and very common in Paths, Passures and moist Places, from Spring will Lutumn.

112. Common Paniel 1820 Care.

till Lutumn.

132. Common Panicled Sea-Grass. Gr. paniculatum marimum vulgatissimum, R. 1286. 12. S. 190. 17. d. 259. 18. Common in Salt Marlbes, tusts in June, &c.

133. Great panicled Sea Grass. Gr. panic. marit. majus, R. 1286. 12. var. S. 190.17. var. d. 259. 18. var. Its Stalks much thicker and taller than the last.

much thicker and taller than the last.

134. Lavander Grass, R. 1288 26. S. 190. 2. d. 260 21.

Gr. pratense spica Lavandula Merret. B. Ox. 201. 22. T. 5. s.

3. opt. CB. 7. p. 7. Th. 98. 7. Trag. 684. 2. Spikes in July in moist Meadows, Woods, &c.

135 Sea Haver-grass, R. 1286. xi. S. 190. 18. d. 259. 19.

Festiva marit, locustis lates. Found plentifully on the Sandy Shoresat Yumouth in June.

136 Ray's great Meadow Haver, R. 1285. 9 S. 188. 8. d.

257.8.

Festuca pratensis major Nob. Gr. sol. junceis brevibus majus rad. nigra, CB. 5. pr. 34. p xi. Th. 2. p. 73. B. Ox. 183. 14. Common in Meadows. Tusts in May and June.

137 Ray's small Meadow Haver, R. 1288. 23. S. ed. 260.

23. Fefluca prat. minor Nob. Gr. foliolis junceis brevibus minus, CB. 5, pt. 35. p. 9. Th. 73. f. 3, Bob. Ox. 182.13. T. 3. f. 1. f. opt. Pluk. 34. f. 2. Lob. 7. lc. 1. B. 9. f. 1. O. 9.2. Jouff. 4. f. 3. P. 1183 f. 4. G. 4. f. 3. Tab. 208. Ic. 1. H. 640. f. 3. l. B. 2. l. 18. p. 463. c. 68, fee. defer, Tufts in May, &c. in dry fandy bib Ground. high Ground.

138 Hard Wall-Grass, R. 1287. 17. S. 189. 14. d.259. 15. Gr. panicula multiplicis, CB. 7. p. 3. pr.xi. p.6. f. Th. 32. f. 7. P. 1157. f.5. B. Ox. 182 9. T. 2. f. 3. f. Barrel. 1212. Ic. 49. Very common on Brick Walls and dry Ground in June, &c.

136 Yellow Spring Grass, R. 1268. c. xi. pl. 1. S. 183 d.

252.
Gr. prat. spic? flavescente, CB. 3. col. 3. p. 3. pt. 9. p. 4. Th. 44 f.3. Chabr. 183. Ic. 1. l.B. 466. f. 1. Barrel. 1181. Ic. 124. f.1. B. Ox. 192. 25. T.4. f 2. & T. 7. f. 3. Appears amongst the first in the Sping in most Meadows.

140 Great Childing Grass. Gr. Proliferum majus fol. latione, Nob. B.Ox. 200. 12. T.5. f 3. f.ult. On Carnedg Libwellm and other Welch Mountains.

141 Small Childing Grass. Gr. Proliferum minus, capillacososio Neb. B. Ox. 200.13. R. 1270.10. S. 184. d. 252. On the Tops of Snowdon, Caderidis, &c. plentifully.

CAP. III. Cyperi, Cyperoidea, & Gram. Cyperoides.

Cyperus's, Cyperoides, and Cyper Grasses.

CAP. III. Cyperi, Cyperoidea, & Gram. Cyperoides.

Cyperus's, Cyperoides, and Cyper Graffes.

SECT. I. Cyperi. Cyperus's, Ray H. Plant. 1299.

142 Long fweet Cyperus, Ray 1299. I. S. 199. I. d. 270. I.

Cyperus odorotus rad. longå, five Cyperus offic. CB. 14. I. phyt.

36. p.5. Th. 216. f. I. Dod. 338. f. 2. Ger. 28. f. I. Tab. 656. f. I.

Hift, 341. f. I. Lob. Ic. 75 2. Ad. 37. Belg. 107. f. 2. obf. 40. f.

I. fonfi 30. c. 24. f. I. Park. 140. f. 3. & 4. Chabr. 149. f. 2. I.

B. 2. I. 18. p. 501. f. I. Or. B. 237. 13. T. 10. f. 2. f. Found in

Purbeck Isle near a Chapel on the Side that looks towards Portland. Island. Flourishes about September.

143 Long Marsh Cyperus, R. 1296. I. S. 197. I. d. 270. 2.

Gr. Gperoides paniculä sparså majus, CB. 6.7. phyt. 65. p. 15.

Th. 86. fig. bonå. Lob. Ic. 20. I. B. 21. f. I. Chabr. 189. f. 3. I.

B. 2. I. 18. p. 495. f. fonst. 30. c. 24. f. 2. Park. 1266. f. 6. Ox.

B. 238 25, T. xi. f. 2. f. Inst. 527. These Panicles spread more than the next, and stand 3, 4, or 5 on a Pedicle. Flourishes in July and August in most Fens, and in the Thames plentifully.

144 Round Marsh Cyperus, R.1300.6. S.200.2. d.270.3. Cyperus rotundus inodorus Germanicus, CB. 14. 3 Prodr. 24. fig. Th. 215 f. 3 Park. 1264 f. 4 Its Panicles are closer set; but its knobby Roots peculiarly distinguish it from the last. Very common in August and September in most Salt Marshes.

Cyperus longue inodorus Germanicus, CB, 14. 4. phyt. 36. p.

7. Theat. 221. f. 3. Lob. Ic. 76. r. B too. f r. Jonst. 29. f. 3. c. 21. Park. 1294. f. 1. Chabr. 195. f. 2. I. B. 2. I. 18 p. 564 f. 1. Ox. B. 237. 24 T. xi. f. 3. f. 24. & prox. Boc. 72. f. opt. Inst. 528. This partaking both of the Rush and Coperus; it's easily known from all others. Flourishes in May and June by the Sea Side

in Cornwal, and on Hinton Moor plentifully.
146 Millet Cyperus, Ray 1301.10. Syn. 200.4.d. 271.5. Gramen Cyperius, Ray 1301, 10. Syn. 200, 4.0.271. 5.

Gramen Cyperiodes Miliaceum, CB. 6.13. phyt. 64. p. 15. Th. 90

f. 12. opt. Lugd. 988. f. & 693. f. 1 Chabr., 195. 3. 1. B. 2. l. 18. p.
504. f. 2. Park. 1171. f. 2. Adv. 38. f. Lob. b. 79 f. 1. B. 109. 1. Jonst.
30. f. 5. c. 21. Ox. B. 237. 15. T. xi. d. 2. f. opt. 15. This is a frecious Cyperus, and easily known by its Figures. Grows in fome
warry Places as about Hamphered and elsewhere in Fune watry Places as about Hampstead and elsewhere, in June, &c.

147 Fonflons's small white Cyperus, Pluk. Tab. 34. f. xi. Gramen Cyperoid, palufire leucanthemum, R. 1295.13. S. 196. xi. d. 266. 15. Gramen Cyp. albis glumis, Merret. 51. Gramen Junceum leucanthemum, Jonfl. l. 1. c.22 pl.7.p. descr. This is a kind of Gramen Luzule Chabr. 188. f. 3. Grows plentifully in a large low Bog between Wickam and Croydon. Flourithmain Angel. rithing in August.

148 Bobart's small brown Cyperus.

Cyperus minor angustifol, palnitris, capitulis fuscis paleaceis, Bob. Ox. 239. 49. T.xi. ter. 3. f. ult. A Variety of the last; I found this plent fully in a By between Southampton and Lim-

ington in August.

SECT. II. Cyperoides Julifera, Ray.

Cyper-jules, or Cyperoides hearing Jules.

149 Cyperoid with thost pendulus Spikes, Ray 1294, 8. Syn.

195.7. ed.265.10.

Cramen Cyperoides, spica pendula herviore, CB.6.5. phyt.63. p.

14. Theatr.85. f. 5. Dod. 339. f Lob. Ic. 76. 2. B. 109. f. 2. Jonft. 29. f.

2. c. 21. P. 1266. f 4. Chabr. 189. 1. f. 6. I. B. 3 1-18 p. 496. f. 3. Ux.

B. 242.5. T. 12. S. 1 f. E. B. 421. Inft. 529. pl 4. Differs from the next in having thost comole Spikes, with one Jule, amongst four or five Spikes. Found in flow running Ditches, in June, &c.

150 Cyperoid with long hanging Spikes, R 1294.6. Syn. 195. 8. ed. 265. xi. Gram. Cyp. Bica pendula longiore & angustiore CB.6.6. prodr. 42. p.13 Th.85 6 Park. 1267.5. Barell. 1255. Ic. 45. Ox. B. 242. 4. T.12. S.1. f. 4. opt. E.B. 421. Inst. 529. pl.5. This is a very tall Grafs, with generally 5 long Spikes hanging at Distances. Found in wet Ditches, in June and July.

151 Ry's Cyperoid with upright Spikes.

Gramen Cyp. polyflachion majus, spicis teretibus eredis, Ray 1294.6. S. 195.6. d. 265.8. Ox. B. 242.8 T 12. ser. 2. f. opt. Its Stalks bluntly square with 3 or 4 upright Spikes, terminating generally with 3 Jules. Grows in thallow Pools.

152 Bohart's Cyperoid with yellowish Wheat ears.

Gramen Cyp. majus præcox, spicis turgidis teretibus flavescentibus, Bob. Ox. 242.6.T. 12 S. 1. f. 6. opt. These Spikes are thicker than the last, resembling a naked Wheat-ear. By Rivers in April. April.

April.

153 Ray's Hoary Cyperoid, Pluk. Tab. 34 f.6. opt.

Gr. Cyperoides polystachion lanuginosum. Ray. 1294. 4. S. 194. 4. d. 265. 6. Ox. B. 243. 10. Its lower Spikes much distant from the upper, the whole Plant somewhat boary. It Flourishes in May and June in moist Meadows.

154 Dr. Lister's Cyperoid, Ox. B. 243. 18 T. 12. S. 1. f. Gr. Cyp. Spicis parvis longistime distantibus, Ray 1295. 10. S. 195 9. d. 266, 12. A tall slender Grass composed of 3 or 4 short Spikes, the lowest much distant from the rest and attended with a longer Leaf. Flourishes about June in a Bog between Wickham and Croyden, near the Viola palustris.

155 Dale's long-spiked Cyperoid.

155 Dale's long-spiked Cyperoid, Gr. C.p. polystachion majusculum latifol. spicis multis longis strigosis, Ray Syn. ed. 265, 9. This is of kind with the next, but very much bigger. First discovered by Mr. Sam. Dale, in a Lane at Black-Notley in the Road to London. It flourishes in Ture. in June.

156 Parkinson's slender-ear'd Cyperoid. Gr. Cyp. sylvarum tenuits spicatum, Park. 1171 f.3 Lob. illustr. 60. Ray 1295. xi. S. 196. 10, &. 266. 14. Ox. B. 243. 9. T. 12. s. 2. f. opt. Very Common in most Woods, where it may be easily distinguish'd by its slender Spikes, which appear in

157 Great broad-leaved Spring Cyperoid R. 1293. 1. S. 194. 1. d. 264. 1. Gr. Cyp. latifol. spica ruffa, sive caule triangulo, CB. 6 1. phytop. 1258 p. 13 Th. 83 f. 1. Park. 1266. f. 1. Jonst. 12. f. 1. Lob. Ic. xi. f. 1. B. 13. f. 1. Tab. 218. Ic. 2. Hift. 655. f. 3. Ger. 29. f. 4. Chabr. 189. f. 1. I. B. 21. l. 18. 9. 494. f. 3. Ox. B. 242. 1. T. 12. f. 1, f. Inst. 529. pl. 1. Bastl. Cat. 7. Fet. 2. 363. We have a grass resembling these Figures, very common by Brooks and River sides, Flourishing in May.

158 Great, narrow-leav'd Spring Cyperoid, Rai 12.932 S.
1042 d. 2042 Gr. Cyp. angustifol. fpicis longis erectis,
CB. 64 pbyt 58 p. 14 Th. 84 f. 4 Lob. Ic. xi f. 2 B. 13
f. 2 Chabr. 189 t. 1 var. I B. 2 l. 18 p. 494 f. 2 Ox. B.
242 3 T. 12 f. 1 f. Inft. 529 pl. 10 The Grafs resembles ling these Figures is smaller then the last tho' near as tall, with narrow Leaves. Flourishes by Water-Sides, in May or fooner.

150 Parkinson's lesser narrow-leav'd Cyperoid.

Gramen Cyp. minus angustifolium, Parks 1266 f. 3.

Mr. Ja. Sherard, first observed this in a Pond near Eltham in Kene, about the End of May; and Mr. Rand, in the Ditches at the King's Arms against White-hall.

160 Small upright Clove-leaved Cyperoid.

Gramen Cyp. fol. Caryophylleis, spicis erectis sessibus, è seminitus confertis compesitis, Ray syn. ed. 2 p. 264 4.

Its Spikes on short or no Footsfalks. Very common in May

and June in moist Meadows.

161 Bobart's Small Cyperoid with Short Heads. Gramen Caryophylleum, angustissimis foliis, spicis fessibus brevioribus eredis non compactis Ox. B. 243 12. This has commonly but two fhort fessil Heads with one Jule at top. Grows Ly Wood fides and moist Meadows.
162 Small pendulous Clove leaved Cyperoid.

Gramen Cyp fol. Caryophylleis, spicis oblongis è pediculis lon-giorebus pendulis. Ray Syn. cd 2 p. 264 5. Differs from the last in having its Stalks more triangular, its Spikes on long

Footflalks, and pendulons when ripe. It Flourishes in May and June in wet Meadows.

163 Roy's Round-grained Bog-Cyperoid.

Gromen Cyp. fol. Caryor hyllers, spicis è varioritus & tumidioritus granis confertis, R. Syn. 194, 3 Var. ed. 264 3 Pluk. 91 f. 3 Ox. B. 243 13. On Chistleburst and other Bogs, in July. Its swelled Grains and Place, differ it from the foregoing.

164 Ray's Vernal Cyperoid.

Gramen Cyperoides vernum minimum, Ray 1294 7 S. 169
12 ed. 266 16 qr? CB. 3 6 phyt. 52 p. 13 Lob. Ic. 10 f.
2 & 3. This spikes early in the Spring, viz. about the Middle of April.

165 Doody's foft-leaved Hampflead Cyperoid.

Gramen Cyp. Bicis brevibus congestis, fol. mollibus, Ray 1910 Syn. 196 14 d. 267 18 Pluk. 91 f. 8 Ox. B. 245 33. It has 2, 3, or 4 small close roundish Heads, with one Jule. Found on Hamfftead Heath, about June.

166 Vernon's small Headed Cyperoid, on long Stalks.

Gramen Cyp. spicis habitioribus, pediculis longinsculis, R. Syn. ed. p.267 18 var. Differs from the last, chiefly in having Pedicles to its Heads.

167 Doody's Croydon bog Cyperoid. Gramen Cyp, una alterave spica rotunda præter cussam, R. Syn. ed. 345 Its Leaves short and soft, the Heads close, round and swelled; each attended with a Leaf. Grows about a Foot high in Croyden-Bogs, with the Viola palustris in July.

168 Ray's yellowish Short-headed Cypercia.

Gramen Cyp. polystachion flavicans, spicis brevibus prope summiratery within R 120d 5 Syn 105 5 ed 265 7. This has

mitatem canlis, R. 1294 5 Syn. 195 5 ed. 265 7. This has commonly but 2, rarely 3 Heads, on short Footstalks near its Jule; its Grains are not so pointed as the next; Heads about

169 Tall Hedgehog Cyperoid. Cyperoides echinatum majus. This has rarely more than 2 fessil or close sharp-grained Heads, near the Top, the low r bearded, with a long Leaf. Grows plentifully in a Meadow with the Cirfum Anglicum near Cambridge, where I got it in company with Mr. Ja. Skerard, about the mid dle of June, 1715.

170 Vernon's diffant-headed Hedgebog Cyperoid.

Cyferoides echinata, capite longe dinante, Nob. Ray Syn. ed. 267 plat. Its Herds very like the last, but the lower one stands on

a Pedicle 6 Inches below the other.

171 Small Marsh Hedgebog Cyperoid, R. 1296 15 S. 196 13 ed. 267 17 Ox. B. 243 19 T. 12 f.2 f. Gramen paluftre echinatum, Lob. Ic. 15 1 B. 16.1. Jonft. 17 f. 1. Our Grafs very well agrees with these Figures, being in all its Parts lets than the 2 last. Very common in July, in Bogs and moist Mea-

172 Flea Cyperoid. C. peroides pulicare, Nob. 172 Flea Cyperoid. C. peroides pulicate, Nob.
Ray 1208 12 Syn. 199 12 ed. 269 12 Ox. B. 244 21. T. 12 fer. 3
f. opt. Puk. 34 f. 10. Its Flea-like Grains hanging downward
differ it from all others. Found in Moorish and Boggy Places
from Mid-my to July End.
173 Pobarts Jule Cyperoid Ox. B 244 22 T. 12 f. 3 f. opt.
Gramen Cyperoides Speia fi implicite coffa, Ray Syn. ed. 247 20
Its Leafe very narrow and fhort, with a flender half inch brow-

nish chaffey Spike, like the Club-rush. It grows about half a Foot high in moorish spongy Soil, about Hockley near Oxford. SECT. III. Gramen Cyperoides spicis tantum Seminiferis.

Cypergraftes without Jules.

Cypergrafies without Jules.

174 Panick Cypergrafs with divided Spikes.

Gramen Cyp. paluftre elatius, spica longiore laxa, Ray 1296;2.

Syn. 197 2 d. 267 1 Ox. B. 244 23 T. 12 fer. 2 f. opt. A very tall Grafs, with divided Spikes; very like when ripe, that of Gramen Panicenm. spic l divisa, CE. without awns. Grows in rotten Bogs and watry Places in June.

175 Panick Cypergrafs with loofe flender Spikes. Gramen Cyp. palustre minus, spices gracilioribus, R. 1296.2, var. Syn. 197.2. var. ed. 268. Lesser than the last, and grows not in so thick Tuss, its Spikes stenderer and thinner sat. Found with

176 Cluster-headed Meadow Cypergrafs.

Gramen Cyp. ex mome Ballon fimile, spica totali è p'uribus spicis composita, R. 1297 5 Syn. 1985 ed. 368 4. Ox. B. 245 34. This has 3 or 4 close rough Heads, ending in a longish spike, the lower Part of which is chastey like a Jule. Found in June in Marshy Meadows.

177 Cluster-headed Sea Cypergrafs.
Gramen Cyp. ex monte Ballon fimile humilius, in maritimis &

Gramen Cyp. ex monte Ballon fimile humilius, in maritimis & arenosis naicens, R.1297 6 Syn. 1986 ed. 268 5 Ox. B.244 30. Its fpike large for the smalness of the Grass. Growing short, and low on the sandy Sea Shores in June.

178 Round Cluster beaded Cypergrass.

Gramen Cyp. spica è pluribus spicis brevibus mollibus composita. R.1296 3 Syn. 1973 ed. 268 2 qr? Gramen Cyp. palustre majus, spica divisi, CB 6 10 phyt. 60 p. 14 Th. 87 f. 10 Lob. Ic. 19 f. 2 B. 20 sig. 2 Chabr. 190 f. 1 Ox. B. 244 29 T. 12 1. 3 f. opt. These Figures (except the last) would better resemble our Grass, had they omitted its bearded Leaf which this never has. Grows pretty common in met Passures.

179 Plucknet's stat Cluster-headed Cypergrass.

Gramen Cyp. spica simplici compressa disticha, Ray 1910 pl. 2 Vol. 3 p.625 20 Syn.199 13 ed.269 13 Pluk. Tab 34f. 4 Ox. B. 245 35. Like the last, but not so tall, and flatter spiked. Found on Chistlehus stand other Bogs in June and July.

180 Ray's 10ugh Cluster-headed Cypergrass.

Gramen Cyp. palustre elegans, spica composita assertore, Ray 1907 Syn. 163 2 cd. 268 6 Plub 24 c On Ray

Gramen Cyp. palustre elegans, fpica composite asperiore, Ray 1297 7 Syn. 138 7 ed. 208 6 Pluk 34 4 Ox. B. 245 31. Its Leaf very narrow, with a naked Stalk a Foot high the Spike composid of 5 or 6 short pale Heads, without any Leaf or Beard. It grows in moilt graffy Places.

181 Greater Prickly headed Cypergrafs, R. 1297 8 S. 198 8 ed.

Gramen Cyp. palustre majus, spica compacti, CB.68 phyt. 59 p. 14 Th.87 f. 8 Tab. 220 Ic 1 Hist. 656 f.3 Ger. 19 f.5 Jond. 21 f.5 Park. 1266 f. 7 Lob. lc. 19 f.1 B.20 f.1 Chabr. 190 f.4 l B.2 l. 18 p. 497 f 1 Ox. B. 244 24 T. 12 s. 3 f. opt. Mr. Bobart's single Spike at the Bottom of his Icon, very well represents his Grass.

which other Authors have made too fhort and fingle. It spikes in M1y and June, near Rivers and watry Ditches.

182 Lesser Prickly spiked Cypergrass.

Gramen Cyperoides spicatum minus, Ray 1279 9 Synt. 198 9 ed.
269 9 qr:Gramen Cyp spicis minioribus minus; compactis, CB.
9 pr. 6 43 p. 13 Th. 89 9 Park. 1268 9 Ox. B. 244 25. The Spikes of this are bearded at each Division with a sine Thread, but smaller and slendarer than the last travers in mois Mandare. smaller and slenderer than the last. It grows in moist Meadows

finaller and itenderer than the last. It grows in mon. measure and Passures in May and June.

183 Small Prickly spiked Cypergras.

Gramen Cyp. spicatum minimum, spica divusta aculata, R. 1207 xi Syn.199 xi ed. 269 xi qr? CB. 7 4 phyt. 83 p. 18 Th. 100 f. 4 Chabr. 101 f. 4 1 B. 2 p. 510 f. 1 Tab. 227 Ic. 1 Hist. 662 f. 1 Ger. 20 f. 9 Ox. B. 244 26 T. 12 s. 3 f. The Tops of this Grass are composed of 2 or 3 small prickly Heads set pretty near together, as all the Figures represent. It spikes in June, in rotten Bogs and Shade Woods. and shady Woods.

184 Ray's Prickly-Spiked Cypergrass with a flender Ear. Gramen Cyp. spicatum minus, spica longa divulsa seu interrupta,

Gramen Cyp. spicatum minus, fpica longa divulsa feu interrupta, Ray 1297 10 Syn. 199 10 ed. 269 10 Ox. B. 244 28. The three lowest of these Heads stand at some Distance, but the rest next the Top closer. It grows in moist Pastures.

185 Soft, brown Wood grass.

Gramen nemorosum, spici rusescente molli, CB. 7 2 phyt. 82 p. 18
Th. 99 f. 3 Park 1183 f. 2 Tab. 226 sc. 1 Hist. 661 f. 1 Ger. 8 f. 2
Chabr. 191 f. 3 I B. 2 p. 509 f. 2 Pet. 2 133 & 380 Ox. B. 244 27
T12 f. 3 f. Its Spikes are soft and reddish. Mr. Bobare has observed this in moist, shady, sandy Woods.

186 Ray's narrow Cypergrass with short selfil Spikes.

186 Ray's narrow Cypergrafs with thort fessil Spikes.

Gramen Cyp. anguttifol spicis parvis sessilibus in soliorum alis,
Ray 1295 14 Syn. 195 15 ed 267 19 Pluk. 34 f. 3 Ox. B. 243 17
T. 12 s. 2. f. opt. Each thort Spike is thinly set in the Bosom

of slong bearded Leaf. Spikes in May and June, in moist and hady Direhes.

187 Buddle's large Cypergrass with short sessil Spikes.

Gramen Cyp. angustif, majus, spicis lessilibus solverum alis. This is is greet than the last in all its Parts. Observed by Mr. Buddle in Sufolk.

188 Pluknet's Cypergrass with double Bur-heads.

Gramen Cyp. minimum. Ranunculi capitalis hinis tarnism.

Gramen Gyp. minimum, Ranunculi capitulis binis ternifve, Or. R. 245 37 Pluk. 91 f. 8 Ray 3 p, 626 22. This much relembles the next, but inflead of one, has two or three Heds.

189 Bobar:'s Cypergrafs with a round Bur-head. Granen Cyp. fpica echinata simplici, Mer. pin. 52 Found frequently in the Spring, in the Bogs on the West Side of Ox-

100 Dr. Sherard's Cypergrafs with a long Bur-head. Gramen Cyp. ininus, Ranunculi capitulo longiore, Ray 3 p.626
23 Syn. ed 270 15. This is a larger Variety of the lift, with
Spikes or longer Heads. Found by Dr. Sherard in the fpungy
logs, in the County of Down, in Ireland.
CAP.IV. Junci & Gr. Juncea. Rushes and its Grasses.
SECT. I. Junci acuti & laves. Sharp and foft Pushes.
191 Large Pficking Sea Rush, Ray 1302 c.3 pl. 1 Syn. 200 1.

Juneus acutus, capitulis Sorghi CB. xi 1 prodr. 21 f. opt. Th. 173 f.1 Park. 1193 f.4. Chabr. 196 f.4 IB.2 r. 520 f. 3 Ox. B. 232 15 T. 10 fer. 2 f. This is a tall thick Ruß with large Sed Veffels, its Point very pungent. On fandy Sea Shores in

192 Paikinson's Sea Rush, Ray 1303 2 Syn. 201 2 ed,272 2. Juncus acutus marit. Anglicus, Park. 1193 f.7 opt. Ox. B.232 14 T.10 f. I f. opt. A tall hard Rush with childing Panicles. Commen in Sea-Marshes about Midsummer till Autumn.

Common in Sea-Marshes about Midjummer till Autumn.

193 Common Hard Rush, Ray 1303 5 Syn.201 5 ed. 272 5.

Juneus acus us panicula sparsa, CB. xi 4 phyt. 28 1 Th. 176 4

Dod.605 f.3 Lob. Ic.85 1 B. 119 3 Tab. 249 Ic.2 Hist. 695 f.2

Ger.31 f.4 Jonsl. 35 f.4 Park. 1193 f.1 Chabr- 196 f.5 1 B. 2 p.

521 f.1 Ox. B.232 13 T.10 f.1. f.opt. A tall slender Rush with little Pith and a loose Panicle 6 Inches below the Tip. Common in Paths and moist Pastures in June and July &c.

194 Small Westmorland Hard Rush, Fluk.40 f.8. Juncus parvus, calamo supra paniculam compastam longiùs poducto, Ray 1305 17 Syn. 202 xi ed. 273 xi. It's a stender sub, not o Inches high. Mr. Newton discovered this near emblesse in West moreland.

195 Sea Rush with round Heads, Ray 1303 2 Syn. 201 2 ed.

Juneus acutus marit. capitulis rotundis, CB. xi 2 Th. 174 f 2 Park 1194 f. 5 Lugd. 987 f. 1 Pluk. 40 f. 4 Ox. B. 232 17 T. 10 f. 2 f opt. The Reverend Mr. Stephens first found this in Devonfopt. The Reverend Mr. Stephens first found bie; it also grows in Somer fet and Hantshire.

196 Pluknet's Childing-Rush with round Heads. Juneus minor acutus marit. prolifer seu ex uno pediculo multa, Ruk. 40 f. 5 Ray 3 p. 629 3. A childing Variety of the

197 Common Bull rush, Ray 1304 7 Syn. 201 7 ed. 273 7.

June maximus f. Scirpus major, CE. 12 2 phyt. 29 5 Th. 178 f.

2 Tab. 249 Ic. t Hist. 696 f. 1 Ger. 30 f. 2 Lob Ic. 85 f. 2 B. 120 f 1

Adv. 44 Jonst. 35 f. 3 Paik. 1191 f. 1 Chabr. 197 f. 2 I B 2 p. 522 f.

2 Ox. B. 231 1 T 10 f. 1 f. opt. Its Use in matting Chairs, Baskets, &c. makes it well known, It heads in July and August, its most River plannishing.

in most Rivers plentifully.

198 Lesser Bull-rush Ray 1304 Syn. ed.273.

Juneus ag medius, CB. 12 3 phyt. 29 6 Th. 181 f. 3 Tab. 250 Ic. 1

Hist. 696 f. 1 Ger 3 1 f. 3 1 B-2 l. 18 c. 240 p. 524 f. v. Ox. B.

231 2 T. 10 f. 1 f. Somewhat lesser than the last. Mr. Dale has observed is in the Sea-diches at Bricklesey and Mersey Islands.

Black of its a grown in Woods.

lt's alfo fild to grow in Woods.

Juncus aq. major carinatus. Differs from the two last, in having a Sulca or Furrow on its Stalk. This Mr. Dood, has observed about Limehouse: I have also found it in his Company near the Thames, on this Side Rattersey Meadows.

200 Triangular Thames Built rush.

Juneus adultis marit. caule triquetro maximo molli procedior nostras, Pluk 40 f. 2 Ox. B. 232 21 Ray Syn. 2013 ed. 272 3. This is much larger and thicker than the next, with rolliferous Heads. Very common in the Thames about Westminson, in July and August.

201 Triangular Tersey Bull-rush, Ray 1303 3
Juneus acutus mari. caule triangulo, CB. xi 3 pr. 22 f. 1 opt. Th. 75 f.3 Park, 1194 6 Ox. B. 232 20 T. 10 s. 2 f.

202 Pluknet's Triangular Clufter headed Bull 1 who Funcus acums marit. caule triquetro molli, fpic glomerata in flylo fessili, Pluk. 40 f.3 Ox. B.232 22. This has our queles Heads at the Top, without any Stalk above. 203 Common tufted, loft or Candle-rulb.

Juneus la vis vulg paniculă spustă nostras, Ray 1304 8 Syn, 202 8 ed. 273.8 qt? Juneus levis paniculă sparsă mijor. CB, 12 4 phyr. 28 t Th. 222 f. 40x, B 2314 T. 1 f. 1 f. opt. They make Candles of the Pub of this, whose Tufts are spread more than N. 205. It grows in Ponds, and by Water-sides, in August and Sept. and Sept.

Juncus levis siter, CB. 12 5 Th. 1825 Ox. B. 231 5 This is t bicker and florier than the lath, and of the same Use.

205 Clofe beaded foft Rush. 205 Close-headed lost Rush.

Juneus lavis vulg paniculà compactiore, Ray 1304 9 Svn. 202 9

cd. 273 9 cm? Juneus lavis, panicula non iparfa, CB. 12 7 phyt.

28.2 Th 182 7 Chabr. 196 f 3 I B. 2 f. 520 f. 2 Ox. B. 231 7 I. 10

f. 1 f. cpt. It grows in Woods and most Passures, in July and Lugust,

200 Black-headed Bog rush.

Lugust Lavis names. Arrival delegant in invisente C. Cont. Box.

Juneus lavis unnor, panieula glomerata nigricante, C. Caut. Ray 1305 16 Syn. 202 10 ed. 273 10 Ox B 233 24 T. 10 f. 3 f. opt. I have gathered this in the Bogs, both about Oxford and Cambridge.

bridze in June and July.

207 Dwarf Rult, R.y 1305 14 Syn. 203 16 ed. 274 16

Juncesus, A.v. 4441. ult. inutilis, CB. 12 9 five Chamelchenos

prodr. 22 3 phyt. 3) x1 Th. 183 9 Park 1192 7 Ox. 8. 232 23 T. 10 f. 2 f.opt. It grows in moift gravely Heaths, &c. in June and

SECT. II. Junei Clavati. Club-rushes. 208 Great, tall, Thames Club rush.

Juneus Clavatus major & altior. This I have observed about Poterborough-House and the seven Elms, on the other fide of the Thames.

200 Common Club Rush, Ray 1305 18 Syn. 202 12 ed. 273 12, Juneus capitulis Equifeti finjor CP, 12 1 phyt. 29 7 II. 186 f. s. Lob Ic 86 1 B, 120 2 O. 8 44 1 Lngd 686 f. P. 11 1. 1196 f. 1 Chahr. 197 f. 6 I B. 2 p. 523 f 3. Very common ly River, and Pond Sides, in May, &:

210 Smill Heath Club-rush, Pluk. 40f. 6. Juneus parvus palufiris, cum parvis capi ulis Equifet i, R. y 1306 22 Syn. 203 15 ed. 27 + 15 qt? Juneus mont. parvus, cum parvis c pitulis luteis, Chair. 197 f. 5 I B. 2 l. 18 f. 523 f. 2 Ox. B. 233 35 T. 10 fer 3 f. On Putney and other moint Heaths in May, 211 Lhwyd's Mountain Club rush.

Juncello accedens Graminifolia plantula capítulis Armeria pro-híeræ Ray Syn. ed. 275 pt. 1 Eob. Ox. 245 40 T. 12 f. 1 f. opt. Its Leaf two menes long and very flender, with a roundish feal; Head, Found on the Welf Mountains and in rotten Ecgs. 212 Plot s leaft Club ruft, Pluk. 40 f. 7.

Juncellus omnium minimus capitulis Equifeti, Plot. H. Ox. 145 5 Tab. 9 f. 3 Ray 1306 20 Syn 202 14 ed. 274 14 Ox. B 234 37 T. 10 f. 3 f. In mont Ditches on Binfey Common, Oxfordshire, and elle where,

213 Floting Club-rush, Ray 1305 19 and 1310 19 Syn. 202 14 and 235 7ed. 274 13 and 276 7 Pluk. 35 f.1 Ox. B 233 36 T 10 fer z f. opt. Juneus capitalis Equiseti minor & finitans, Ct. 12 2 prodr. 23 6 Th. 87 2 Puk. 1196 3. In Ponds on Wandfworth Common, near Putney, and ellewhere in July.

SECT III. Grinina Juncea, Rush graffes. 214 Moss rush or Goose-corn. 214 Mols rush or Goole-corn.

Juneus montanus paiustris, Ray 1303 6 S. 201 6 ed 272 6 Gr. Junceum solitis & pica Junci, CB. 5 so phyt. 70 p. 16 Th. 78 f. 10 Tab.
224 Ic. 1 Hit. 64 6 f. 3 Ger. 18 f. 4 Lob. 18 f. 1 B. 19 f. 1 Jonst.
21 f. 4 Vark. 1103 fig. 2. This well resembles these Figures.

Frequent on Turney Heath and such like Places. It stourishes from Midsummer till Winter.

215 Round headed Rush grass. Gramen Junceum maritimum vel palustre cum pericarpiis totun4 dis, Ray 1307 6 Syn 204 5 ed. 276 5. Funcus parvus cum pericarpiis rotundis, I B. 21.18 c. 232 p. 552 fig. * & detc. Chabr. 191 lc. 5 fecund. Fig. 97 Gramen Juceum polytachion, CB. 5 7 phyt. 74 p. 16 Th. 75 f. 7 Tab. 223 lc. 2 Hiti. 650 f. 2 Ger. x: f. 2 qt? Park. 1190 f. 7 * N.B. The Synonyms in Mr. R. y & Histor., are mixt with the last The Rush here meant, grows with us, both in Sea Marshes and moit Meadows...

216 Long leaved Rush grass with a white Root.

Gramen foliolis juncers oblong is, rad. albaCB.5 t phyt. 34p. o
Th.73 f. 1 * Gramen Junceum Dileon. Lugd 425 f.2 Park. 1189
f 3 Ray 1309 14 Syn.ed. 277 xi. Mr. Bobart has olderv'd this
at the Head of the Bogs at Wooton Heath three Miles from Oxfor !.

217 Toadgrafs, Ray 1309 13 Syn 204 6 ed. 276 6.
Gramen nemorofilm eals culis paleaceis, CB. 7 5 phyt. 14 p. 18 Th.
100 f. 5 Tab. 225 Ic. 2 Hist. 660 f. 2 Ger. 4 f. 4 Park. 1190 f. 8 1 ob.
Ic. 18 f. 2 B. 19 f. 2 Jonst. 4 f. 4 Ox. B. 228 14 T. 9 f. 2 f. Very
common in wer and muddy Flaces in May, June, &c.
218 Ray's long: threaded Toadgrafs, Fluk 35 f. 2.

Gramen Junceum five Holofteum minimum paluftre, capitulis quatnor longissimis, staminibus donatis, Ray 1319 185yn. 105 8 ed. 276 8 Ox. B. 230 30 T. 9 s. 3 f. ult. opt. The Figures of this ed. 276 8 Ox. B. 230 30 T. 9 f. 3 f. ult. opt. The Figures of this Grafs exceed any Description. I have found it in Flower on Hownslow Heath about the Middle of July.

Hownslow Heath about the Middle of July.

SECT. IV. Gramen Juncea, fol. articulatis. Knot-rushes.

219 Tall Wood Knot rush, Ray 1307 4 Syn. 204 4 ed. 275 4.

91? Gramen Junceum fol articulato sylvaticum, CB. 5 8 phyt. 73

p. 16 Tb. 76 f. 8 Tab. 223 Icon 1 Hist. 659 f 1 Get. 20 f. 10 Lob. Icon 12 f. 1 B. 13 f. 3 Jonst. 22 f 9 opt. Park. 1270 f. 5 1 Ox. B.

227 1 T. 9 f. 1 f. This is much lesser than the next, its Panicles more fpread, yet smaller Seed Veffels. I's often found in moift

Moods; floursbing from Midsummer till Eutumn.

220 Common Knot rush Ray 1307 2 Syn. 204 2 ed. 275 2.

Gramen Juneum folio articulato aquaticum, CB. 5 9 phyt. 66 p.
15 Th. 76 f. 9 prodr. 12 f. 1 opt. Lob. Ic. 12 f. 1 B. 13 3 Tab. 214 Ic.
2 Hist. 252 f. 1 Ger. 12 f. 1 Ox. B. 227 2 T. 9 f. 1 f. Less than the last both in Height and Panicles. In wet marshy Places from Midsummer till Automa

Midfummer till Autumn. 221 Common Knot-rush with Utricles,

Gramen Junceum fol. arziculato aq cum utriculis, CB 7 5 prodr. 12 2 f. opt. Th 77 f 9 Park. 1270 5 f. 2 Ox. B 227 2 T. of. 1 f. ad Rad. A Variety of the last in having red leafy Heads, which appear in Autumn in marshy Ground.

222 Small Knot-rulb.

Gramen Junceum capitulis triangulis minimum, Ray 1307 3Syn. 204 3 ed 275 3 qr? Gramen Junceum, Ger. 4 f. 4 Tab. 225 lc. Hift. 660 f. 2 Ox. B. 227 3 T 9 f. 2 f. opt. qr? Park. 1270 f.xi. Its much lefs than the last: Common in moist and boggy Places at Hampfled, &c. flourishes about Midsummer.

223 Dale's Childing Knot-rush, R. 1307 5 S. 204 3 var. ed. 275 3 var. Gramen Junceum minimum paniculis foliaceis Bob. Ox.227 4 T.9 s. 2 f. opt. Pluk. 32 f. 3. Differs from the last in having small Leaves coming out of the Panicles: Time and

Place the fame.

Place the lame.

SECT. V. Gramen Juncea hirluta. Hairy Rush-grasses.

224 Small hairy Rush-grass, Ray 1291 I S. 193 I ed. 262 I
Gramen bifutum capitulis Psyllii, CB. 7 3 phyr. 81 p. 18
prodr. 15 f. 2 * opt. Tb. 103 f. 3 Leb. Ic. 15 f2 Belg. 16 f. 2 Jonst. 17
f. 3 Park. 1185 f. 6 Chabr. 188 f. 6 I B. 2 p. 494 f. 1 Tab. 219 Ic.

2. Hist. 656 f. 2 Ger. 16 f. 2. Ox, B. 225 4 T. 9 f. I f. It's the
least of all the following, Flowers in April and May in Mea.

down and Passurer frequently.

dows and Passures frequently.
225 Close-headed hairy Rush-grass.
Gramen hirsuum elatius, panicula juncea compassa. Ray 12012 S.1932 ed 2632 Ox B.2251 T.9 f.1 f. Gr. nemorofum histutum majus angustifolium, Park. 1165 f. 4 ad marg.opt. Its L. much longer and taller then the last with longer, and closer Heads Flourishes in May in Cain Wood plentifully.

226 Common hairy Wood Rush-grass.

226 Common hairy Wood Rush-grass.

Gr.nemorosum hirsutum vulgare, Ray 1292 3 S. 193 3 ed. 263 3 qr? Gr. hirsutum latifol. majus, CB. 7 1 Th. 101 f.1 Tab. 227 lc. 2 Hist. 662 f. 2 Ger. 17 f. 1 Park. 1184 f.1 Jonst. 19 f.1. These Figures easily distinguish this from others. It grows frequently in hilly Woods, Flowering, in the Spring.
227 Great hairy Wood Rush-grats.

Gr. nemorosum birsutum latifol. maximum, Ray 1292 5 Syn. 193 4 ed. 263 4 Chabr. 188 f.4 Th: 102 f.2 Park. 1185 f.3 Ox. B.
225 T.95 1 t. IB. 2 493 f.2 qr. CB. prodr. 15 f. 1. Its large spread panicles, and broad hairy L. distinguish it from the rest. Grows with the three foregoing in Cain-wood at Hampsted with the three foregoing in Cain-wood at Hampfled

plentifully in April and May.

SECT. VI. Gr. Juncea tomentofa. Cotton-Rushes. 228 Cotton-Rush, Ray 1306 1 Syn. 203 1 ed. 275 1.

Gr. prat. tomentosum, panicula sparsa CB. 4 2 phyt. 122 p. 26 Th.

61 f. 2 Tab. 230 Ic. 2 Hist. 665 f.1 Ger. 27 f.1 Chabr. 195 f.6 I B.

2 p. 514 f. 1 Trag. 683. Lob. Ic. 87 f. 2 B. 122 f.1 Adv. 44

Fonf. 29 f. 1 Park. 1271. f.1 Ox. B. 224 1 T.9 f.3 f. opt. Its many Heads of fine Cotton when ripe, easily distinguish it from the rest. It Cottons in June and July in boggy Moorish grounds as at Hampstead &c.

229 Hares-tail, or fingle- headed Cotton-rush, R. 1306 21 & 1308 xi, Syn. 203 17 ed. 274 17 Juncus Alpinus capitulo lanuginolo, seu Schænoglaurus, CB. 12 1 prodr. 7 p. 23 f. * opt. Th. 188 f. 1 phyt 30 pl. 10 Pink. 127 2 f. 6 Chabr. 199 f. 1 IB. 2 p. 514 f. 2 Ox. B. 244 6 T. 9 f. 3 f. Lob. illustr. 62. Grows in the Moors of Shrophire and Lancashire, &c.

SECT, VII. Gr. Juncea Miscellanea : Miscellany Rushes. Gramen Junceum spicatum seu Triglochin, CB. 6 16 phyt. 67 p.
15. Th. 81 f. 6 Tab. 224 Ic. 2 Hist. 65 f. 4 Ger. 18 f. 1 Park. 1187 f.
6 Lugd. 431 f. 2 Ox. 228 18 T. 2 fig. Lob. Ic. 17 f. 18. 18 f. 1 Jonst.
20 f. 1 Chabr. 190 f. 3 I B. 2 p. 508 f. 2 It flourishes in May, &c. in boogy Grands. in boggy Grounds.

or Doggy Grounds.

231 Sea Spiked Rush, Ray 1308 9 Syn. 249 to ed. 276 to Gramen Junceum & ipicatum alterum CB. 6 17 phyt. 68 p. 15

Th. 82 f. 17 Tab. 215 Ic. 1 Hist. 652 f. 2 Ger. 12 f. 2 Lob. Ic. 162 B. 17 2 Jonst. 13 f. 2 Park. 1278 f. 9 Chabr. 190 f. 2 I B. 2 p. 508 f. 17

Ox. 228 19 T. 2 fig. Very common in Salt Marshes; ipiking in May and line.

May and June.

CAP. V. Equiseta. Horsetails, &c. 232 Great Marsh Horfetail, Ray 128 1 S.20 1 ed. 42 1 Equiferum palustre longioribus setis, CB.15 2 phyt. 31 1 Matt. 725 f. 1 Theatr. 241 f.2 Tab. 251 f.2 Hill 698 f.2 Ger. 955 f.t Cam. 770 f. Dod. 73 f.1 Lob. Ic. 793 f. B. 688 f. obf. 461 f.1 Jonst. 1113 f. 1 Park. 1200 f. 1 Chabr. 551 f. 1 Ox. B. 621 1 This grows upright and is much the large first with us: Its Jules diffing from its learn Scalles. Except for the set of States. flinct from its leavy Stalks. Found in boggy Rills and marshy Grounds.

sny Grounds.

233 Lesser Marsh Horsetail, Ray 129 4 Syn. 21 4 ed. 42 4
Equiseium palustre, brevioribus setis CB. 15 3 phyt. 32 3 Th. 242
3 Lob. Ic. 795 f. 1 B. 970 f. 1 Jonst. 11 14 f. 4 Park. 1200 f. 3 Ox. B.
621 3 These Jules grow at the Top of its Leaves when they first spring up. Found in spongy rotten Bogs234 Corn Horsetail, Ray 128 2 Syn. 20 2 ed. 42 2
Equipment of compressions for CP.

Equiletum arvense longioribus setis, CB. 16 9 phyt. 32 5

Tb. 247 f. 9 opt. Park. 1202 12 Dod. 73 f. 2 Lob. 705 f. 2 B. 970

f. 2 T. 252 Ic. 1 Hist. 698 f. 3 Ger. 956 f. 3 Ox. B. 621 2 qrt Trag.
694 f. Its separate Jules spring in April, in moist Stubble and
Corn Fields. It grows supine and sometimes with secundary
Resolve Briftles.

235 Fine Wood Horsetail, Ray 128 3 Syn. 20 3 ed. 42 3
Equiserum sylvaticum, senuishmis setis, CR. 167 Th 245 f. 7 Tab.
253 sc. 1 Hist. 699 f. Chabr 551 f. 5 Ger. 957 f. 5 Park. 1201 f. 5
The Fineness of its Bristles differs it from others. It Jules at the Top, whilst the Leaves are young; in Cain-Wood or low wet Grounds.

236 Naked rough Horfetail or Shavegrafs, Ray 129 7 S.21 6 ed.437 Equisetum foliis nudum non ramosnm seu Junceum, CB. 16 10 phyt. 32 6 Th, 248 f. 10 * Chabr. 550 f.6 Trag. 692 f.1 This rises generally with but single Rushes and is known by its peculiar Roughness. Found in Warwicksbire and Wiltshire.

237 Doody's naked smooth Horsetail. Equiferum nudum lævius nostras, Ray 3 p. 103 Syn. ed. 426
Differs from the last in being smooth. Heads in May in moist
boggy Ponds and Ditches.

238 Buddles naked smooth Horsetail,

Equifetum lave penè nudum Nob. qu? Equifetum nudum, Tabe 251 Ic.1 Hist. 698 f.1 Ger. 955 f.2 Jonst. 1113 f.2 Park. 1201 7 f. These Figures pretty well resemble this Plant. Found by the Reverend Mr. Buddle on a new dug Bog on Hounstow-Heath.

230 Branched naked Horsetail, Ray Syn.ed 438

239 Branched naked Horsetail, Ray Syn.ed 43 8 qr? Equiseum foliis nudum ramosum, CB. 16 xi phyt. 32 7 Th. 250 s.xi Dod. 73 f. 3 * Lob. Ic. 794 f. 1 Jonst. 1114 f. 3 quoad Icon Park. 1201 f. Ox. B. 621 13 Found plentifully in Bocking River in Essex by Mr. Dale.

240 Mares-tail Ray 129 5 Syn. 21 5 ed. 42 5 Equiseum palustre, brevioribus foliis polyspermon, CB. 15 6 phyt. 31 2 Th. 243 f. 4 Tab. 835 Ic. 2 Hist. 2 505 f.r Ger. 957 f. 6 Chabr. 553 f. 2 Lob. Ic. 792 4 B. 967 f. 4 Jonst. 1114 f. 6 Park. 1201 3 Ox. B. 621 6 These Leaves are short and plain with Seeds at each Joynt. In Ponds and slow Waters.

241 Long leav'd Mares-tail. Equisetum palustre majus.

241 Long leav'd Mares-tail. Equifetum palustre majus, foliis longioribus. I have observed this in some Rivers.
242 Stinking Water Mares-tail, Ray 130 1 Syn.21 7 ed.

Equifetum fatidum sub aqua repens, CB. 16 13 prodr. 25 f. 5 opt. * phyt.35 8 Th.251 f.13 fonst.1115 f. Park. 1201 f. 10 Chabr. 551 f. 6 1 The Figures will give you an Idea of the Plant, it's commonly green under Water, but gray and brittle when dry. It grows in muddy Waters and has that sinking from flinking scent.

243 Warty Water Mares. tail, Ray 191 2 Syn. 35 pl. ult. ed.

Equiletum palustre ramosum, aquis immersum s. Millefolium aquaticum Equisetisolium, Ray Cat. Cantab. 49 descr. opt. I have sound this with its warry Heads in Battersey Rivulets plentifully.

244. Coralline Horsetail. Ray Hist. Plant. 130. xi. Synops.
21. 8. edit. 2. p. 43. 12. Equisetum seu Hippuris Coralioides, Jonst. 1115. 7. Pluk. Tab. 29. f. 4. In the Ditch about Chistleburst-Bogg.
245. Cluster headed Coralline Horsetail.
Equisetum Coralloide, seminibus in summitate conglobatis & folis absconditis, Buddle. 'Tis larger than the last. Observed by Mt. Mt. Buddle, in the Ponds about Heals near Islanich in

by Mr. Ad. Buddle, in the Ponds about Henly near Isswich in

offolk. 11/6 water Moss-Horsetail. Ray 3. 104. pl. 15. Syn. ed.

246. Irish water Moss-Horsetail. Ray 3. 104. pl. 15. Syn. ed.
43. xi.
Hippuris muscosus ex Hibernia, Pluk. 193. f. 6. Ox. B. 621.
10. First found by Dr. Sherard in Turf-bogs in Ireland, and since by Mr. Dandridge in England.
CAF. VI. Filices Britannica. British Ferns.
SECT. L. Filices foliis fere integris. Ferns with mostly whole Leaves or Wings.
47. Common Harts-tongue, Ray 134. 1. Syn. 22. ed 44. Lingua Cervina Officin. C. B. 353. 1. Cam. 579. f. Chabr. 557.
18.748. f. 1. Clus. 213. f. 2. Dod 407. f. 1. Lob. Ic. 2. p. 265. f. 2. Belg. 209. f. Obs. 486. Lugd. 219. f. 2. Jonst. 1138. f. 2. Park. 1046. f. Ox. B. 556. 1. Tab. 1. f. Ger. 976. f. Trag. 549. f. Its whole Leaves and slant feed lifts, differs it from all other British Ferns. Grows in stony shady Roads and moist Rocky places.

248. Fingered Harts-tongue, Ray 134, S. 22. d. 44. Ox. 557.

2.T. 1.
Lingua Cervina multifido folio C. B. 253. 2. Matth. 606. f. Chabr. 557, f. 3. I.B. 749. f. 2. Cluf. 213. f. 3. Dod. 267. f. 2. Lob. 805. Ic. 2. & 2. p. 226. 2. B. 981. & 300. f. Jonst. 1138. f. 2. Park. 1047. f. 2. Lugd. 1219. f. 2. Tab. 481. f. Icon. 801 1. Ger. 976. f. Varies from the Common in being Finger-tipt and found with it, but more rare.

249. Tree Polypody Ray 137. 1. S. 22. 1. ed. 45. 1.
Folypodium vulgare CB. 359. 1. Matth. 887. f. 1. Cam. 993. f. Chabr. 558. f. 3. Dod 464. f. 1. Lob. 814. 2. B. 992. f. O. 475. f. Lugd. 1229. f. Tab. 478. f. Ic. 799. 1. Ger. 972. f. Jonst. 1132. f. 1. Trag. 546. f. opt. Ox. B. 562. 1. T. 2. f. Very well known and common in Woods at the 1001s of old Oaks and other Trees.

other Trees.

other Trees.

250. Wall Polypody. Polypodium murale, pinnulis serratis
Novis Botan. Hort. Numb. 1. pl. 1. The wings of this are
somewhat narrower than the last, and lightly serrated. I gathered this Plant May 24. 1709. on the Walls of Windfor-Castle
inthe Apartments of the Revd. Dr. Manningham then Dean of
that place, whose Son a very Nice Botanist, first observed it
and shewed it methere.

251. Jagged Welfb Polypody, Ray 136. 2. S. 22. 2, ed. 45. 2. Polypodium Walliæ, pinnis altè laceratis Nobis Piuk. 30. f. 1. Difers from the Common in having wings deeply jagged Discovered AD. 1668, by Richard Kayse, of Bristol, on a Rock in a Wood near Dennis-Powis-Castel by Caerdist in Glamorgan-

Type.

252. Common Ceterach R.ay 139.c. 3. Syn. 23. ed 45.

Ceterach Offic. CB. 354. Matth. 646. f. 1. Cam. 640. f. opt.

Chabr. 556. f. 6. IB. 741. f. Dod. 648. f. Lob. 807. 1. B. 983.

O. 470. Jonst. 140. f. 1. Park. 10. c.46. f. Ox. B. 561. 1. T. 2.

f. Get. 978. f. 1. Trag. 551. f. A small Fern and being whony rushy below, differs it from others. Growson old Charches, Castles, and Stone Walls, particularly about Bristol, where I have seen it in plenty.

253. Lhwyds jagged Spleenwort. Filicula Alpina Pedicularis rubra folis. subra folis. subra solid.

rubræ foliis. subtus villosis, Ray 3. p. 63. S. 27. ed. 46. Pluk. 89. f. 5. Ox. B. 573. 28. descr. Mr. Limyd firit found this in the Clefts of moist Rocks on one of the highest Mountains in Wales, it grows also on Snowden, but rare.

254. Sea Penny Fern, Ray 140. c. 4. S. 23. ed. 46. 2. Trichomanes maritima major Nob. CB. 358. Lob. Ic. 814. 1. B. 992. f. O. 474. f. Lngd. 1226. f. 2. Jonft. 1143. f. 4. Park. 10. f. 7. Grows on Sea-rocks, and being much larger than the three following, is eafly diffinguisht.

three following, is easily distinguish.

255. Shop Maiden-bair or Penny-Fern Ray 140. C. 5. S. 23. ed. 46. Trichomanes five Polytrichum offic. CB. 356. Matth. 853. f. 3. Cam. 925. f. opt. Dod. 471. f. 1. Jonit. 1146. f. 1. Iak. 1051. f. Lugd. 1211. f. Tah. 477. f. 1. 16 801. 2. Trag. 530. f. 1. Its an Officinal and used in Pettoral Decoctions &c. Growson old Walls, Churches, and shady Rocks.

256. Branched Penny-Fern, Ray 140. & Vol. 3. p. 61. S. 27. ed. 46. 2. Ox. B. 591. xi. T. 3. t. Trichomanes dentata Nobis Chabr. 556. f. 1. IB. 741. f. 1. Ger. 985. f. 2. Tab. 477. f. 3. Ic. 802. Dr. Richard Kichardson, a Currous Botanis, has found this frequently on the high and moist Rocks of Wales as

also on Inglebrough, but could rarely observe it branched; Dif.

allo on Inglebrough, but could rarely observe it branched; Differs from the Common in being more serrated.

257. Fagged Venny-Fern. Trichomenes fol. eleganter incisis Petiver Ray 3. p. 62. Puk. T.b. 73. f. 6, qr? Ox. R. 501. 12. & 13. T. 3.f. opt. Ray 3. p. 61. This is much deeper cut than the Last, especially its lower wings.

258. Matthioles Lancet Fern Ray 138. 1. S. 27 ed. 45. 2. Lonchiris aspera C. B. 350. 1. Matth. 661. f. 1. Cam. 664. f. Chabr. 556. 3. 1. B. 736. fig. Lugd. 1220. f. 1. Ox. 566. 1. T. 2. f. Jonst. 1140. f. 3. Park. 1042 f. 1. Gen. 979. f. 1. Tab. 480. f. 1. Ic. 799. 2. That most accurate Northern Botanist and my Worthy Friend Dr. Richard Richardson sent me this which my Worthy Friend Dr. Richard Richardson sent me this which he gathered on the Rocks of Phainon vellon in Wales. He tells me, its also pretty common on the high Rocks of Snow-

259. Lhwyds Lancet-Fern, Ray 3. 84. Syn. ed. 48. 4. Filix aculeata Lonchiridi emula noffras Pink. 180, f. 3. Ox. B. 567. 7. var. I am oblig'd to Mr. Jacob Bobart, Botanick Frofesfor of the Physick Garden at Oxon for a Specimen of this, which differs from the last, its wings being nuch jagged and more prickly, Grows on the Welch Mountains.

260 Pluknets Welfh Lancet Fern, Filix parva Cambrobrit. pinnulis latis auriculatis brevibus aculeata Pluk. Tab. 89.

f. 6.

261. Hysvae Fern. Filicula Alpina tenerior, alis latiusculis brevioribus integris profunde dentatis O.c. B. 576. 23. descr. Pluk. 179. f. 4. secund. Ray S. ed. 48. 4. On the tops of Snowden and Hysvae rocks near Penrice Church in Caernar-

262. Great Prickly Fern. Ray Syn. ed. 48. 3 Ox. P. 580 16. Lonchitis aculeata major Nob. Each stem has 15 or 16 wings, and each wing between 30 and 40 Feathers or pair of leaves,

Grows in shady ditches.

263. Common Prickly Fern, Lonchitis aculeata vulg. Ray 143. 3. Syn. 24. 2. ed. 48. 2. Pluk. 179. f. 6. Ox. B. 580. 15. T. 3. S. 3. Jonft. 1130. 4. In moist and shady displace. ditches.

264. Earless Prickly-Fern, Lone bitis aculeata fine auriculis, Ry. Syn. 24. 2. ed. 48. 2. var. A Variety of the last. 265. Pluknets rusty Prickly-Fern. Lone bits aculeata, lanugine ferruginea. Pluk. 180. f. 1. Ray 3. p. 86. Less and paler than the lait, with rusty stems. Grows in rotten hogs.

SECT. II. Filices non ramosa. Unbranched Ferns.

266. Great dented Male Fern. Filix non ramofa latifolia dentata Inft. Rei Herbir. 536.7 Hist. Plant. Faris 443. Schol. Bot z. These wings are much larger than the next, as is the whole Fern. The Figure in Camerarius on Matthiolus p. 994. does not ill repretent this plant.

267. Common Male-Fern. Filix mas Dod.462.f. 1. Cam. 991.f. Trag 546 f.Chabr. 553.f. i. It. 737.f. i. Park. 1036.f. Lugd 1222.f. i. Filix non ramofa dentata CB. 358.3. Muth. 886.f. i. Jonft. 1129. 2. defer. Tab. 2.p. 471.f. 1. Ic. 791. 2. Ger. 968., f1. Ray 143. 1. Sy 11. 24.

1.ed. 47. 1.

268 Mars Fern. Filix palustris feu aquatica. Ray 146. c.41.

Syn. 24 4.ed. 48 6. Its Feathers plain and seed edged, by which and its lower wings gradually lessening, its known from the last. Grows in Bogs and Moorish places.

269 Rugby Fern. Filix palustris major Nobis Ray Syn.ed. 341. pl. 2. Seems to be a larger kind of the last, Found on Duns-moor-heath in the Ditches which parts Rugby from Dunchurch

in Warwick-frire.

270. Male Stone Fern. Filix faxatilis non ramofa, nigris maculis punctata CB, 358.3. Cluf. 212.f. 1. Pan. 706 f. Park. 1044. f.1 Pluk. 179.f. 5. Ox. 581.28. Ray 146 2. Syn. 24.5. ed. 48.7. Differs from the Mar/b Fern, in its foil and being lefs. Found plentifully on the high rocks in Darby-spire, Yorkshire and Westmoreland.

271. Forked Stone Fern. Filix minor Britan. pedic. palli-

271. Forned Stone Fern. Falix minor Britan. pedic. panidiore, alis inferioribus deorium spectantibus Ox. Rob. 571.17. defer. gr? Pluk. 284.2, Its lower wings bend downwards. Grows about the Peak in Darbysbire.

272. Goodyers crisp Fern. Rsy 144.4. Syn. 24.3. ed. 48.5. Filix mas non ramosa, pinnulis angustis rais profunce dentatis Font. 1033. defer. Pluk. 180.f. 4.Ox. B. 579.8. T. 3. Its Finne or Feathers are deep notcht not unlike curl'd Tanfy. Grows in wet Shady places. Shady places.

273. Fingertipt crifp Fern. Filix minor non ramola, alis foliorum alternatium positis bishis & multisidis, Ray 3. p. 78. Syn.ed.46.6.var. Pluk. 284.f.3 Ox. B. 580.21. deler. Differs from the left in being for the different control of the property of the control of the property of the control of the contro the last in being fingertipt. Sr. Tho. Willoughby observed this on Litchfield Mintter.

SECT. III. Filices ramosæ. Branch wing'd Ferns or Brakes. 274. Common Female Fern or Brake. Filix sæn. vulg.

L. 1. 1.5845

Offic. Ray 150, c.2 Syn.25 1 Ed. 49.1. CB. 357.1. Matth. 886. f.2. Lugd 1222. f.2. Cam. 992. fig. 18.3. 1.37. 727. Dod 462. f.2. Jouft. 1128. f.2. Park. 1037. f. Ox.B. 583 3. Tab. 2471. f.2. 1c. 793. f.1. Ger. 969. f.2. So well known, and very common on Heaths, it needs to Deteription.

275. Wall-Brake. Filix ramofa, five Fæmina minor noffras, Ox.B. 584.18. defer. Ray 151. c.4. Syn. 25, 4. ed. 50.5. I have formerly observed this on several Walls in London; it's also found on the Sea-Rocks in the West of Free land.

276. Small Brake. Filix ramofa minor nobis. Chabr. 555.

276. Small Brake. Filix ramofi mivor nobis, Chabr. 555. F.z. 1B.733. f.1. Ray 151. c.7. Syn.25.5. ed. 50. 6. Cluf. 212. f.1. Pan 704. f. Lugd.1225. f. 3. opt. Jonit. 1135. f. 3. Park. 1044. f. 2. Trag. 538. f. qi? Tab. Ic. 794. i. H. 472. f. 3. In they Places on the Sides of Mountains in Westmortand and Wales. Wales.

277. Common dented Brake. Filix fxm. pinnulis dentatis Nobis Ray 150. c 2. defer. Syn. 25.2. ed. 49.2. Ox.B. 579. xi. defer. T.3. f. Grows in shady moist slony Places, and often at the Posts of Father P.O. k.

the Roots of rotten Oaks.

278. Cicely Brake. Filix Alpina Myrrhidis facie Cambrobritannica, Pluk. 89. f.4. Ray 3. p 94. Syn. ed. 49.3. Ox. B. 584. xi. The lower Wings next the Stalk are much the lengelt. Mr. Lbwyd first discovered this on the Top of Mount Glyder.

279. Hemlock Brake. Filicula Cambrobritannica pinnulis Cicntaria divitura, Bob.Ox. 581.31. Ray 3. p.84. Syn. ed. 50.8. It's composed of about 8 pair of Wings with deep cut Leaves.

Grows on the Top of Snowdon.

280. Small brittle Brake. Filix faxatilis, caule tenui fragili, Ray 152.1. Syn. 26.6. ed. 50.7. Ox. 581. 29. 41? Chabr. 555.4 IB. 734. f 2. On Old Stone Walls and Rocks in Derby-fhire, Torkbire, a. d Westmorland.

281. Round brittle Fern. Filim faxatilis, caule tenui fragilishumilis a familia poundi ithan P. Rochardon. That and

li humilior, pinnulis rotundionibus R. Richardson. Under this Title I received it from Dr. Richardson, who discovered it on Snowdon, and plainly differs from the last in having rounder Leaves.

282. Common llack Maidenbair, or Fern. Adiantum nigrum vulgare Nobis. Ray 151.2. Syn. 26.7. ed. 50. 9. CB. 355. 2 1B. 3, 1.37. p. 734. f. Ded 466. f. 1.0b. 870 Ic. 2. B. 988. f. Adv 361. John. 1137. f. Park. 1049. f. 2. Ger. 975. f. r. Tab. 1c. 796. 2. H. 474. f. 2. Ox. B. 588. 76. T. 4. f. Grows at the Poster of Transact Shark. in Shada Market and an Ott the Roots of Trees and Shrubs, in Shady Woods and on Old

283. Great black Fern. Adiantum nigrum fol. duplo majoribus faturatæ magis viriditatis, cauliculis quoque longioribus & nigrioril us, Sibbald. Prodr. 7. Tab. 4. f. 1. Found in Lockcoat-

Hills in Linkithgow,
284 Natiow black Brake. Adiantum nigrum segmentis an

gustionibus, Ray Syn.ed. 51. 9. var. 9? Fluk. 124. 6.4 Ox. B. 588.17. It's narrower and deeper cut than the Common. 285. Prenton's Fifeness black trake. Adiantum nigrum fol. Lunarme minoris. Ray 3, p. 61. Its Leaves broad and tomewhat dented, the Seed differ like Hartflongue. Dr. Fresson such discontinuous descriptions of the Seed of the Seed difference of the Seed of the See

vered this on the Sea Coasts at Fifeness in Scotland.

286. Round Jesse, black Brake. Fiti elegans, Alianto nigro accedens, segmentis rotundioribus, Ray 3. p. 85. Syn. ed. 55.xi. Ox.581.25. de.cr. Dr. Sherard and discovered this on 55.xi. Ox.581.25. de.cr. Dr. Sherard with discovered this on the Rocks in Justey; and Mr. Jac. Bubare has finee observed it on Addition Curch in Oxfordfine.

287 Fine Irish Fern, or Mourn Brake. Filix minor longisolia, tarsis raris, finnulis longis tennissimis & ol longis laciniis simbriatis, Ray 3, p.79. Syn. ed. 51.12. Its Leaves retemble Dr. Pluknet's Tab. 89. 11g. 3. but its Tinfi, or Brancher stand more distant. Dr. Sherard found this on the

Branches fland more diffant. Dr. Sheraid found this on the Mountains of Mourn in the County of Down in Ireland. 288. Shop l'enushair, Ray 147 1. Ox.B.587.6. T.5. f. Adantum tobo Coriandri, Ch. 355.1. Matth. 832. f. 1. Cam 924. f. Chabr. 555. f. 3 IB. 3. 741 fig. Dod 469. f. 2. Lob. lc. 809.2. B. 986. f. .. dv. 302. Tab. ic. 795.1. H. 473. f. 1. Ger. 982. f. 1. Lugl. 1208. f. Trag. 530. f. 1. Found in Wales by Mr. I. hwyd.

289. Adianto vero isfine, Adiantum notiras majus, Coriandri folio in apice nounihil rotundo, pediculo pallide rul ente Sibbald prodr. 7. Ox. 587.7. Dr. Irvin found this in the Grots or Faults about Weyms, Scotland.

290. Adianto vero affine n' firas mirus, fol. obtufo, faturatè vindi alian incho, cum pediculo vindi minore Sibald. pr.8. Found in the Gap of a Rock on Leebesat lake near Limnuck, Scotland.

SFOT. IV. Filleulæ. Dwaf Ferns.
291. White identiair. Filleulæ ella valgaris.
Ruta natratia, Cl. 354. R 146. c.5. 11.1. Syn. 24 ed. 48. Ox. 585.
22.T. 5. f. Com. 785. t. Chair. 555. f. o. IB. 745. t. Lob. Ic.
812. I. B. 988. f. Olf. 472. f. Lod. 470. f. Jonft. 1144. f. 3.

Park. 1050. f.4. Tab. 10.796.1. H 474.f.1 Ger. 983. f. Trag. 530. f.2. On old Stone-Bridges, Walls, and Clefts in Rocks. 292 Forked Maidenhair, Ray 141.c. 7. Syn. 24. ed 47. Ox.B.

585. 23. T.5 f. Filix fax:t. five Filix las it. five cornicul ita CB. 358.5. Fonft. 1045. f. 8. Lugd. 1226. f 1. Trag. 437. f. Grows on the Rocks in Edinburg-Park. and on some Mountains in Wales.

and on some Mountains in Wales.

293 Date's Fern Darea Tunbrigensis minor Mus. nost. 761.
Ray 1583. descr. Bocc. Mus. 24. T. 2. Iluk. 3. f. 59. I have otherwes this in relaty on the Rocks near Tunbridge-wells.

294 Newton's Dare-Fern. Advintum petræum perpusiblium Anglicum, sol. bissids veltrissids Newtoni, Ray 141. c. 6. Syn.

23.ed. 47. Pluk. 3. f. 6. Found near Wrenose in Westmoreland among Mess, Mr. Lhw, d has observed this in Ireland half a span-high.

SECT V. Filices Florida. Flowring Forns.

SECT V. Filices Florida. Flowring Ferns.

295 Ofmund R yall. Oliminda Regalis Offic. Ray 151. c. 5.

Syn. 25.3 ed. 50.4. Ox. 593.1 T.4. fig CB. 357.3. Chabr. 554 fd.

1B.728. t Dod. 463. f Lob. 813. IB 992 f Obf 474. Lugd. 1225.

i 1. p. 9. Ger. 971 f Jonfl. 1131. f Park. 1038.f. Trag. 543. f.

opt. Flowers in July and Aug. on rotten Bogs and moist Woods.

216 Flowring Spleenwort. Lonchitis. Florida vulg. nobis.

Lonchitis minor, CB. 359. 2 Matth 661. f. 2. Ray 138 2. Syn.

23 ed. 45 1 Cam. 665. f. Chabr. 556. f. 4. & 5 IB 737. f. 1822. Cluf.

213. f. i. Lob. fc. 815. 2. R. 994 f. O. 475. f. Lugd. 1276 f.

opt. & 521. f. 2. Jonfl. 1140 f. 2. Park 1042. f. 2. Tab. lc. 800. I. H.

480. f. 2. Ger. 978 f. 1. Irag. 550. f. opt. Its feed bearing reaves are much narrower than the other, and grow upright in moith Heaths and Woods, and shady Ditches. Seeds in July.

297 Otton Flowring Fern. Ofmunda Westmorlandica foliss.

297 Orton Flowring Fern, Ofmunda Westmorlandica foliss, tenuis ne dinectis Mus. nost. 792. Ray 153. 3. yn. 26. 8. ed. 51.10, Ox. B. 593. 4 F. 4 f. Pluk 3. f. 2. opt. Grows on Snowden-bill in North Wales, and Inveral places in Westmorland, particular of the contraction of the contraction of the contraction of the contraction.

I ray at Orton.

Try at Orion.

298 Moonwort. Lunaria recemble minor & vulg. CB-354 r.

Matth.647 f.2. R.y 127.1.8)n. 2ed 44 Ox.B.594 r. T.5. Cam.
643 f.C. ab. 544 f.6.1B.710 f. Dod. 139. f.2. Lob. lc. 807.2:B.983.
f. Cibi. 470. Jonit.405.2. Park 507.1 Fab. lc. 137.1. H. 513. f.2.
Ger. 320. t.2. Trag. 914. f. Flowers in May, &c. in cold pattures, in ly ground and old Gravel pits.

299 Jagged M. onwort. Lunaria minor, fol. diffectis Ray 3.
p. 101. Its larger than the common, and Flowers more branched Found with the last.

Found with the lift.

300 Addertto stie, Ophioglossum vulgatum, CB.354.1. Matth. 447.f.5. Ray 126.2.10.5yn,21.ed.44.0x.B 595.T.5 f. Cam.364.f. Cipibr.541 f.4 p. 126.2.10.5yn,21.ed.44.0x.B 595.T.5 f. Cam.364.f. Cipibr.541 f.4 p. 126.2.10.2.7 f. Park. 506. Tab. Ic. 849.1. H. 519. 1.2. Get. 327 Trag. 323. f. It tongues in May in moist dieadows and Pastures.

CAP. VII. De Mufcis. Of Earth, Tree and Water Mois. SECT. I. Lycopedii. Wolfs-claws or Club-mofs. 301 Common Wolfs claw, Ray 120.1 Syn.16.1. ed.25.1.

Muscus terreliris clavatus, CB360.x. Chabr. 559. f.4. IB. 3. l. 37. p. 758. t. Cord. 111. f.1. Dod. 472. f. 2. Lob. lc. 2. p. 244. f. 2. Belg. 251. f. Obl. 645. f.1. Lugd. 1324. f.1. Jont. 1502. f. xi. Park. 1307. f.4. Ger. 137. f. x1. Tab. 814. Ic.z. Hift. 814. f.3. Ox. B. 645.1.

A Creeping Moss with 1, 2, or 3, Claus standing on long stoutialks, Heads in July and August, in Heaths and Moun-

302 Creeping small Club moss with Upright heads. Lycopodium repens minus, clavis erectis Nob. Ray 121.c.3.Syn. 17. 3. ed. 27. 1 Pluk. 205. f. 5. fig. mala Ox. 624 x1 Tab. 5. fer. 1.f. x1. Its many fibrofe roots keep it clefe to the ground, it's hardly known till in bead, thele grow upright and about two Inches high in boggy Heaths, about July and August.

303 Jumper Wolfs claw Ray 120. var. Syn. 16.2, & 18:10.

ed.26.2. Musicus terrestris fol. retro restexis, IB. 3. 1 37 p. 759. Ox. B. 624. Tab, 5. ser. 1. f. 3. opt. Differs from the Common in having shift and restex Leaves and res claws without stalks, Found on the Welfh Mountains.

304 Upright Firr Club-mofs, Ray 121. c,4 defer. opt. Syn. 17.4. ed. 27.2.

L)copodium erectum Abietiformis majus Nobis, Ox. B. 624.9. Tab. 5. fer. 2. fig. 9. opt. where its very well express. It's upright and monthy forked, with Leaves very green and close to the flalks, which are much thicker than any of the foregoing; it feeds in May and June, they are small and round. This has no diffinit claws as most authors has Figured it. Grows on the high Mountains in Wales, as also on Ingleborough in Yorkshire and the Peak in Darbyshire.

305 Rays small upright, Firr Club-moss Ray 122. c.5. Syn. 17. 5. ed. 27. 3. Lycopodium erectum Abietiformis minus Nob.

Pluk. 47. f. 7. Ox. B. 624. 10 T. 5. f 2. f. 14. Much lefs than the last and lower, its Leaves not so close, but more starting off, under which lie its seed along the most part of the stalk. Found in the Rills and moist places on Snowden, Ingleso-

good Cypress Club moss, Ray 12. 2. Syn. 17 2. ed. 26.3. Ox. B.024.5. Where its well described but no where truly Figured, its twigs rarely branched and mostly fingle claw'd without footstakes. On the Mountains of Wales and Ingleborough and Anderkin in Scotland plentifully.

307 Bobart's Cypress Club moss, H. Ox. B.624 5. var. Tycopodium Cupress folis, 4 julis, nudis pedicules Nob. Its naked footstak a Hand long, ending with 4 claws on each stake. This should rather resemble the Figures which Authors give of the

196. Clusius his Portugal M of Ray 112. 3. Syn. 13, 2. ed. 21.
2. Lycopodium repens squamesus Nob. Muscus denticulatus misor CB. 360.8. Clus. 249. f. 2. Lob. 2. p. 244. f. 1. O. 644. Luzd.
1326. f. 1. Dod. 473. f. 1 Forst 1162. f. 10. Ger. 1373. f. 9. Tab.
184. f. 2. 806. lc. 1. H.
186. f. 187 he found this at the Lead mines in Somerfeishire,

where I have fearcht for it; but neith rethere, nor any other part of England have I yet met with it, but from several places abroad I have had it sent me.

apronal have had it tent hie.

1309 Bobart's Tree Club mofs, Ox.B.6248. T.5. f.2 f.8 opt.

Ray Syn. ed. 338. pt 5. qt. Kay 3. p 47. Syn. ed. 2. p. 39. 16.

The tips of this Mofs for the mon part of Winter are full of Powder ; it grows on the bodies of theses &cc. frequ nt-

SFCT. II. A usci terrest. majores. Greater Earth Mosses. 310 Fern Moss, Ray 112,1. Syn.13.1. ed.21.1.
Museus Filicinus major CB.360.4. Park. 1310. f. 3. Jonst. 1561. f. Chabr.759. f.5. 18.757. f.1. Ger. 1372, f.7. Tab. Ic.607 1. Hist.83. f.4. Ox. B.625.20 T.5. ser. 3. f.20. Common in pady Woods, it beads in Dec. and Jan. but rarely.

21. Small Fern Ajos. Museus Filicinus minor CB.360.2.

311 Small Fern Ajofs. Muscus Filicinus minor CB. 360.2. prodr. 151.2. ex Sent. Bob. Ox.625.21. T.5. f.4. f.21. Fine

leaved like the last, but less.

312 Vernon's Tamarisk-Moss. Muscus ramosus repens spicatus CB. 361.4. pr. 151.2. Ox. B. 625.6. T.5 1 3. t. 6. Its heads crooked, longish and end pointed. Found at the roots of

313 Braynes crested-Moss. Ray 3. p.47. Syn. ed 38-13.
Museus crittam califensem representant flivescents & c. Mus nost.
84 Ox. B. 625.8. T.5. s. 3. f.8. Flor. Prus. 166. f. In moss

Woods and Boss.

314 Bonavert's crifp Moss. Mnscus squamosus superficie crispu Nob. Ray 3. p.44. 391. ed. 337. pl. 3. Ox B 625. 10. T.5. 13. 110. It Lears its beads on the stiller, rearthe top of its Leaves. Thave gathered this on the Chalk Banks by the Thames fide, between Purfleet and Gravelend.

315 Cou mon Wood Mofs. Muscus terrestris latioribus foliis 315 Con mon Wood Moss. Mulcus terrestrus satisfibus fossis major, seu vulgatis, Ray 122.1. & Vol.3 p.43. pl 8 624.1. H. Ox.B. Syn.17.6. ed.36.1. But the Syronyms there to CB. IB. &c. belong to No 329. This Moss is very common in stady Woods and Hedges. It Heads in Dec. and Jan.

316 Sherard's common Irish Moss. Mustus eredus major, solid angustioribus acutis, Ray3 p 44. 11.1 Syn. ed.237. Like the last, but with sharper and thinner Leaves. Common in beland.

317 Doody's hook-leaved Moss. Mufeus erectus folius reflexis, Ray3. p 44. Syn. ed. 337. pl. 4. Ox. B.624.2. T.5. f. 2. Its crooked Leaves and being tomewhat lets, differ it from No. 315. Grows in wet Graff, Grounds.

318 Doody's hook leav'd Bog-Mois. Muscus palustris scorpides ramofus erectus. Its Leaves and Tips are both reflex. Found on the Bogs behind Charlton.

310 Ray's long pointest Eco. Moss. Muscus valustris valde.

319 Ray's long pointed Eog-Moss. Muscus palustris valde ramolus, furculis erectioribus, foliolis in tennes & longos mu-

crones productis, Ray 3 p 47 Syn ed 39 14. 3 320 Vernon's, Hinton-Moor Mets. Museus palust, major, fol. membranaceis acutis, Ray 3 p 47. Syn ed 39.19. Its Stalks long and golden, with a crooked Cap and Head, growing from the Bottom of the Leaf, in May and June, in most Bogs and

321 Our common Copress Moss. Muscus terrest. vulg. Currest solins, Mus, nost. St. Ray 3 p. 45 & 47. 116. Syn ed 39. 20. Muscus Currest formis ranolus, CB 361. 9. pr 152 9. ex sent. Inst. R. H. 55 4 pl 7. Ox B 626.27. T 5. 14 t 27. common in Woods and some Passures. Heads in December.

222 Buddle's sender Cypress Mots. Muscus Cuprest formis, sagelis tenuioribus & fol. acutis.

323 Buddle's thick Cypress-Mots. Muscus Cuprest formis, rinculis brevioribus & crastioribus ferè rotunds, & suppriss.

vinculis brevioribus & crassioribus fere rotundis, & supinis.

324 Doody's upright Flat-Mols. Museum erroins sol. anguatis caulibus appressis, Ray 3, p.44; pl. 1. Syn. ed. 327; pl. 1. It. Museum ramotus eresus may i CB.3616. frod 1515, ex. lent. Instit. 553; pl. 3. Common in harren Woods.

325 Buddle's Suffolk House-Mols Museum downstine non-standard received a president of the synthesis o

Buddle First found by him and only in Suffel's on Barns or

Thucht Hales.

Bouch Honfes.

326 Pluknet's flender Mountain-Mofs. Mult in Cup. offines, Bocc. Mult, p. 161. F. 1. 108. Infinisterreft. Achorum Lipiti us advances major & erect or Ray 3. p. 47. Syn ed. 30. 15. & R.3. p. 33. & 38. 11. 42. Ox B. 624. 7. F. 5. f. t. f. 5. P. uk. 17. f. 1. Int. 555. pl. z. Lis Heads long and upit by on the Stalks, but rare-555, pl 2. Its Heads long and upit be on were states, but rasely feen. I have observed it plentifully in fome deep fluidy, chalky Lones.

**The Act of the form of the value limiting fus.

327 Hairy Welch Moss. Nus us terr st. vulg. Linuvinosus Alpinus, Rry 3, p 45, pl 3. Syn 18.xi. ed 37.5. Fink 47. sig 5. Ox B 625.7. P 5. 82. f ult. gr. CB 300.9, Get 1300 f 1. Its Stalks and Heads thort, very common on the Mountains of

328 Our Hairy Mountain-M. s. Muss is industrial lunginosus Alginus, I us. nost. 83. It's much smaller than the last. This I found in the Sweedland Holls in Leicestershire.

329 Common White Pog-Moss, Ray 122, Syn 18.9 ed 37.4.

Ox B 625,4 Muscas palustris in Ericetis nations forilus Pluk tor. f 1. Muscus terrestris vulg it simus, C \(\frac{3}{6}\), t. (3) \(\frac{4}{7}\), f. \(\frac{7}{2}\), p. \(\frac{7}{2}\), p. \(\frac{7}{2}\), f. \(\frac{7}{2}\), p. \(\frac{7}{2}\), p. \(\frac{7}{2}\), g. \(\frac{7}{2}\), where its green, but white when dry. It bears round black stables Heads in June.

330 Red Bog Moss. Museus palustris rubedine tinclus, Normegicus, Nob. Ox B:625,4. Var. Its a Variety of the last tur-

megicus, Nob.Ox B:625,4. Var. Its a Variety of the last turning red when dry.

331 Brotherton's fine white Water-Moss. Musicus erestus palustris albus, foliis capillaceis, Ras. 3 p. 45. pl. 1. Syn ed. 338 pl. 1. Found by this Curious Gentlem in in Lancastrice.

332 Ray's Tree, like Moss. Musus Dendroides sylvarum, erectis cannolis Kili mmulis, rad. repeate, Ray 3, p. 35. pl. 23. Syn ed. 32.22. Ox B. 626, 30. Tab 5. f. 4. f. opt. Is f. 554 pl. 5. This has a long niked scaly Stalk spreading above like a Tree. Heads in March at the Roots of Feld Trees in moss two des.

333 Vernon's Tree like Miss. Musus Dendroides elation, ramusis crebris minus surculosis, capitus strevila, fediculus insidentilus, Ray 3 p. 36 pl. 23. Syn ed. 32.23. Ox 6.6.6, 37. T. 5. f. 4. f. cpt. Instead pl. 23. Syn ed. 32.23. Ox 6.6.6, 37. T. 5. f. 4. f. cpt. Instead pl. 23. Cheeved by Mr. Kennon at Gamlingay, and by Mr. Buddle in barren Fields near Information.

SECT. III. Musci to restres minores. Small Earth Mosses. 334 Velvet Moss. Multus terrestris vul aris minor, Adameticarei cipiulis, Ray 122. Syn 18,8. ed 36,2. R 3, p.43. Pl Paris 502. pl 2. Inft 553.pl x. Meads in December & . from the Sides of its branches in burren Pattures, and in Woods at the Bottom of Trees and Shrubs.

335 Ray's fmailer Earth M Is. Muscus terrestr. repens, 1

freeier familis, fed multo minor, Ray 3. y 44. pl 4 Syn ea 38 8 B Ox 625,5 T 5. f 2. f Intl 533 pl 5. The Leaves in this are paler and fmaller than the lath, with which it is also

found.

336 Turnfort's crookbeaded Moss. Muscus squemosus ramedus crassion capitulis incurvis, Inst 553, pt 12. Listers from 334, in being thicker. Heads in December and January, at the Roots of Trees and Strubs.

337 Ray's small mort headed Moss. Muscus terrestr's parvus, capitulis brevibus tumidis nominal means & rotundis, Riy 3. p 45. Syn. ed 38, 10. Its Heads shirt, swelled and somewhat bending, coming out of the Sides, on a half-inch Stalk on a bulbous Base. Found with the last in December or some

338 Our creeping Moss with shining Leaves, Must nost 83. Mulcus terrestris luteo viridis sericeus repens, Ray 3 45 S. n. ed 38, 12. Ox B 626 5. T 5. s 4. s. Common on Brick Walls near condon; it's easily known by its creeping and Silky Gloss. Heads in Winter.

330 Doody's shining Tree M. s. Museus repens serici modo lucens, viticulis longloribus electis, Ray 3. P. 44. 1 lult Syn ed 338, 5. Ox B 626, 26. T 5. f 4. f Snines like the last, with upright thicker Sprigs. Grows on Trees, &c.

340 Bobart's curled Godflow Mols. Muscus terrestris repens jubility was, foliolis cripus innormalis ramididus centures conferens, Ox 625, 12. T 5, 13. f. Its Spris tofe, y bowth, filly and curled. Grows on the Walls of Godflow Charles and 2 Miles from Oxford.

341 Balles curled Rog. Moss. Muscus scorpoides palustris, fol. crissis pyramidalibus, Ray 3, p.28 & 35. Syu ed 22, 36. Ox 625.9. A yellow Moss with curled and coyling Sprigs.

Ox 625.9. A yellow Mots with curred.

Found by Mr Davies on Hinton-Moor.

342 Bokart's finall Fir-moss. Must us terrest. furculis filamentosis tenacibus Abietinis semel tontum divisis, Bob Ox.

11. f Inst. 554.4. Its Sprigs single brancht,

mentons tenacibus Abietinis iemei tontum divins, Boo Ox. 626.22 T 5. f 1. f Infl. 554.4. Its Sprigs fingle brancht, creeps among ft the Grafs on dry Hills.

343 Ray's finall Fern-mofs. Muscus terrest, parvus supinus Filicis modo interdum penuatus, Ray 3. p 45. Syn ed 38.9. Fl. Pruss. 167. pl 454. f It's like N°334, but less, and ends in longer Sprigs. Found in Brainbly Places.

344 Ray's small Moss with big Heads. Muscus terrest. omnium minimus, capitulis majuscutis oblongis erectis, Ray 3. p 45. Syn ed 38. xi. Ox 625.14 T 5. f 3. f 6. On Sheds Pales, & c. Mr Buddle has alto observed it at the Bottom of Trees in moist Places in April.

345 Ray's Stonecrop Moss. Muscus terr. surculis Kali aut Illeccbra amulis, foliolis subrotundis squamatim incumbenti us Ray 3. p 45 Syn,ed 37.6. Ox 626.28. Common in Woods at the Roots of Trees.

346 Ray's sharp-pointed midling Mols. Muscus terr. medius, inpirus & repens, foliolis crebris in acutos mucrones productis, R 3. p 45. Syned 37.7. Common on Thatch and Tree-Roots in Nov. and Decemb.

347 Doodyes small flat faced Moss. Muscus fol, caulibus appressis uno versu dispositis, viticulis minoribus. Refembles

the last, but less, and found with it.

348 Doodyes common Flat Tree Mols. Muscus vulg. minoi, cauliculis compressis, Ray 3. p 47. Syn. 244. pl 3. ed 39.18. Inst 555. 3. Heads in Dec. and Jan. at the Roots of imall Trees.

349 Dr. Richardsons Club-faced Bog-moss. Muscus terrest. repens, Lycopodii sermè facie, Ray 3. p 43.6. Syned 337. pl 5. Ox 626.24. T 5. [4. f. 1 have met with this pietty Moss in some most Ditches in Bishops Wood Hampsted. Heads about Midsummer.

350 Buddles Wood Feather-mofs. Muscus pennatus sylvations. q1? Ox. 626. 35. T 6. 1 r. f. Grows on the fandy Banks of a dry Dirch between Cain-Wood and Mother Huffs.

Heads in May and June.

351 Scamptons Peak Moor-moss. Museus aquat. pileis acutis Mus. nost, 74. Fig. Ox 626, 29. T 5. f.4. f. First discovered by this Gentleman on the large Stones which Le in the Rivulets amongst the Peak-moors in Desbyshire.

352 Buddles Feathered Water-moss. Muscus pennatus a-

quaticus. Grows on Boughs that lye in the Water in Ponds,

&c. Heads in Winter.

353 Doodyes Feathered Water-moss. Muscus pennatus aq. Ray Syn ed 388. Observed by him in the Thames, and Dr.

Richardson in the North.

354 Dr. Sherards Irish Floating Moss. Muscus fluitans, fol. in flagellis longis tenunbusque, Ray 3. p 45. pl 12. Syn ed 338. Found by him in the Pitts of the thaking Boggs in Ireland.

M.I. Buddle has also observed it in August in our Winter-Ponds, viz. fuch as are near dry in the Summer.

SECT. IV. Musci scl ferè pellucidis. Glass-mosses.
355 Sherards Bushy Glass-moss. Muscus trichoides ramofus, fol. longis pellucidis & veluti ramofis, R.3. p 43 pl 4. Syn 243. I ed 36.6. Pluk 49. f 1. Ox B 630. T 6. I ult. f r. Mulcus Polygoni folio, Pl I ar 504. Inft 555. Flor. Pruff. 168. pl 457. f. 46. This Moss branches at Top, out of which comes 3 or 4 Heads together, each on long Stalks, in March and April. I have sound this in Bellsize Grove near Hampstead, and others the like Places.

rne like Places.

356 Pluknets dry curled Glass moss. Muscus Polytrichoides pellucidus seré, fol. denticulatis ad margines veluticrispis, Ray 87n 240 Pluk 68. f. 8. Ox B 630 2. T 6. f These transparent L. curl and quickly sade; at the Endos each single Sprig comes one Stalk, ending in a long crooked Cap and Head. Found in divers shady Places about Hampstead. Heads in No-

veml'er.

257 Rofe Glass-mols. Mufeus parvus flellaris, CB 361. xi. Ray 113.4. Vol 3. p. 39.7. Syn ed 35.7. Ox B 630.9. 1 7. f 1. f 7. opt. Jonft. 1563. f 14. Park. 1308. f 9. Lob. Ic 2, p. 243.

1. B 179. f. Muscus electus fol. in ordern fparsis, Flor. Prust. 168. pl 457. f 47. This is known by its Rosey Tops, its Heads somewhat 6.00ked, on Inch Stalks. In most Heaths and low Woods

Woods.

358 Upright Oval Glafs-mess. Muscus tricheides fol. Seignth roundis, erectus major, R.3. p 43. pl 3. Syn ed 338. pl 6. Ox B 627. 39. T 6. 11. fult. Its upright and larger than the next. Grows in Bogs and Moors, &c. in Ireland,

Wales and elewhere.

350 Ray's Oval Glass-moss. Museus fol. paucis subrotundis pellucidis, Nob Ox B 627. 40. T 6. s. f 1. opt. Pluk. 47. f 7. Adiantum aureum humilius, fol. latis subrotundis, Ray Cat. Angl. & Hist. 124.5. Vol. 3. p.40.1. Syn. 19. 5. ed 35.

1. Its Leaves oval and thin set. Grows in thady Lanes and moift low Woods, &c.

360 Pluknets tound Glass moss. Muscus trichomanoides Supimis nofter elegans, Pluk. Algmagest. Ox. B. 627. 41. T 6. f 2. f 2. opt. Its commonly brancht with thin set round al-

ternate Leaves. In moist Ditches and shady Places.

361 Losels round Glass moss. Musius Trichomanoides, fol. roundioribus pellucidis, squamatim conjuncte sibi incumbentibus, BOx 627.42. Tab 6. 12. 13. Losel. 167. pl.456. 145. opt. These Leaves are round, cles let and often brancht. I have met with this in moist shady Places,

362 Cornwall Glefs-mois. Mufcus aq. Cornubienfis, plurimum capillaceus, foliolis exiguis alternis per totam capillorum laugitudinem adnatis, Bob Ox 627 48. Its Leaves very small, roundish, alternately set and thin First discovered by the Re-

verend Mr. Stevens in Dartmore Lake, Cornwall.

363 Lhwyds purple Glass moss. Musus Trichomanoides purpureas, Alpinis rivulis innascens, R.S. ed 41. 1. Ox B627. 48. var. T 6.1 3. fult. Found by Mr Lhwyd in the Rivulets on the highest Mountains in Wales.

364 Bobart's great Feathered Glass-moss. Muscus pennatus capitulis Adiani, Bob Ox 626, 36. T 6. s 1. f 5. R.S 236. qr? ed P 35. z. Found in mady moist Woods, &c. Heads in December

and January.

365 Rays small Feathered Glass moss. Muscus Filicifolius f. pennatus minor, pinnulis plurimis ad mediam costam latiusculis crebris, Ray 3. p 40.6. Syn ed 35.3. Ox B 627. 37. Like the last, but in all Paris less. On the Banks of shady Ditches

in Woods, &c. Heads in Dec. and Jan.

366 Vaillants Pin-feathered Glass-moss. Muscus pennatus omnium minimus, Instit. 556, Muscus Polytrichoides perexiguus, capitulis in summis surculis s. foliis subrotundis erectis, Ray 3. p 40 7. Syn ed 35.4. Smaller than the Two last, and particularly differs in bearing its Head at the Top of its Leaves. Heads with the last, and grows in the like Places.

SECT. V. Musci Trichoid s, capitulis ferè eredis.

Pike Mosses.

367 Bobarts Hearth-moss. Adiantum aureum medium, fol-tenuissimis, capit. erectis acutis, Ray Syn 237. Bob Ox 630. xi.

T 7. 1 1. f xi. opt Polytrichum aureum medium, Ray 124. 3. è fent. Bobart. Inst. 551.6. Common Bistops Wood, Hampsted. 368 Doodyes long-headed Moss. Muscus trichoides minus, capitulus longissims, Ray 3. p 34.4. Syn 243. ed 29. 3. Ox B 631.13. T 7. f 1. f 13. Inst 551.7. Its very long Heads for to small a Moss diff is it from others. On the dry sandy Bonks about Mather Hist. See Hampsted. Heads in February and the sandy and the sandy Banks, about Mother Huffs, &c. Hampsted. Heads in Febr. and March

369 Our Extinguisher Moss. Muscus trichoides pileis magnis acutis, Mus, noit, 89, sig. Ray 3, p 37.24. Syn ed 32.24. Ox B 630.12. T 7. s 1. f 12. Inst. 552. Its large long coned Cap differs it from others. I this observed this on the old Flint Wall in the Road near St. Albans. Heads in Decemb.

370 Dr. Richardsons hook leaved Bog-moss. Muscus trichoides palustris, capit. erectis, fol. reflexis, Ray Syn ed 338.

Its also found both in Ireland and Wales.


371 Common Wall Pike moss. Muscus trich. parvus, fol. Musici vulg. capit. longis acutis, Doody Ray Syn 243. pl 5. ed 28.1. Hist. V.3. p 34. pl 1. Ox B 628.4. T 6. 13. f qr? id. 628.1. T 6. f. lts Leaves are mixed with white Hairs. Very common on all the Brick Walls about London. Heads in Dec. Jan. &c.

372 Bobarts Wall Pike-moss. Muscus tectorum densis cel-372 Bodarts Wall Pike-mois. Mujeus tectorum densis celpitibus, eapit. oblongis, fol. latiusculis in pilum desinentibus, Bob Ox 028. 2. T 6. 13. f. Leaves broader and roundish, with pointed Heads, bus not so crooked as the last. 373 Mr. Dyers Devon-moss. Museus trithoides humilis ramotus, eapitulis oblongis, tumidiusculis donatus, Bob Ox 628.3. T 6.13. f This differs from the foregoing, in being branched

branched.

374 Bobarts Rofe House-moss. Museus muralis minimus Roleus, seu stellaris, capit. longiusculis acutis erectis, Bob Ox 629.12. T 6. 1 3. f Each Sprig ends in a Rose. Grows scarce half an Inch high, on Walls and Medge sides plenticular

375 Doody's Globe moss. Muscus trich. medius capitulis sphericis Ray Syn 243. pl 2. cd 30. xi. Ox B 628. 6. T 6 13. f opt. Adiantum aureum minus, capitulis rotundis Bobaris, Ray Syn 237. Init. 551. An elegant Mois, and pretty com-mon on the landy Banks in the Lane from the Spaniards at Park-Gate towards Highgate. Heads in March.



Botanicum Anglicum,

The English Herball:

Wherein is contained a curious Collection of Real Plants, being the true Patterns of fuch Trees, Struls, and Herbs, as are observed to grow Wild in England.

By which any one may most casily attain to the Speedy and True Knowledge of them.

With an Account (affixed to each Plant) of their Names, Places where Growing, and Times of Flourishing: As alfowhat Parts and Preparations, of each Phylical Plant, are most in Use.

And for the farther Instruction and Satisfaction of fuch, who are Lovers of Plants. The Composer of this Collection, chose to make his chiefest Reserences to the General History Catalogue and Synopsis of that Leanned Author, and most Judicious Botanist, Mr. John Ray: As also, to our Two most Esteemed English Herballs. Johnson upon Gerard and Parkinson; and for your more speedy finding each PLANT; he hath quoted the Page, wherein you may observe its Name, Figure, or Description.

An Explication of the Words and Authors which are here Abbreviated.

C. B. Prodr. & Pin. Casper Bauhine his Prodromus & Pinax, both Printed at Basil. A. D. 1671. Quarto.

J. B. Joannis Bauhini Plantarum Historia Universalis Ebrodun. 1650. 3. Vol. Fol.

Cat. Cant. Mr. Rey's Catalogue of Cambridge-Plants 1660. Octavo.

Ger. Gerards Herbal, Corrected, and Enlarged by Mr.

Thomas Johnson an Apothecary, Lond. 1636. Fol. L. Pl. Phys Leonardi Plucknetij Phytographia. This most compendious Herbal confifts or near 800 new and unheard of Trees, Strubs, and Herbs; not only of European growth, but from both the Indies, &c. all which are very curroufly Engraven on Copper Plates; with fuch adapt Names from their Proper Characteristicks as are truly descriptive of the things themselves. Lond.

Mer. Pin. Dr. Marret's Pinax, being a fuccinct Account of what Protables, Animals, Feffills, &c. hath been observed to be Natives of England, Lond. 1667. 800.

Offic. Shows you by what Name that Plant is generally Prescribed by the Learned Colledge of Physicians, and

known in the Apothecaries Shops.

Park. Parkinfon's Theatrum Botanicum : or, Theatre of Plants Lond. 1640. Pol.

Park. Parad. The same Authors Paradise, or Garden of Flowers. Lond. 1619. Fol.

Phyt. Brit. Phytologia Britannica, being an Alphabeti-

cal Catalogue of English Plants, 1650. 8vo.

Prep. & Usu. Denotes what Parts and Preparations of that Plant are most frequently used.

Raij. Cat. Angl. Mr. Ray's Catalogue of English Plants! Lond. 1677. 800.

Raij. Fascic. Fasciculus Stirpium Britannicarum, being a Supplement to his English Catalogue. Lond. 1688. 8vo.

Raij. H. Pl. Historia Plantarum Raij, being an Univerfal History of Plants, digested in a Methodical manner; with the Characteristick Notes of each different Genus, and most Acurate Descriptions of every Plant: As also, an ample Account of their Use and Vertucs. Lond. 1686. 2 Vol. Fol.

Raij. Synops. Synopsis Methodica Stirpium Britannicarum. Wherein is contained an account of what Plants hath hitherto been observed to grow Wild in England, compofed after the fame Method with his General History: A Book very inseful to such as are desirous to know Our Native English Plants. Lond. 1690. 8vo.

BOTANICUM ANGLICUM; |

Tipe English herball:

W' erein is contained a curious Collection of Real Plants, being the true Patterns of fuel Trees, Shrubs, and Hrbs, as are observed to grow Wild in England.

Wi han Account affixed to each Plunt, of their Names, Places where Growing, and Times of Flourishi.g. As also, what Parts and Preparations of each .
Intofact Diant the most in Use.

Sold by Samuel Smith, at the Princes-Arms in St. Taul's Church-yard. Louden.

Fig. 14.0 Ray Cot. Appl. St H. H. 1277. Synop. 19. It Flowers in Acrol. and Appl. and to Fruit is 147 a bout Michael note, is grows in Woods and Thought The Soils or well Scrope Ire.

Hortus Siccus Pharmaceuticus:

A Physick Garden of Medicinal Herbs.

Wherein is contained a curious Collection of Real Plants, being the true Patterns of such Trees, Shrubs, and Herbs, both Nivive and Foreign, as are most frequently Prescribed by the Learned Colledge of Physicians, and daily Used in the Apothecaries

With an Account (affixed to each Plant) of their Names, Places where Growing, and Times of Flourishing; As also, what Parts, and Preparations of each Plant are most Use.

Sold by Samuel Smith, at the Princes Arms in St. Paul's Church-Yard. London.

Trifolium nodiflorum glomerulis mollioribus & roun lioribus femine magno Raij Cat. Angl. parvum hir 10t un floribus parvis dilute purpureis in glomerulis molloribus & oblongis semine magno Ejusa. Synops. 134. 11.11. 945. It Flowers and Seeds in July, &c. in barren Pattures. Knotted Trefeil, with foft Heads, and large Seeds. Alsine parva palustris tricoccos Portulaca aquatica similis Raij Synops. 140. & H.Pl. 1035. Alsine formis paludosa tricarpos flosculis albis inapertis. L. Pl. Phys. Tab. 7. Fig. 5. It Flowers and Seeds in March, April, &c. in moist places, and sometimes amongst Corn. Blinks.

Anagallis lutea Ger. Fig. 618. Fl. luteo Park. Fig. 558. Raij. Cat. Angl. & H. Pl. 1024. Synops. 144. It Flowers about Midsummer in shady moist Woods. Tellow Pringernel.

Ascyrum supinum villosum palustre Park. 574. supinum ελωθες Clusii scu 2. Ger. 542. Raij Cat. Angl. & H Pl. 1020. Synops. 144. It Flowreth about July in moorish and boggy places. Hoary St. Peters-wort.

Bursa Pastoris minor. Park. Fig. 866. minima Ger. Fig. 276. & Nasturtium petraum Ejusd. Fig. 251. & Park. Fig. 828. Rais Cat. Angl. & H. Pl. 827. Synops. 116. It Flowers and Seeds in April and May on Rocks, and Sandy Banks. The lesser Spepheards-Purse.

Cauda Muris Ger. Fig 426. Myosuros Cauda Muris Park. Fig. 500. Raij Cai. Angl. & H. Pl. 1332. Synops. 209. It Spikes in April and May near Highways, and amongst Corn. Monse-tail.

Gramen parvum præcok spicâ lakâ canescente Raij Cat. Angl. 150. & Synops. 191. L. Pl. Phyt. Tab. 33. Fig. 9. It Flourishes in the Spring in barren and gravelly grounds. Pt. Kap's small Vernal Grass with a soft loose Spike.

Juncus parvus palustris cum parvis capitulis Equiteti Raij H. Pl. 1306. & Synops 203. L. Pl. Phyt. Tab. 40. Fig. 6. It Flowers in May, on Heaths and Boggy Places. Dw. wf Rush with small Aglet-Heads.

Lichen parvus crectus foliolis profunde laciniatis L. Pl. Phyt. Fig. 2. Tab 42. It was first observed by that most Ingenious and Skillful Botanist Dr. Plukenet in a Boggy rill of Water, near Dullidge-Wells, where it grows plentifully. Dr. Plukenet's small upright Liverwert, with deep Jagged Leaves.

Lysimachia galericulata minor Raij Cat. Argl. & H. Pl. 572. Synops. 84. Gratiola latisolia Ger. Fig. 581. latisolia seu nostras minor Park. Fig. 220. It Flowereth in July and August in Boggy Places. Hooded Loose-strife.

Muscus marinus denticulatus minor ramulis in creberrima capillamenta sparsis L. Pl. Phys. Tab. 48. Fig. 3. It's frequently found cast on the Sea-Shoar. Bushey Teathered Sea-Moss.

Nummularia flore purpurascente Ger. Fig. 635. minor fl. purp. Park. Fig. 555. Raij. Cat. Angl. & H. Pl. 1100. Synops. 161. It Flowers in fully in moist Boggy Grounds. Purple Flowered Money-wort.

Plantaginella pallustris C. B. Pin. 190 Raij. Cat. Angl. & H. Pl. 1077. Synops. 158. Alfine palustris repens soliis lanceolatis floribus albis perexiguis L. Pl. Phys. Fig 4. Tab. 74. It Flowers and Seeds in May, in dry Ponds and muddy Lanes. Water-Plantain Chickweed.

Sagitta aquatica omnium minima Raij. Syneps. Append. 242. Observed by Dr. Plukenet to grow in the Thames, by Lambeth-Bridge, and Peterborough-House, Flowering in June. The least Arrow-head.

Saxifraga graminea pusilla slore parvo tetropetalo Raij. H. Pl. 1026. & Synops. 145. Anglicana Alsine solia Ger. Fig. 567. Anglica Occidentalium Lobelis Park. 1 ig. 437. & Caryophyllus, minimus muscosus nostras Fig. 1. 134. It Flowers and Seeds great part of the Summer in Garden Walks, and moist Barren Pastures. Chiekwich Break-Stone or Pearl-Wort.

Speculum Veneris minus Ger. Fig. 439. Park. 1331. Raij Cat. Angl. & H. Pl. 748. Syneps. 104. It Flowers in May, amongst Corn. The lefter Corn-Violet or Venus Looking Glass.

Tithymalus leptophyllos Paris Fig. 193. Estila exigua Tragi Ger. Fig. 503. Ruj. Cat. zingl. 101. H. Pl. 828. Syrops. 125. It Giovers and Seeds in June, &c. amongst Corn. Dwarf comuni Spurge.

Fig. lops bromoide, Ger. Fig. 77. bromoides Belgarum Park. Fig. 1148. Raij. Cat. Angl. 107. 80 H. Pl. 1254. Synopf. 179. It Flourishes with the common Octs amongst Corn. Bearded Hill Oats, or Haver.

Anagallis aquatica rotunditolia Ger. Fig. 620. aq. 3. Lob. tol. fubroundo non crenato Park. Fig. 1237. Rasj Cat. Angl. 105. & H. Pl. 1101. Symppf. 162. It Flowers and Seeds in June and July, in Boggy Grounds. Round leav'd Water Fungernel.

Anemone nemorum alba Ger. Fig. 383. Ranunculus nemorofus albus fimplex Park. Fig. 325. Raij Cat. Angl. & H. Pl. 624. Synopfig3. It Flowers in March and April in moilt low Woods. Wood 21 emony.

Aftragalus sylvaticus Ger. Fig. 1237. Lathyrus sylve-stris lignolior Park. Fig. 1062. Raij Cat. Angl. & II. Pl. 916. Ernor. 131. It Placers and Peds in April and May, in thady Wood. Wood Peof.

Cardamine impatiens altera hirfutior Raij Cat. Angl. & H. Fl. 815. Stroops, 114. It Flowers in April and May, in Ditches and moist sandy Places. The lesser hairy Ladies Smock.

Carvophylin pratentis Ger. Fig. 594, prat. noster major Park. 1330 Rais Cir. Angl. 27., & H. Pt. 991. Synops. 139. It Flowers in June and July, in Sandy and Gravelly Grounds. Depiserd Pink.

Centaurium luteum perfoliatum C. B. Fin. parvum luteum Lob. Ger. Fig. 547. minus luteum vulgare Park. Fig. 272. Reij H. Pl. 1093. Synopf. 160. In Challey and Flowntainous Places, Flowering in July and Laguft. Tellow Centaury.

Filix palustris seu aquatica Raij Cat. Angl. & minor palustris Ejusd. H. Pl. 146. & Synops. 24. Dryopteris Pena: & Lob. Ger. Fig. 1135. It's sound in Boggy and Moorish Places. Alarsh Ferm.

Fumaria alba latifolia claviculata Ger. Fig. 1088 latifolia alba Park. Fig. 288. Raij Cet. Angl. & H. Fl. 974. Synops. 137. It Floriers and Pods in June, July, &c. on Hedges and the Banks of Rivers. Climbing Fumitory.


Gramen hirfotum elatius paniculà juncea compacta Ran Car. Lingl. 146 & H. Pl. 1201. Streep. 193. an Gr. nemorum hirfut un mains angultaolium Park. Fig. 1185. It Thuriffes in May in mi ist Ditches near Hedges and Woods. Close headed harry Grafi.

Gramen nemorosum hirsutum latisolium maximum Rais Cat. Angl. 156. & H. Pl. 1292. Synops. 133. an Gr. nemorum hirsutum latisolium minus sunceà paniculà Park. 1185. Whose Figure very well operes with this Plant. It Flurishes in Lipril and Mar, in thady Woods and on Rock. The greatest bread have I ba. y Wood-Grass.

Heileborine valustris nostras Roif Cat. Logl. & H. Pl. 1221. Synops. 178 It Flowers in Jone and July, in Boggy Places. Marsh Hellwers.

Pentaphyllum rectum minus Pak. 400. Cuinque-folium Tormentille facles les. Per 988. Raij Cat. Anel. 8:11. Pl \$13. 85 cm ges. le elever in Barron Sendy Patitutes. Tormente Ca jusfail.

Peplus sen Esula rotunda Ger. Fig. 503, Esula rotunda sen Peplus Park. Fig. 194. Raij Cat. Angl. & H. Pl. 869. Synops. 125. It Flourishes from Midjummer till Autumn, in Gardens amongst the Weeds. Petty Spurge.



The transfer of the second of

Hieracium Dentis Leonis folio hirsutum Ger. Fig. 303. foliis & floribus Dentis Leonis bulbosi Park. Fig. 788. Dens Leonis hirsutus rationale, Hieracium dictus Raij Synops. 46. Cat. Angl. 160. & H. Fl. 245. It Flowers in May and June, very frequently in Pastures and Meadows. Kough Danielion, commonly called Dandelion Dawsweed.

Hieracium fruticosum latisosium hirsutum Park. 802. Raij Cat. Angl. & H. Pl. 238. Synops 44. about High-gate and Hamstead, and elsewhere in Eushy Woods: Flowring in July and Angust. Scool rough leabed Bully Pawaweed.

Hieracium fruticosum augustisolium majus Park. Fig. 801. Intybaceum Ger. Fig. 298. Raij Cat. Angl. & H. Pl. 238. Synops. 44. In stony, gravelly, and shrubby places: Flowring in July, &c. Parrew leabed bushy Hawkweed.

Hieracium luteum glabrum, seu minus hirsutum I.B. Tom. 2. lib. 24. p. 1029. Cichoreum pratense luteum levius Park. Fig. 777. an potius Hieracium aj hacoides Ger. Fig. 297? soliis & facie Chrondrillæ Lobelii Park Fig. 794? Raij Cat. Angl. & H. Pl. 234. Synops. 43. It Flowers from Midsummer till August, in Pastures, and on Ditch banks. Sinooth Success Baubwied.

Hieracium minus præmorta radice Park. Fig. 794. minus scu Leporinum Ger. Fig. 296. Ranj Cat. Angl. & H. Pl. 230. Synops. 42. It's very frequent in Meadows and Pasture-Grounds: Flowring in fuly and August. Pellow penils-bit, or Pawames with bitter Roots.

Hieracium longius radicatum Ger. Fig. 298. Park. Fig. 790. Raij. Cat. Angl. & H. Pl. 230. Synops. 42. It Flowers from Midsummer till Autumn, very common in Meadows and Paftures. Long rooted Park need.

Oenanthe Cicutæ facie Lobelii Park. Fig. 894. Filipendula Cicutæ facie Ger. Fig. 1059. Rasj Cat. Angl. & H. Pl. 441. Synopf. 66. It Flowers between Midsummer and August in Rivers and simal running Waters, but not very frequent. Deniet Dropwest.

Orchis five Cynosorchis purpurea spica congesta pyramidali Cat. Cant. Raij Cat. Angl. & H. Pl. 1215. Synops. 170. an Cynosorchis militaris spica rubente conglomerato Park. 1345? & C.B. prodr. 28? It Flowers in July in dry and chalky Grounds. Burple late flowing Dechis.

Orchis palmata minor fl. rubro Park. Fig. 1358. Serapias minor nitente flore Ger. Fig. 222. Raij Cas. Angl. & H. Pl. 1224. Synopf. 173. It Flowers in June in wet Pastures and Meadows. Med banded Dathis, with long Spurs.

Orchis palmata fl. luteo viridi Raij H. Pl. 1224 Synops. 173. vid. Cat. Lingl. 217 an palmata fl. viridi C. B. prodr. 30? & Park. 1359? Scrapias Batrachites altera Ger. Fig. 224? It Flowers about the end of May in dry Grounds. Banded Dichis with a green flower, called by some the frog Dichis.

Orchis hermaphroditica Ger. Fig. 211. Orchis serapias bifolia vel trifolia minor Park. Fig. 1350. Raij Cat. Angl. & H.
Pl. 1221. Synops. 172. It Flowers in July in Pasture Grounds
and low Woods. Buttersty Satyrion.

Ornithopodium minus Ger. Fig. 1241. Park. Fig. 1092. Raij Cat. Angl. & H. Pl. 931. Synopf. 132. It Flowers from June till September in Sandy Heaths and Gravelly Grounds. Small Bitosolost.

Pecten Veneris seu Scandix Ger. Fig. 1040. Scandix vulg. sive Pecten Veneris Park. Fig. 916. Rais Cat. Angl. & H. Pl. 428. Synops. 64. It Flowers in May amongst Com. Menus Comb, or Shepherds Reedle.

Pedicularis five Crista Galli lutea Park. Fig. 713. Crista Galli Ger. Fig. 1071. Raij Cat. Angl. & H. Pl. 769. Synops. 106. It Flowers from Midsummer till August in dry Paitures and Meadows. Pellow Kattle, or Cochs-Comb.

Pedicularis palustris rubra elatior Raij Cat. Angl. & H. Pl. 770. Synops. 106. It Flowers in May and June in Marshy and Boggy places. Ray's Breat Parth red Kattle, or Lousewort.

Pedicularis Ger. Fig. 1071. pratensis rubra vulgaris Park. Fig. 713. Raij Cat. Angl. & H. Pl. 770. Synops. 106. It Flowers in May and June in moist Meadows and Moorish Heaths. Red-Kettle.

Persicaria Salicis solio Potamogiton angustisolium dicta Raij H. Pl. 184 & Synops. 32. Cat. Angl. 241. Potamogeiton angustisolium Ger. Fig. 821. Fontalis major longisolia Park, Fig. 1254. It Flowers in June and July in Ponds and large Ditches frequently. Parsow leaved grandweed or Arsmatt.

Plantago aquatica major Ger. Fig. 417. Park. Fig. 1245. Raif Cat. Angl. & H. Pl. 618. Synopf. 92. It Flowers from June till August in small Rivers and Ponds frequently. Great Emater-Plantane.

Plantago aquatica humilis Ger. Fig. 417. aq. minor Park. Fig. 1245. Raij Cat. Angl. & H. Pl. 618. Synopf. 92. It grows in the same places with the greater of this kind, but is not so common, The Lesser Mater-Plantane.

Polygala Ger. Fig. 563. minor Park. Fig. 1332. Raij Cat. Angl. & H. Pl. 1335. Synops. 209. It Flowers in May and June on dry and hilly Pattures. Spils-most

Polygonum Germanicum sive Knawel Germanorum Park. Fig. 447. selenoides sive Knawel Ger. Fig. 566. Kaij Cat. Angl. & H. Pl. 213: Synops. 40. It Florishes about Midsummer in Sandy barren places. Knawel, or German Knotgtass.

Populus alba Ger. Fig. 1486. Parki Fig. 1410. Raif Cat: Angl. & H. Pl. 1418. Synopf. 214. It beareth its Jules or Catkins in April and May in wet Woods, and by Water-fides. White Poplar, or Abele-Tiee.

Populus alba foliis minoribus Ger. Fig. 1487. Park. Fig. 1410: Raij Cat. Angl. & H. Pl. 1419. Synopf. 215. It is in such places as the greater kind. The Smaller leabed Tablite Poplar.

Populus Lybica Ger. Fig. 1487. Park. Fig. 1411. Raij Cat. Angl. & H. Pl. 1419. Synopf. 215. Frequently amongst Birch-Trees in moist Woods, bearing its Catkins in the Spring. The Asp, or Tremblings popular.

Potamogeiton latifolium Ger. Fig. 821. Fontalis major latifolia vulgaris Park. Fig. 1254. Raij Cat. Angl. & H. Pl. 188. Synopf 34. It Flowers and Seeds in June and July in Fish-Ponds and standing Waters. Legond leaded Posidenced.

Potamogeiton aquis immersum solio pellucido lato oblongo acuto Raij H. Pl. 188. & Synops. 34. Cat. Angl. an longis acutis soliis Ger. Fig. 822? It Flowers in June, in large slow running Waters. Long leabed pellucid Pond weed.

Potamogeiton perfoliatum Phyt. Brit. & Raij H. Pl. 188. Syn. 34. & Cat. Angl. Pot. 3. Dodonei Ger. Fig. 822. It Flowers at the same time with the others in Ponds and slow running Rivers. Perfoliate Pond-weed.

Potamogeito affine gramen aquaticum Cat. Cant. Raij Cat. Angl. & H. Pl. 190. Synopf. 35. In Ponds and standing Waters. Ray's mater-grais, with small crooked Pods.

Pscudomelanthinum Ger. Fig. 1087. Lychnoides segetum sive Nigellastrum Purk. Fig. 632. Raij Cat. Angl. & H.Pl. 998. Synops 139 It Flowers in June and July, too frequently amongst Com. Curis.

Pulmonaria Gallica five aurea latifolia Ger. Fig. 304. Raij Cat. Angl. & H. Pl. 239. Synopl. 44. It Flowers in June and July, on old Walls, shady Banks, and in Woods. French or Golden Lunguett.

Ranunculus aqua'ilis omnino tenuifolius I. B. Millefolium five Maratriphylion flore & semine Ranunculi aquatici Hepaticæ facie Ger. Fig. 827. Millefol aquat. Ranunculi fl. & capitulo Park. Fig. 1256. Raij Cat. Angl. 200. & H. Pl. 586. Synops. 87. It Flowers in April and May in shallow Ponds and Ditches. Fine seased destates Crowfoot.

Ranunculus aquatilis Ger. Fig. 829. aq. Hepaticæ facie Park. Fig. 1216. Raij Car. Angl. & H. Pl. 586. Synops. 87. It Flowers in April and May frequently in standing muddy Waters. Water Crowfoot with differing leaves.

Ranunculus palustris rotundisolius Ger. Fig. 962. palustris Sardonius lævis Park. Fig. 1215. Rais Cat. Angl. & H. Pl. 585. Synops. 87. It Flowers in May, June, &c. and is very common in all Watry Ditches. Kound leaded Mater Trowsoot.

Ranunculus flammeus serratus Ger. Fig. 962. Park, Fig. 1215. Raif Cat. Angl. & H. Pl. 587. Synops. 88. It Flowers in June in Watry and Marshy Meadows. Indented Spearwort.

Ranunculus hederaceus rivulorum se extendens atrâ maculâ notatus I. B. aquatilis varietas altera Ger. 830. Raij Cat. Angl. & H. Pl. 586. Synops. 87. It Flowers from May till Mid-summer in Ditches and shallow sandy Rivulets. The isabed scient-Crowsot.

Ranunculus bulbosus Ger. Fig. 953. Park. Fig. 329. Raij Cat. Angl. & H. Pl. 381. Synops. 86. It's very common in Meadows, flowring all the Summer. Mound-rooted, or Bulbous Crewfoot.

Ranunculus pratensis repens Park. Fig. 328. prat. etiamque hortensis Ger. Fig. 951. Raij Cat. Angl. & H. Pl. 581. Synops. 86. It Flowers from May till August in moist Meadows. Common creeping Crowfoot, or Butters Cups.

Ranunculus pratensis erectus acris vulg. Park. Fig. 328. surrectis cauliculis Ger. Fig. 951. Raij Cat. Angl. & H. Pl. 583. Synops. 86. It's very common in moist Meadows, flowring nost part of the Summer. Apright speadow-Crowsoot.

Raphanus aquaticus Ger. Fig. 240. aq. Lobelii Park. Fig. 1228. both these Figures are ill expressed. Raij Cat. Angl. & H. Pl. 818. It Flowers in June and July in Watry Ditches, and muddy places.

Rapistrum arvorum Ger. Fig. 233. Park. Fig. 862. Raij Cat. Angl. & H. Pl. 802. Synopf. 111. Amongst Corn too frequently, flowring most part of the Summer. Pellow-Charlock, a caile-sputate.

Rapistrum album articulatum Park. Fig. 863. Raphanus lylv. Ger. Fig. 240. Raij Cat. Angl. & H. Pl. 805. Synops. 111. It Flowers about Midsummer plentifully amongst Corn. Withite source Charlock, with joynted Cods.

Rapistrum luteum siliquâ glabrà articulatà Cat. Cant. desc. Raij Cat. Angl. & H. Pl. 805. Synops. 112. It Flowers at the same time with the other kinds amongst Corn. Pellow Charlet, with a smooth joynted Cod.

Rubeola arvensis cærulea repens C. B. prodr. 145. Rubia minor pratensis cærulea Park. 276. Raij Cat. Angl. & H. Pl. 483. Synops. 74. It Flowers in June, &c. in Untilled or Falow Fields. Little Kield Madder.

Salix Ger. Fig. 1389. arborea angustisolia vulgaris Park. Fig. 1430. Raij Cat. Angl. & H. Pl. 1419. Synops. 215. It Flowers early, before the Leaves appear. The most common mbite millow.

Salix Caprea rorundifolia Ger. Fig. 1390. latifolia rotunda Park. Fig. 1432. Raij Cat. Angl. & H. Pl. 1422. Synopf. 216. It prows in Hedges, flowring in the Spring. Mound leabed pallow.

Salix Caprea latifolia Ger. Fig. 1390. latifolia oblongior ark. Fig. 1432. Raij Cat. Angl. & H. Pl. 1422. Synops. 217. in Watry Hedges frequently. Common Sallow.

Salix folio longissimo Cat. Cant. Raij Cat. Angl. & H. Pl. 1423. Synops. 217. In moist Grounds, and by River-sides. the Diet.

Sambucus aquatilis sive palustris Ger. Fig. 1424. palust. sive 14. Park Fig. 209. Raij Cat. Angl. & H. Pl. 1586. Synops. 225. I Flowers in May, and the Berries are ripe in Antumn in moist Hedges and wet Woods.

Sorbus sylvestris sive Fraxinus bubula Ger. Fig. 1473. Ornus sive Fraxinus sylv. Park. Fig. 1419. Raij Cat. Angl. & H.Pl. 1457. Synops. 219. It Flowers in May, and the Fruit is ripe in Septemb, in Marshy places and Hilly Woods. Duicken-Lee.

Spergula I. B. Saginæ Spergula Ger. Fig. 1125. Saginæ Spergula major Park. Fig. 562. Raij Cat. Angl. & H. Pl. 1034. Synops. 148. It Flowers in June and July in Corn-Fields. Spurrey.

Spergula purpurea I. B. Alfineformis five 5 Ger. 1125. Sagiræ Spergula minima Park. 561. Raij Cat. Angl. & H. Pl. 1034. Synoof. 148. It Flowers in May and June in fandy and gravelly Grounds. Surple Spurrey.

Taxus Ger. Fig. 1370. Park. Fig. 1412. Raij Cat. Angl & H. Pl. 1416. Synopj 214. It Flowers in the Spring, and the Fruit is ripe from Autumn till Christmas, in Stoney and Mountainous Woods and Hedges in the West and Southern parts of this Land plentifully. The Devi-Liee.

Thalictrum majus vulgare Park. Fig. 264. Thalietrum five, Thalictrum majus Ger. Fig. 125. Raij Cat. Angl. & H. Pl. 403. Synopf. 62. It Flowers in July and August near Rivers, and in moist Meadows. Deadow-Kie.

Tithymalus Helioscopius Ger. Fig. 498. Park. Fig. 189. Raij Cat. Angl. & H. Pl. 869. Synops. 125. It Flowers and Seeds about Midsummer in Kitchin-Gardens and wast Grounds. Sun-Spunge, or Wasterwort.

Tragopogon luteum Ger. Fig. 735. Park. parad. 514. Raij Cat. Angl. & H. Pl. 252. Synops. 46. It Flowers in June and July in Meadows and Fertile Paltures frequently. Pellom Coates beato, or Go to Bed at Boon.

Tribulus aquaticus minor Muscatellæ floribus Ger. Fig. 824. aq. minor alter Park. Fig. 1248. Raij Cat. Angl. & H. Pl. 189. Synops. 34. It Flowers in June and July in Rivers and and Ditches of clear Water. Small Frogs-Lettice.

Tribulus aq. minor Quercûs floribus Ger. Fig. 824. aq. minor prior, Park. Fig. 1248. Raij Cat. Angl. & H. Pl. 189. Synops. 34. It Flowers in June and July in muddy Waters, as Ponds, and slow running Rivers. Curled Frogs-Lettice.

Trisolium Fragiserum Ger. Fig. 1208. Frag. Frisicum Park, Fig. 1109. Raij Cat. Angl. & H. Pl. 946. Synops. 135. It Flowers in July and August in Clay Pastures, where the Water hath stood all the Winter. Strawberry Liesoil.

Trisolium luteum Lupinum Ger. Fig. 1186. Raij Cat. Angl. & H. Pl. 949. Synops. 135. It Flowers and Seeds in June and July in Corn-Fields and Sandy-Grounds. Depontresoil.

Trisolium Lupulinum alterum minus Raij Cat. Angl. & H. Pl. 949. Synops. 135. It Flowers about Midsummer in Meadows amongst the Grass. The Lesser Bop-Tresoil.

Trifolium luteum minimum Ger. Fig. 1186. Park; Fig. 1111. Raij Cat. Angl. & H.Pl. 952. Synopf. 135. It Flowers in June and July in Barren Pastures. Small Pellom or Melitat Exceosi.

Trisolium cordatum Ger. Fig. 1190. Medica Arabisa Camerarii sive Trisol. cordatum Park. Fig. 1115. Raij Cat. Angl. & H. Pl. 963. Synops. 137. It Flowers and Seeds in May and June in dry and sandy Pastures. Claber, or Deatt-Aresoil.

Trifolium filiquosum minus Ger. Fig. 1190. Raij Cat. Angl. & H. Pl. 967. Synops. 137. It Flowers in July and August, frequent in most Pastures. Smooth creeping codded Tresoil.

Trisolium corniculatum 3. Dod. Trisolii siliquosi varietas major Ger. 1190. Raij Cat. Angl. & H. Pl. 968. Synops. 137. It Flowers from June till August in moist Meadows and near Hedges. Poary birds-spot Lieseil.

Vaccinia nigra Ger. Fig. 1415. nigra vulg. Park. Fig. 1456. Raij Cat. Angl. & H. Pl. 1488. Synops. 222. It Flowers in May, and the Fruit is ripe in August, in Boggy Heaths and moist Woods. Black Emports, Embortle-berries, or Bil-herries.

B. Tall too

A tree eage will read with the Care Say Cat. With the second of the second The state of the s t elemente di E kan polsiti a di Lini Lan Ligazante Lann The state of the s The Late of the La 189-18.C. .; I 3 I ;. है हिन्द्र १ से AL PERSON Color of the state तिहरी हेडा दें ते पुरुष के बीचू भीचन स्टेंग हुन napens T 15 1 rg. ma. r inûrellark. 1995 - i it Flo M. the said. ❖ 31 3 5 F well it is cib mile et : 92 PL 172. C So the second second 311 3 41. LE F.D. e figsterlichen bis Fig. 1:40. The first of the William Field.

1. 1. 18. Line 19. Line 19. William Willia i continue de la cont L Theren 4129 C 12 1 3. T. eg & M 1250 Justin valg. Park eg & M 1250 Jp J. 180. Its Wall of the Walls of the state of the series of the ser Comparison and the Sections and the comparison of the comparison o 二十 天主 门门 ា មានក្រុ 11: 4 To the second of 心花园 The Mark to the second And e short किन्द्री किन्द्री केन्द्रीत । 10 (2% 5 . 22 313 y) . Distribution English Nount bay europad y . . 5 J. 71 . 20

Muscus ramosus floridus Ger. sig. 1560. Quercinus sæniculaceus Park. sig. 1312. Raij Cat. Angl. & H. Pl. 115. Synops. 14. On old Trees frequently. Eree-moss with Rundles.

Muscus arboreus ramosus I.B. Raig Cat. Angl. & H. Pl. 115. Synops. 13. Very common on most old Trees, and Pales. The most common, of flat branched Erree-Pols.

Muscus terrestris latioribus soliis major seu vulgaris Raij Cat. Angl. & H. Pl. 122. Synops. 17. In most shady places very frequently. The greater common Earth-Sposs.

Filix faxatilis ramosa maritima nostras Raij H. Pl. 151. Fa-scieul. & Synops. 23. On old Churches and Stone-Walls. Small branched Stone-Fern.

Equisetum nudum lævius nostras D. Doody Raij in Append. Synops. 244. an nudum Ger. sig. 1113? junceum sive nudum Park. sig. 1201? In Rivulets and Watry Ditches. 24-ke Posse-Lail.

Lactuca marina tubulosa Raij H. Pl. 77. Fascic. 14. & Symops. 6. In such Ditches and Rivers as the Sea-water flowerh into every Tide. Ray's Long bladered Dyner-green.

Fungus ramosus niger compressus parvus, apicibus albidis Raij Car. Angl. & H. Pl. 104. Synops. 10. It grows to the Roots of Trees and rotten Wood. Ray's Black bosned southout spushrome, with white tips.

Fungus 3 five Orbicularis Ger. 1582. F. ovatus 3 Park. 1324. Raij Cat. Angl. & H. Pl. 104. Synopf. 11. It's found in dry Meadows and Pastures in Autumn frequently. Duty Quiffrom, or Puff-Balls.

Fungus maximus rotundus pulverulentus, dictus Germanis Pfossis. I. B. Raij Car. Angl. & H. Pl. 105. Synops. 11. On Dunghils, and in Fertile Pastures. The larger out pushrom.

Fungus Campanisormis niger multa semina plana in se continens D. Merrer in pin. 41. Fungi calyciformes seminseri Mentzel. pugil. Fig. Tab. 6. It's sound Spring and Autumn in Rainy and uncertain Weather in shady places, growing to putrissed Wood. Seeding Cup Mustrom.

Fungus arborum & lignorum putrescentium coloris varii Raij Cat. Angl. & H. Pl. 109. Synops. 12. imbricatim dispositis versicolor Park. Fig. 1322. It's found growing to the Bodies of divers Trees. Party-coloured 20000/99usprom.

Fungus spongiosus niger reticulatus doliolis vinosis adnafeens D. Doody Raij Synops. 12. It's often sound in Wine-Cellars, sticking to the Casks. Doody's punk-like Mustrem.

Muscus Filicinus Ger. Fig. 1561. Park. Fig. 1310. Raij Cat. Angl. & H. Pl. 112. Synops. 13. It's common in shady Woods. Fern-Spots.

Muscus corniculatus Park, Fig. 1308, Raij Cat. Augl. & H. Pl. 113. Synops. 13. In barren Heaths. Dozned Doss.

Muscus multiformiter pyxidatus apicibus coccineis D. Plos. Hist. Nat. Stafford. cap. 6. p. 199. Fig. Tab. 14. Raij in Append. H. Pl. 1851. & Synops. 13. In Hilly and barren Heathy Grounds. Byrches Scatlet-headed Cup-Moss.

Muscus tubulosus ramosissimus fruticuli specie Raij Cat. Angl. & H. Pl. 113. Synops. 13. Muscus Corallinus sive Corallina montana Ger. Fig. 1572. Musc. Corallinus montanus Park. Fig. 1312. On Spungy and Boggy Heaths. Cozalline spots.

Lichen cinereus terrestris Raij Cat. Angl. & H. Pl. 117. Synops. 14. In dry Pastures frequently. Mr. Ray's Asp-colouted ground Liber-wort.

Lenticula aq. trifulca 1. B. C. B. Hederula aq. Ger. Fig. 830: Ranunculus Hederaceus aquaticus Park, Fig. 1216. Raij Car. Angl. & H. Pl. 118. Synopf. 15. In Fenny Ditches and muddy Waters. 3by leabed Duck-meat.

Muscus aq. terrestri vulgari similis sed major Raij Cat. Angl. & H. Pl. 122. Synops. 17. an Fontalis minor lucens I.B? In Rivers and standing Waters, growing to Stones or Wood.

Muscus palustris rerrestri similis Raij Cat. Angl. & H. Pl. 122. Synops. 18. terrestris vulgatissimus Park. Fig. 1306. In moist Boggy Heaths. Boggy Earth-Hoss.

Adianthum aureum humilius soliis latis subrotundis Raij Cat. Angl. & H. Pl. 124. Synops. 19. By Rills of Water, and in moist shady places. Kound leaved Golden Paiden-Bair.

Lichen seu Hepatica minor umbellata Park. Fig. 1314. Hepatica umbellata Ger. Fig. 1567. Raij Cat. Angl. & H. Pl. 127. Synops. 20. It Flowers in May and June, in moist, shady, and Boggy places. Kound headed Liber-Wort.

Lichen seu Hepatica minor Stellata Park, Fig. 1314. Hepatica Stellata Ger. Fig. 1565. Raij Cat. Angl. & H. Pl. 125. Synops. 20. It Flowers about Midsummer in Gravelly and Shady Walks, in Gardens and moist places. Star headed ground Libertweet.

Lichen alter minor folio calceato Park. Fig. 1314. Raij Cat. Angl. & H. Pl. 125. Synopf. 20. It Flowers and Seeds about May, near Rills of Water, and in moist shady places. Small Liverwort, with crumpled leaves.

Lichen seu Hepatica lunulata epiphyllocarpos D. Dale. Raij H. Pl. 125. Fascicul. & Synops. 20. It Flowers and Seeds in the Spring, on moist and shady Banks. Mr. Dale's spoon Liberwegt.

Lichen minimus foliolis laciniatis Raij H. Pl. 126 Fascicul. & Synops. 20. Near Rivulets, and in shady moist places. Mr. Ray's Jagged leabed Liberwort.

Equisetum 8 Ger. 1115. fætidum sub aqua repens Park. Fig. 1201. Raij Cat. Angl. & H. Pl. 130. Synops. 21. In silthy and muddy Waters, frequently from Midsummer till Autumn. Stinking water Bosse-Tail.

Equisetum majus Ger. Fig. 1113. majus palustre Park, Fig. 1200. Raij Cat. Angl. & H. Pl. 128. Synops. 20. It Flowers about Midsummer in wet and Marshy Grounds. Seest parts Dogle-Lail.

Equisetum omnium minus tenuisolium Park. Fig. 1201. Raij Cat. Angl. & H. Pl. 128. Synops. 20. In the low wet Grounds, and Woods about Highgate, and in other the like places. Finest leabed Posse-Lait.

Equisetum 7. sive Hippuris Coralloides Ger. 1115. Raij Cat. Angl. & H. Pl. 130. Synops. 21. It grows in wet Boggy places. Gerard's Cozalline Hogses Tail.

Adianthum nigrum vulgare Park. Fig. 1049. Onopteris mas Ger. Fig. 1137. Raij Cat. Angl. 92. & H. Pl. 152. Synopf. 26. On old Churches, and to the Roots of Trees in shadowy Woods. Common black Paiden-Pair.

Lonchitis aspera Ger. Fig. 1140. aspera minor Park. Fig. 1042. Raij Cat. Angl. & H. Pl. 138. Synops. 23. In Woods, Heaths, and moist places. Mough pricen-west.

Hordeum spurium Ger. Fig. 73. spurium vulg. Park. Fig. 1147. Raij Cat. Angl. & H. Pl. 1258. Synops. 180. It's very common on Mud-Walls, and by Path-ways, being in Spike most part of the Summer. Walls Barley, or rather will-

Gramen Secalinum Ger. 29: Secal majus Park. Fig. 1144. Raij Cat. Angl. & H. Pl. 1258. Synopf. 180. In Meadows, it Spikes in June and July. Peadow Kie Grafs.

Gramen Secalinum palustre & maritimum Rais H. Pl. 1258. Fascic. & Synops. 180. In Salt Marshes, frequently Spiking in June, July, &c. Mr. Ray's sparsh are Grass.

Spartum nostras parvum Lobelii Ger. in Append. Fig. 1631. parvum Batavicum & Anglicum Park. Fig. 1199. Raij Cat. Angl. & H. Pl. 1260 Syn. 180. It Spikes in Angust in Mountainous Heaths, and Boggy places.

Lolium rubrum Ger. Fig. 78. sive Phoenix Park. Fig. 1145. Raij Cat. Angl. & H. Pl. 1263. Synops. 181. It's very common in Pastures, Meadows, and near Path-ways, Spiking in May, June, &c. Red Darnel-Grass.

Lolium album Ger. Fig. 78. Park. Fig. 1145. Raij Cat. Angl. & H. Pl. 1262. Synops. 181. It Spikes in July and August amongst Corn, especially Wheat. Darnel, or May-Brais.

Gramen Alopecuroides majus Ger. Fig. 10. Gr. Phalaroides majus Park. Fig. 1164. Raij Cat. Angl. & H. Pl. 1264. Synopf. 182. It Spikes in April, May, &c. very common in most Meadows. Great For tail Grass.

Gramen fluviatile spicatum Ger. Fig. 14. aq. spicatum Park. Fig. 1275. Raij Cat. Angl. & H. Pl. 1264. Synops. 182. It Spikes about Midsummer, in Pools and Watry Ditches. Spikes Flote grass, or For-tail mater-grass.

Gramen Alopecurinum minus Ger. Fig. 11. Gr. Alopecuroides spica longa minus Park. 1169. Raij Cat. Angl. & H. Pl. 1265: Synops. 182. It Spikes in May and June amongst Corn. Long stender eared Bastard For-tail Grass.

Alopecuros maxima Anglica Park. Fig. 1166. altera maxima Anglica paludosa Ger. 87. Raij Cat. Angl. & H. Pl. 1266. Synops. 183. It Spikes in June and July in Salt Marshes. Great Marsh foresail-grass.

Gramen pumilum hirsutum spica purpuro argentea molli D. Dale. Raij H. Pl. 1265. Fascicul & Synops. 182. On Banfead-Downs and such like places, Spiking in June. Warf for-tail-grass, with a Purplish Bilber coloured Spike.

Gramen Typhinum majus seu 1 Ger. 2. medium seu vulgatissimum Park. Fig. 1170. Raij Cat. Angl. & H. Pl. 1267. Synops. 183. It Spikes in June and July in moist Meadows, near Hedges and Path-ways. The greater Cats-tail-grass.

Gramen Typhinum minus Ger. Fig. 12. Park. Fig. 1170. Raij Cat. Angl. & H. Pl. 1267. Synopf. 183. It Spikes at the same time, and is to be sound in the like places as the greater kind. The Lesser Cats-tail-grass.

Gramen vernum spica brevi laxa Raij H. Pl. 1268. & Synops. 183. It's very common in Meadows and Pastures Flowring in the Spring. Spring-grass, with a loose rellewish spine.

Gramen Cristatum Anglicum Park. Fig. 1160. Crist. Bauhini seu 3 Ger. Raij Cat. Angl. & H. Pl. 1269. Synops. 184: It Spikes about Midsummer plentisully in most Meadows and Pastures. Smooth crested grass.

Gramen tremulum I. B. Phalaris pratenfis Ger. Fig. 86. Raij. Cat. Angl. & H. Pl. 1274. Synops. 185. In Meadows, frequently bearing its Panicles about May. Qualing grass, Cow-quakes, and in some places Ladies hair.

Arundo vallatoria Ger. Fig. 36. Harundo vallatoria seu vulgaris Park. Fig. 1209. Raij Cat. Angl. & H. Pl. 1275. Synops. 185. It Flowreth about September, very frequent in Rivers and Fish-Ponds. Common-Keed.

Gramen dumetorum panicula acerosa semine papposo Raij H. Pl. 1287. & Synops. 185. Cat. Angl. 149. Calamogrostis nostras sylvæs. Johannis Park. Fig. 1181. It Flowers from June till September in Bushy Woods. Reed, or Bushegrass, with a downy panicle.

Gramen Arundinaceum aquaticum Park. Fig. 1274. majus aq. Ger. Fig. 6. Raij Cat. Angl. & H. Pl. 1280. Synopf. 186. It Spikes in June, July, &c. in Rivers frequently. Great Content of Treed-grafs.

Gramen Arundinaceum aquaticum panicula Avenacea D. Doody, Raij in Append. H. Pl. 1909. Fasc. 9. & Synops 186. In the River Thames; between London and Chelsey, and other the like places, Flowring in June and July. That is grass, botth an Dat-like panicle.

Gramen Arundinaceum acerosa gluma nostras Park. Fig. 1273. Rais Cat. Angl. & H. Pl. 1280. Synops. 186. It Flowers from Midsummer till August in Watry Ditches, and Rivers. Parkinson's Great Reed-grass, with Chassy heads.

Gramen aquaticum cum longissima panicula I. B. sluviatile Ger. Fig. 14. Park. Fig. 1275. Raij Cat. Angl. & H. Pl. 1281. Synopf. 186. It Spikes in May, June, &c. in Rivers and running Ditches. Chater-grass, both bery slender and long Panicles.

Gramen paniculatum aquaticum minus Raij Cat. Angl. & H. Pl. 1281. Synops. 186. Miliaceum fluitans suavis saporis D. Merret. in Pin. 53. It Flowers in June, on the Banks of Rivers, and in standing Waters. Dr. Merrer's Water Pillers grass, with the taste of Liquorish.

Gramen segetale Ger. Fig. 5. segetum panicula speciosa Park. Fig. 1158. Raij Cat. Angl. & H. Pl. 1283. Synops. 87. It Spikes in June and July in moist Woods and Pastures, and amongst Corn. Fair panicled Corn-grass, or Bent-grass.

Gramen agrorum venti spica Park, sig. 1158. harundina-ceum Ger. sig. 5. Raij Cat. Angl. & H. Pl. 1284. Synops. 87. It Spikes about Midsummer amongst Corn. Cozu Reed-grass, or Bent.

Gramen Avenaceum pratense elatius panicula slavescente locustis parvis Raij Cat. Angl. & H. Pl. 1284. Synops, 187. It Spikes about Midsummer in Meadows and Pastures. Mr. Ray's meadow Dat-grass, with a yellowish manicle.

Gramen pratense minus seu vulgatissimum Raij H. Pl. 1284. Synops. 87. an prat. minimum album Park. sig. 1156? minimum album Ger. sig. 3? It Spikes from Spring till Autumn, in Pastures, and near Path ways. The Soul common Speadow-grass.

Gramen pratense Ger. sig. 2. prat. vulgatius majus Park. sig. 1156. Raij Cat. Angl. & H. Pl. 1285. Synops. 188. It Flowers in June and July, in Pastures frequently. The greatest Peadow-grass.

Gramen pratense paniculatum molle Park. fig. 1155. Ger. 29. Raij Cat. Angl. & H. Pl. 1285. Synops. 188. It Flowers in June and July in Pasture Meadows. Soft tusted Speadow-grass.

Gramen Caninum paniculatum molle Raij H. Pl. 1285. Fasc. 11. & Synops. 188. It Flowers in June amongst Corn. Mr. Ray's Soft tusted Dogs-grass.

Gramen pratense panicula duriore laxa unam præcipue partem spectante Rais Synops. 188. H. Pl. 1285. & Cat. Angl. 149. It Spikes in June, very common in most Pastures. Mr. Ray's Peadow-grass, with harder sparsed Austs.


Gramen pratense vulgare spica sere Arundinacea I. B. Raij Cat. Angl. & H. Pl. 1286. Synops. 189. It Spikes in June, in most Pastures. Speadow grass, with a Reed-like Spike.

Gramen murorum spica longissima Goodyeri Ger. sig. 29. spica nutante longissima Park. sig. 1162. Raij Cat. Angl. & H. Pl. 1286. Synops. 189. It Spikes in May and June, in dry barren Grounds, and on Walls srequently. Goodyer's Capon-tail-grass.

Gramen paniculatum bromoides minus paniculis aristatis unam partem spectantibus Raij Cat. Angl. & H. Pl. 1287. Synops. 189. It Spikes in June on Walls, and in Sandy grounds. Mr. Ray's Small tusted Dat-grass, with awas.

Gramen minus duriusculum Ger. fig. 4. panicula multiplici Park, fig. 1157. Raij Cat. Angl. & H. Pl. 1287. Synops. 189. It Spikes in June in dry places, and on Walls frequently. Small Baid-grass.

19/ 1 7 100 - 100 100 . Programs and and the state of t or we have a more raides i acada a 1879 raide raide i company acada (the fitzgall. D. Dog - 1 1 新教 (1) Carr C h e effitis - Builty 18 The Lines Az Pija. 1 14) Ray Com. - he per large land . 1 . 151222 Gerrich til rsy. y c. - in the state of the Control of the contro Tep Co. 7 1 1 1 E 3 Cow grakes in a maller dineate Volgar : Parl Turing Spreech St. 16. Craming Same H. Pl. 129 . & E . no ites its second A. A. Tree age bearing



Conyza minima Ger. Fig. 482. Raij Cat. Angl. & H. Pl. 262. Synops. 48. It Flowers in July and August, in muddy places, where the Water hath stood all the Winter. Dwarf or Small Fleabane.

Convolvulus major albus Park. Fig. 163. Smilax levis seu lenis major Ger. Fig. 861. Raij Cat. Angl. & H. Pl. 725. Synops. 102. It Flowers in July and August, in Hedges, Orchards, and moist places. Great Bindweed.

Convolvulus minor vulgaris Park. Fig. 171. Smilax lenis minor Ger. Fig. 861. Ray Cat. Angl. & H. Pl. 725. Synops. 162. It Flowers from Midjummer till Autumn, in Gardens, plowed Lands, and by Path-ways. Small Budweed.

Convolvulus minor Atriplicis folio Park. Fig. 172. Volubilis nigra Ger. Fig. 863. Raij Cat. Angl. & H. Pl. 182. Synopf. 32. It Flowers and Seeds from July till Antumn: It's a Weed in Gardens, Vineyards, and amongst Corn. Black Bindweed.

Cornus sæmina Ger. Fig. 1467. Park 1521. Raij Cat. Angl. & H. Pl. 1591. Synops. 225. In Hedges and Woods, slow-ring in May and June; the Berries are ripe in Autumn. The Dogsberry, Gatter-Erw, or semale Cornel.

Coronopus Ruellii Ger. Fig. 427. recta vel repens Park. Fig. 502. Raij Cat. Angl. & H. Pl. 843. Synopf. 112. It Flowers and Seeds from May till Angust in moist places, and on Ditch-banks. Swines Cresses.

Cotula fætida Ger. Fig. 757. Park. 86. Raij Cat. Angl. & H. Pl. 355. Synops. 57. It begins to Flower about Midsummer, amongst Corn and in plowed Fields. Paithes, or Stinking Parsund.

Elaphoboscum Matth. Pastinaca latif. sylvest. Ger. Fig. 1035. Park. 944. Raij Cat. Angl. & H. Pl. 409. Synops. 63. It Flowers and Seeds in July and August, on the banks and borders of Fields, especially in Salt Marshes. This Parsneps.

Erica vulgaris hirsuta Ger. Fig. 1380. vulg. hirsutior Park. Fig. 1480. Raij Cat. Angl. & H. Fl. 1713. Synops. 230. It's often found with the common, Flowring from Midsummer till August. Common rough leabed Beath.

Erica tenuisolia Ger. Fig. 1382. virgata seu 6 Clus. Park. Fig. 1483. Rais Cat. Angl. & H. Pl. 1718. Syn. 231. It Flowers most part of the Summer, in wast and Heathy Grounds. Fine leabed Beath.

Eruca aquatica Ger. Fig. 248. Park. Fig. 1242. Raij Cat. Angl. & H. Pl. 808. Synops. 112. It Flowers in June and July, in Ditches, and Water courses. Conster Mocket.

Eruca sylvestris Ger. Fig. 246. sylv. vulgatior Park. Fig. 818. Raij Cat. Angl. & H. Pl. 807. Synops. 112. It Flowers and Seeds from Midsummer till Autumn, on Brick, and old Stone-Walls frequently. Edilo Rochet.

Erysimum latisolium Neapolitanum Park, Fig. 834. Raij Cat. Angl. & H. Fl. 811. Synops. 113. It Flowers about Midsummer, on Walls, and near Path ways. Smeoth bread-leabed Dedue-Dustaro.

Eupatorium Cannabinum Park, Fig. 495. Cannab. mas. Ger. Fig. 711. Raij Cat. Angl. & H. Pl. 293. Synopf. 51. It Flowers in July and August, on the brinks of Rivers, and in running Waters. Dutch or Common Demp Agrimony.

Eupatorium Cannabinum sæmina Ger. Fig. 711. aquat. duorum generum Park. Fig. 595. Raij Cat. Angl. & H. Pl. 360. Synops. 57. It Flowers in July and August, in standing Pools and most Watry places. Canater Henips Agrimony, with a divided Leaf.

Eupatorium aquaticum fol, integro Park, Fig. 595. Cannab. 2m. varietas altera Ger. 711. Raij Cat. Angl. & H. Pl. 361. 3nopl. 57. It Flowers in July and August in moist Watry laces, and standing Pools. Chater Bemps Agrimony with n undivided Leaf.

Galeopsis vera Ger. Fig. 704. legitima Dioscoridis Park. Fig. 608. Raij Cat. Angl. & H. Pl. 548. Synops. 80. It Flowers about Midsummer, in Hedges and Bushy places. Deoge- Pettle.

Gallium album Ger. Fig. 1126. Mollugo vulgatior minor fl. albo Park, 565. Raij Cat. Angl. & H. Pl. 481. Synopf. 73. It Flowers in June in moist Warry places. White Laoies Ecotratu.

Genista spinosa vulg. Ger. Fig. 1319. Park. Fig. 1004. Raij Cat. Angl. & H. Pl. 1729. Syn. 232. It Flowers from May till Autumn, in Heathy and Barren places abundantly. Common furze, whins or Gosse.

Genista spinosa minor Fark. Fig. 1004. aculeata minor sive Nepa Theophrasti Ger. Fig. 1321. Raij Cat. Angl. & H. Pl. 1729. Synops. 232. It Flowreth a little before Autumn, and is to be found with the common. The Lesser Fitze Bush.

Genista aculeata Ger. Fig. 1320. Genistella aculeata Park. Fig. 1004. Raij Cat. Angl. & H. Pl. 1731. Synops. 233. It Flowers about Midsummer, in Barren-Grounds, as Hamstead-Heath, &c. Peedle-Furze, or Petty-Whin.

Genistella tinctoria Ger. Fig. 1316. Genista tinct. vulgaris Park. Fig. 229. Raij Cat. Angl. & H. Pl. 1725. Synops. 232. It Flowers in July and August in most Barren Grounds. Green-weed, Dyers-weed, or Moodwaren.

Geranium Columbinum majus dissectis foliis Ger. 938. Malacoides laciniatum five Columbinum alterum Park 706. Raij Cat. Angl. & H. Pl. 1059. Syn. 155. It Flowers in the Spring in Barren Pastures, and near Hedges. Dobes foot Cranes-bill, with jagged Leaves.

Geranium moschatum inodorum Park. 708. Cicutæ folio inodorum Ger. Fig. 945. Raij Cat. Angl. & H. Pl. 1057. Synops. 154. It Flowers about Midsummer on Sandy Banks, and in Gravelly Grounds. Mnsaboty Field Cranes-bill

Gladiolus palustris Cordi Ger. Fig. 29. Juncus storidus Park. Fig. 1197. Raij Cat. Angl. & H. Pl. 701. Syn. 107. It Flowers in June and July, &c. in standing Muddy Waters. The Flowring-Mush, or Mater-Gladiole.

Gramen Leucanthemum Ger. Fig. 47. majus Park. Fig. 1325. Raij Cat. Angl. 162. H.Pl. 1027. Synopf. 146. Frequent in Woods and Buthy places; flowring in the Spring. Sptitche wort, with large Flowers.

Gramen Leucanthemum alterum Ger. 46. minus Park. 1324. Raij Cat. Angl. 163. H. Pl. 1027. Synops. 146. It Flowers in May and June in Sandy and Bushy places. The Lesses Ditchwort.

Hypericum supinum glabrum Ger. Fig. 541. minus supinum Park. 572. Raij Cat. Angl. & H. Pl. 1019. Synops. 143. It Flowers in June and July, in sandy barren Grounds. The least Leasing St. John's-wort.

Hypericum 5. seu pulchrum Ger. 540. minus erectum Park. 572. Raij Cat. Angl. & H. Pl. 1019. Synops. 143. In low Woods and Heathy places; flowring about August. Smell Expright St. John's-wort.

Jacea nigra Ger. Fig. 727. nigra vulg. Park. Fig. 468. Raij. Cat. Angl. & H. Pl. 325. Synopf. 155. It's very common in Meadows, beginning to Flower about Midsummer. Pattellon or Bnapweed.

Jacobea Senecionis folio incano perenne Raij Cat. Angl. Edit. 2. & H. Pl. 285. Synops. 50. an vulg. minor Park. 608? It Flowers in July, &c. on dry Banks, and in Bushy places. Poary perennial Kagwozt, with Groundsel leabes.

Jacobea latifolia palustris seu aquatica Raij. Cat. Angl. & H. Pl. 385. Synops 50. an latifolia Ger. Fig. 280? It Flowers in June and July in Trenches, and Watry Meadows. Parth Ragmost.

Jacea major Ger. Fig. 727. & Scabiosa st. purpureo ejusd. Fig. 721. nigra vulg. laciniata Park. 470. Raij Cat. Angl. & H.Pl. 325. Synops. 55. It Flowers in July and August in Corn-Fields. Steat Anapweed, or Spatsellen.

Lactuca sylv. major odore Opii Ger. Fig. 309. sylv. Endiviæ soliis odore viroso Park. Fig. 812. Raij Cat. Angl. & H. Pl. 219. Synops. 41. It Flowers in July, in Hedges, Lane-sides, and Stony places. Bread leaded strong scented will Lettice.

Lagopus vulg. Park. Fig. 1107. Lagopodium, Pes Leporis Ger. Fig. 1193. Raij Cat. Angl. & H. Pl. 948. Synops. 135. It Flowers in June, July, &c. amongst Corn, and in Sandy barren Grounds. Pares=stoot, or Pares=stoot Trefoil.

Lamium luteum Ger. Fig. 702. Park. Fig. 606. Raij Cat. Angl. & H. Pl. 560. Synopf. 82. It Flowers in May and June, in Woods and shadowy places. Pellow Archangel.

Lamium perfoliatum vulgare Nobis. Alsine Hederula altera Ger. Fig. 616. Hederulæ solio major Purk. Fig. 762. Raij Cat. Angl. & H. Pl. 560. Synops. 82. It Flowers about Mid-summer in Kitchin-Gardens and Tilled Grounds. Great Penbit.

Lapathum acutum minus Park. 1226. acuti varietas folio crispo Ger. 387. Raij Cat. Angl. & H. Pl. 173. Synops. 30. It Flowers and Seeds from June till September, in Rivulets and Watry Ditches plentifully. Sharp pointed Dock, with curbed Leaves.

Lapathum pulchrum Bononiense sinuatum 7. B. Tom. 2. Lib. 23. pag. 688. Raij Cat. Angl. & H. Pl. 174. Synops. 31. It Flowers and Seeds from June till Autumn in several barren Grounds about London and elsewhere. Fiddle Dock.

Luteola Ger. Fig. 594. vulg. Park. Fig. 603. Raij Car. Angl. k. H. Pl. 1054. Synops. 153. It Flowers in June and July, on Walls, waste Grounds, and in Fallow-Fields. Sailo school, ellew stated, and Diers-weed.

Lychnis sylvestris sl. albo Ger. 468. Park. Fig. 631. Raij Sat. Angl. & H. Pl. 994. Synops. 140. It Flowers from May ill August, in the borders of plowed Fields, Meadows, and Ditch banks.

Lychnis sylv. rubello store Ger. Fig. 469. sylv. st. rubro sylv. Fig. 631. Raij Cat. Angl. & H. Pl. 994. Synops. 140. If Flowers the most part of the Summer in Hedges and moist pitches frequently. Red stowed with Campion.

Lysimachia galericulata Ger. Fig. 477. Gratiola cærulea w latisolia major Park. Fig. 631. Raij Cat. Angl. & H. Pl. 1972. Synops. 84. It Flowers in July and August in running Ditches, and by Waters sides. The Greater booded waitlow Herb.

Lysimachia lutea Ger. Fig. 474. lutea maj. vulg. Park. Fig. 44. Raij Cat. Angl. & H. Pl. 1021. Synops. 144. It Flowers om June till August in moist Meadows, and by Waters sides. Ellow Millow Beth, or Loosestic.

Lysimachia purpurea spicata Ger. Fig. 476. Park. Fig. 546. bij Cat. Angl. & H. Pl. 1036. Synops. 149. It Flowers in me and July in moist Ditches, and near running Waters. urple spiked Willow Berb, or Loosestrife.

Lysimachia purpurea trifolia caule hexagono Spigel. Isag. if Cat. Angl. & Synops. 149. It grows in the same places, a Flowers at the same time with the Common Purple spiked low Herb. Three leaved Purple spiked Loosearise.

Lysimachia campestris Ger. Fig. 478. siliquosa major Park. § 548. Raij Cat. Angl. & H. Pl. 861. Synops. 123. In moist evelly Gardens, Walks, and Woods, slowring a little after summer. The Greater smooth leabed, codded existion ath, or Loosestrise.

Lysimachia siliquosa glabra media sive minor Ger. 479. if Cat. Angl. & H. Pl. 867. Synops. 123. It's found usually Rivers and Ditch sides; slowing in July and Angust. pole smooth leabed codded willow Derb, or Lossestrise.

Lysimachia siliquosa glabra angustisolia Ger. 479. Raij Car. Angl. & H. Pl. 862. Synops. 124. It Flowers in July and August in Marsh and Watry Meadows. The Least smooth teabed Cooked willow Perb, or Loosestrise.

Lysimachia siliquosa hirsuta magno slore C. B. pin. 245. siliquosa Ger. Fig. 476. Raij Cat. Angl. & H. Pl. 861. Synops. 123. Near running Dirches and River sides; Flowring in July and Angust. Great hairy codded Locsestrife, by some Coolings and Cream.

Lysimachia siliquosa hirsuta parvo slore C. B. pin. 245. sylvatica seu 12. Ger. 479. an siliquosa sylv. hirsuta Park. 549? It Flowers about the same time, and is to be sound in the same places with that called Codlings and Cream. The Lesser bairy codded Loosestrie, or willows yerb, with small Flowers.

Malva sylvestris pumila Ger. Fig. 930. sylv. minor Park. 299. Rasj Cat. Angl. & H. Pl. 599. Synops. 88. It Flowers from June till Autumn, in Kitchin-Gardens, Untilled places, and near Pathways. Small wild Spallow, or Dwarf Spallow.

Marrubium aquaticum Ger. Fig. 700. aq. vulg. Park. Fig. 1230. Raif Cat. Angl. & H. Pl. 535. Synopf. 79. It Flowers in July and August in Rivers and Watry places. Water Duzehound.

Melampyrum luteum latifolium C. B. pin. 234. Crateogonon vulg. Park. Fig. 1326. album Ger. Fig. 91. Raij Cat. Angl.& H. Pl. 775. Synops. 108. In low barren heathy Woods, as Hampstead-Heath, and such like places; flowring from May till August. Common rellow Cow-Waheat.

Mentha aquatica rubra Park. Fig. 1243. aq. five Sifymbrium Ger. Fig. 684. Raij Cat. Angl. & H. Pl. 1101. Synopf. 162: It Flowers in July and August in watry Ditches, and on the fides of Brooks frequently.

Millesolium aquaticum floridum sive Viola aquatica Park. Fig. 1256. Viola palustris Ger. Fig. 826. Raij Cat. Angl. & H. Pl. 1101. Synops. 162. In muddy and flow running Waters; flowring in May and June.

Millefolium fl. rubro Ger. Fig. 1072. rubrum vulg. Park. 693. Raij Cat. Angl. & H. Pl. 346. Syn. 56. It Flowers from June till Antumn in the same places with the common. Putple slowered Parrow, or spilifoil.

Mollugo montana minor Gallio albo fimilis Raij Cat. Angl. & H. Pl. 482. Synops. 74. It Flowers about Midsummer, on Mountainous and Heathy Grounds, and also in Marshy places. Small Sountain Bastard Padder.


Myosotis scorpioides arvensis hirsuta Ger. Fig. 337. scorp. hirsuta Park. Fig. 691. Raij Cat. Angl. & H. Pl. 504. Synops. 77. It Flowers in June and July in Gardens and dry barren Grounds. Souse-ear Scorpion-grass.

Myosotis scorpioides minor flosculis Iuteis Park. 692. Raij Cat. Angl. & H. Pl. 504. Synops. 77. It Flowers in April and May in sandy barren Pastures. Simal yellow flowered Scozpionsgrass.

Myosotis scorpioides palustris Ger. Fig. 337. scorp. repens Park. Fig. 691. Raij Car. Angl. & H. Pl. 504. Synops. 77. It Flowers in June and July in Rivulets and Watry Ditches. Water Scoppion-grass.

Myrrhis sylvestris Neapolitana & etiam Anglicana Park. 935. Ceresolium 5 sive Myrrhis nova Æquicolorum Columnæ Ger. 1038. Raij Car. Angl. & H. Pl. 469. Synops. 71. It Flowers and Seeds in May and June, on Ditch-banks and near High-ways plentifully. Small Bemlock Cherbil, with rough Seeds.

Oenanthe palustris sive aquatica Park. 895. & aq. major Ejusal. Fig. 1233. Filipendula aq. Ger. Fig. 1060. Raij Cat. Angl. & H. Pl. 441. Synops. 65. It Flowers in May and June in Marshy Meadows, and near Rivers. Eyeat water Dropwort.





Gramen afperum I.B. pratense spica multiplici rubra Park. sig. 1160. Raif Cat. Angl. & H. Pl. 1287. Synops. 190. It Spikes in May & June, in Meadows & Pastures frequently. Mough-grass.

Gramen Cyperoides spica simplici compressa distica D. Plukenet, Raij Fascie. 10. & Synops. 199. It Spikes in June and July in Boggy and Moorish places. Dr. Plukenet's Soggy Cyperus-grass.

Bromos sterilis Ger. fig. 76. herba sive Avena sterilis Park. fig. 1148. Raij Cat. Angl. 105. & H. Pl. 1289. Synops. 191. It Spikes in May and June near Path-ways, and in the Borders of Fields. Great wild Datograss, or Drank.

Festuca Avenacea hirsuta paniculis minus sparsis Raij Cat. Angl. & H. Pl. 1289. Synops. 191. It Spikes in May and June, on Walls, and in the borders of Fields very common. Rough Dat-grass, with closer Tusts.

Festuca altern Dod. Raij Cat. Angl. Avenacea spicis habitioribus glumis glabris Raij H. Pl. 1289. & Synops. 192. It Spikes in June and July amongst Corn. Dat-grass, with looser and smoother Tusts.

Gramen exile hirsutum Ger. fig. 17. nemorum hirsutum minus angustisolium Park, fig. 1185. Raij Cat. Angl. & H. Pl. 1291. Synops. 193. It Flowers in April and May, in Meadows and Pastures frequently. Small Bairy-grass.

Gramen nemorosum hirsutum vulgare Raij Cat. Angl & H.Pl. 1292. Syneps. 193. an hirsutum nemorosum Ger. sig. 19? an nemorosum hirsutum majus Park. sig. 1184? In hilly Woods frequently. Flowring in the Spring. Common hair 2000 grass.

Gramen Cyperoides foliis Caryophylleis vulgatiffimum Raij H. Pl. 1293. & Synops. 194. Graminis nigro-lutei verni varietas minor I. B. It Flowers in May and June in moist Meadows, and near Rivers. Pany spiked Cyperus grass, with July-Flower leaves.

Gramen Cyperoides spicatum 1 Ger. sig. 22. Cyp. vernum minimum Raij Cat. Angl. & H. Pl. 1294. Synops. 196. It Spikes early in the Spring in Earren Pastures frequently. The Least Mernal Cyperus-grass.

Gramen Cyperoides polystachion lanuginosum Raij Car. Angl. & H. Pl. 1294. Synops. 194. an Gr. Cyp. Norwegicum parum lanosum Park. 1172? It Spikes in May and June, in moist Meadows. Daary spikes Cyperus-grass.

Gramen Cyperoides spica pendula breviore C. B. pin. 6. Pseudo-Cyperus Ger. sig. 29. Cyperus sive Pseudo-Cyperus spica brevi pendula Park. sig. 1266. Raij Cat. Angl. & H. Pl. 1294. Synops. 195. It Spikes about Midsummer in moist watry Ditches. Batard Cyperus, with short pendulous heads.

Gramen Cyperoides spica pendula longiore Park. 1267. Raij Cat. Angl. & H. Pl. 1294. Synops. 195. It Spikes in June and July in moist Dirches. Battard Typerus grass, with long pendulous heads.

Gramen Cyperoides sylvarum tenuius spicatum Park, sig. 1171. Raij Cat. Angl. & H. Fl. 1295. Synops. 196. It Spikes in April and May in Woods frequently. Lobel's Slender ear'd Wasod Experus-grass.

Gramen Cyperoides palustre majus Park, sig. 1266. palust. Cyp. Ger. sig. 19. Raij Cat. Angl. & H. Pl. 1297. Synops. 198. It Heads in May, June, &c. near Rivers, and in Watry Ditches. The Greater spines Cyperus-grass.

Gramen Cyperoides spicatum minus Raij H. Pl. 1297. Synops. 198. Cyp. spicis minus compactis Park. 1268. Cyp. parvum Ger. 20. (agrecing with his Description.) It Spikes in May and June in most Meadows and Pastures. The Lester spiked Cyperus-grass.

Gramen Cyperoides spica è pluribus spicis brevibus mollibus composita Rais H. Pl. 1296. & Synops. 197. It Spikes in May and June in moist Pastures and Woods. Mr. Ray's Experise stees, with many roundish heads.

Gramen caninum nodosum Ger. sig. 23: can. nodos. bulbosum vulgare Park. 1175. Raij Cat. Angl. & H. Pl. 1284. Synops. 191. It Spikes in June, in Woods, Hedges, and amongst Corn. Enterty Degs grass.

Gramen tomentarium Ger. fig. 29. Juncoides lanatum sive Juneus bombyeinus vulgaris Park. fig. 1272. Raij Cat. Angl. & H. Pl. 1306. Synops. 203. It Flowers in June and July in Boggy Moorish Grounds. Cotten-grass.

Gramen junceum aquaticum Bauhini Park, fig. 1270. aquat. Ger. fig. 13. Raij Cat. Angl. & H. Pl. 1307. Synops. 204. It Flourishes from Midsummer till Autumn in wet and marshy places. Rull-grass, with differing heads.

Gramen junceum aquaticum magis sparsa panicula Park. fig. 1269. an Gr. junceum sylvaticum Ger. fig. 22? Raij Cat. Angl. & H. Fl. 1307 Synops. 204. It Flourishes in June and July in moist Woods, and watry Places. The Greater must grass with large Tusts and joynted Leaves.

Gramen junceum capitulis triangulis minimum Raij Cat. Angl. & H. Il. 1307. Synopf. 204. Graminis juncei varietas minor Ger. 4. It Flourifles about Midsummer in moist boggy places. The Least mush-grass, with triangular heads.

Gramen junceum capfulis triangulis cauliculis tenuibus foliis ad nodos & panicularum divaricationes prælongis Raij H. Pl. 1307. & Synopf. 204. It Flourishes in July in Marshy and Boggy places. Mr. Dale's Least triangular headed Rushegrass, with a foliaceous head.

Gramen junceum maritimum Ger. fig. 21: marit. majus Park. fig. 1270. Raij Cat. Angl. & H. Pl. 1307. Synopf. 204. It Flourishes about Midsummer in moist Meadows, and in Salt Marshes. Rush-yrass, with round Seed-Vessels.

Gramen junceum Ger. fig. 4. junceum parvum five Holoftium Matthioli Park. fig. 1190. Raij Cat. Angl. & H. Pl. 1309. Synops. 204. It Flourishes in May, June, &c. in wet and muddy places. Toat-grafs.

Juncus acutus Ger. fig: 35. acutus vulg. Park. fig. 1193. Raij Cat. Angl. & H. Pl. 1303. Synopf. 201. It Heads in June, July, &c. in moist neglected Pastures. Common Bard-Rush.

Juncus montanus palustris Raij H. Pl. 1303. Synops. 201. an Oxyschenos five Juncus acutus Alpinus Cambrobritannicus Park. sig. 1193? In Hilly, Wet, and Moorish Grounds. Flourishing from Midsummer till Spring. Solstus, or One-Com.

Juncus aquaticus maximus Ger. fig. 35. lævis maximus Park. fig. 1191. Raij Cat. Angl. & H. Pl. 1304. Synopf. 201. It Flourishes in July, August, &c. in most Rivers plentifully. Common Bull-ruth.

Juncus caule triangulati D. Merret. in Pin. 67. an acutus maritimus caule triangulo C. B. prodr. 22. fig.? Raij Cat. Angl. & H. Pl. 1303. Synopl. 201. It's found in the Thames, both above and below London, Flourishing in July, August, &c. Will-1989, with a triangular Stalk.

Juncus lævis vulgaris panicula sparsa nostras Raij H. Pl. 1304. & Synops. 202. an lævis Ger. sig. 35? an lævis panicula sparsa major Park. sig. 1191? In Ponds, and by River-sides, Florishing with the common kind. Tusted soft Rush.

Juncus Levis vulgaris panicula compactiore Raij H. Pl. 1304. & Synops. 202, an levis vulg. Park.? & Levis glomerato flore Ejusa. fig. 1191? It Florishes in July and August, in moist Pastures and Woods. Close headed soft Rush.

Juncus minor capitulis Equiseti Ger. fig. 1631. aquaticus capit. Equiseti Pack, fig. 1196. Ra. jCat. Angl. & H. Pl. 1305. Symps. 202. It Florishes in May and June, in Ponds and Rivers frequently. Club, or Eglet headed Rush.

Juncellus Lobelii Park. 1192. Raij Cat. Angl. & H. Pl. 130A. & Synops. 203. Gramen sparteum capite bisido vel gernino D. Merret. in Pin. 58. It Florishes in June, July, &c. in Moist. Sandy, and Boggy places. The Least Rush.

Alnus Ger, fig. 1477. vulgaris Park. fig. 1409. Raij Car. Angl. & H. Pl. 1409. Synopf. 213. Near Rivers, and in moist Warry places, bearing its Cones from May till September. The Common Coniferous Alder.

Erica major fl. purpureo Ger. Fig. 1381: pumila Belgarum Lobelio, Scoparia noltras Park. fig. 1482. Raij Cat. Angl. & H. Pl. 1714. Synops. 231. In Boggy Heaths, Flowring in June, July, &c. Deary Beath, with pale Purple Flowers.

Euonymus vulgaris Park, fig. 241. Theophrasti Ger. fig. 1468. Raij Cat. Angl. & H. Pl. 1621. Synops. 228. It's sound in Hedges, Flowring in May, and the Berries are ripe in September. Prick-wood, or Spindle-Tree.

Fagus Ger. fig. 1444. Park. 1403. Raij Cat. Angl. & H. Pl. 1381. Synops. 212. It Flowereth in April and May, and the Fruit is ripe in September, in Woods and Hedges. The Beach-Aree.

Gnaphalium minimum I. B. Filago minor Ger. 641. (whoso Figure is ill exprest; the Icon of the Gnaphalium majns Germanicum Park. 685. is much more like it.) Raij Cat. Angl. & H. Pl. 206. Synopf. 52. It Flowers in May, June, &c. in Sandy Barren Grounds, and sometimes amongst Corn. The Least Cudweed.

Lampsana Ger. fig. 255. vulgaris Park, fig. 810. Raij Cat. Angl. & H. Pl. 256. Synops. 47: It Flowers in June and July, in Tilled Fields and Gardens frequently. Ripple-wort.

Lapathum sylvestre vulgatius Park. sig. 1225. sylv. solio minus acuto Ger. sig. 388. Raij Cat. Angl. & H. Pl. 173. Synops. 30. It Flowers and Seeds from June till September, in moist Fields, and near Path-ways. The Most common broad leabed Wilds Dock.

Lapathum acutum minimum Ger. 388. Park. Fig. 1224. Raij. Cat. Angl. & H. Pl. 174. Synops. 31. It Flowers and Seeds in July, August, &c. near Ditches, and in moist places. Small sharp pointed Dock.

Napus sylvestris C. B. pin. 95. Bunias sive Napus sylvestris Park. sig. 865. sylv. Lobelii Ger. sig. 235. Raij Cat. Angl. & H. Pl. 802. Synops. 110. It Flowers in May, on Ditch-banks, and amongst Corn frequently.

Rapunculus Scabiosæ capitulo cæruleo Park. 646. Scabiosa minima hirsuta Ger. sig. 723. Raij Cat. Angl. & H. Pl. 744. Synops. 104. It Flowers in June and July, on Walls, and in Barren chalky Grounds. Base-Botket.

Salix pumila angustisolia recta Park. sig. 1434. Chamæitea sive Salix pumila Ger. sig. 1391. Raij H. Pl. 1422. & Synops. 215. In Heaths and Boggy places. Common creeping Dwarf-willow.

Salix folio longo latoque splendente fragilis Cat. Cant. Raij Cat. Angl. & H. Pl. 1420. Synops. 215: an Salix fragilis Park. 1431? In Osier-holts and Watry places, Flowring in the Spring. The Crack-Wiltow.

Sedum minus hamatodes Ger. fig. 512. Vermicularis hamatodes Park. in margine 733. Raij Cat. Angl. & H. Pl. 691: Synopf. 99. On old Houses and Walls, Flowring in June and July. Didinary yellow Prickmadam, or Stone-crop.

Sedum minus Officinarum Ger. fig. 512. Vermicularis fl. albo Park. 733. Raij Cat. Angl. & H. Pl. 1040. Synopf. 150. It Flowers about Midsummer on old Houses and Walls. Anglie flowered Stone-crop, with round pointed Leaves.

Selinum Sii foliis Ger. fig. 1018. segetale Park. fig. 932.
Raij Cat. Angl. & H. Pl. 443. Synops. 66. It Flowers in July and August in moist Claie Corn-Fields. Goodyer's Donewort.

Sium majus angustisolium Ger. sig. 256. Raij Cat. Angl. & H. Pl. 444. Synops. 66. It Flowers and Seeds in July and Angust in Rivulets. Common upright Waters Parsnep.

Sium 3 seu umbellatum repens Ger. 256. Raij Cat. Angl. & H. Pl. 444. Synops. 66. It Flowers in June and July in small Rivers. Creeping Mater Darsnep.

Sium minimum Raij Cat. Angl. & H. Pl. 444. Synopf. 67. It Flowers and Seeds in May, June, &c. in Ponds and Watry Ditches. The Least mater partnep.

Sphondylium Ger. fig. 1009. vulgare Park. fig. 953. Raij Cat. Angl. & H. Pl. 408. Synopf. 63. It Flowers in June and July in moist Meadows and near Rivers. Cowsparsnep.

Trisolium pratense album Park. sig. 1110. minus pratense sl. albo seu 2 Ger. 1185. Raij Cat. Angl. & H. Pl. 942. Synops. 133. It Flowers about Midsummer, in Meadows frequently. Thite slowered Speadow Tresoil.

Trisolium pratense Ger. sig. 1185. prat. vulgare purpurcum Park. sig. 1110. Raij Cat. Angl. & H. Pl. 943. Synops. 134. In Meadows frequently, Flowring most part of the Summer. Common Purple, or Honey-Suchle Tresoil.

Trisolium purpureum majus sive sativum Raij Cat. Angl. pratense purpureum majus Ejusa. H. Pl. 944. & Synops. 134. It Flowers from June till September in Meadows and near Hedges. The Greater Purple Deadow Tresoil.

Valeriana minor Ger. fig. 1075. fylvestris minor Park. fig. 123. Raij Cat. Angl. & H. Pl. 388. Synops. 61. It Flowers in May and June near Rivers, and in Marshy Grounds. Small mild Malerian.

Valeriana sylvestris seu palustris minor altera Raij Cat. Angl. & Synops. 61. aq. minor sl. minore Ejusd. in H. Pl. 389. It grows in Watry places, and Flowers with the other. Ray's water-malerian, with small Flowers.

Verbascum fl. albo parvo I. B. mas foliis longioribus Park. 60. Verbascum Lychnite Matthioli Ger. fig. 775. Raij Cat. Angl. & H. Pl. 1095. Synops. 160. It Flowers in June and July in Barren Fields, and by Way sides. Aprillein, with small white Flowers.

Viburnum Fark. fig. 1448. Lantana five Viburnum Ger. fig. 1490. Raij Cat. Angl. & H. Pl. 1590. Synops. 225. In Hedges, Flowring in May, and bearing ripe Berries about September. The petiant wealy, or way faring-tree.

Vicia Ger. fig. 1227. vulg. fativa Park. fig. 1072. Raij Cat. Angl. & H. Pl. 900. Synopf. 129. It's frequently Sown in Fields, and Flowers in May. Common Tare, or Hetch.

Vicia fylv. five Cracca major Ger. fig. 1227. Aracus seu Cracca major Park. fig. 1071. Raij Cat. Angl. & H. Pl. 902. Synops. 129. In Hedges, on Ditch-banks, and in Sandy places. Flowring from April till Midsummer: Strangle-tare, or wild-Metch.

Vicia maxima dumetorum Ger, fig. 1227. Park. 1072. Raij Cat. Angl. & H. Pl. 901. Synops. 129. It Flowers in May in Woods and Hedges frequently. Buth-Wetch.

Vicia multiflora five spicata Park. 1072. Raij Cat. Angl. & H. Pl. 903. Synops. 130. It Flowers about Midsummer, in Meadows, Woods, and Hedges. Tusted Hetches, or Birds tares.

Vicia segetum cum siliquis plurimis hirsutis C. B. pin. 345. sylv. sive cracca minima Ger. sig. 1228. Arachus sive cracca minor Park. 1070. Raij Cat. Angl. & H. Pl. 902. Syn. 129. It Flowers about May, is a Weed among Corn. Small-tare, with hairy Pods.

Vicia minor segetum cum filiquis paucis glabris Morif. Hist. Oxon. fig. 64. Viciæ sive craccæ minimæ species cum filiquis glabris I. B. Raij Cat. Angl. & H. Pl. 902. Synops. 129. It Flowers and Pods in May and June, amongst Corn, and in moist Woods. Small-tare with smooth Pods.

Urtica minor Ger. fig. 707. Raij Cat. Angl. & H.Pl. 161. ta2 cemifera minor annua Raij Synops. 29. It Flowers and Seeds in July and August, in Kitchin-Gardens and untilled Grounds. The Letter Stinging Pettle.



Fortus Siccus Chirurgicus.

In quo Continetur

Collectio Plantarum quæ maxime à Chirurgis usurpantur in Componendis Balfamis, Oleis, Unguentis, Emplaftris, &c.

N II O

Synonymis Necessariis, Locis Natalibus & Tempore Vigendi.

Accesserunt etiam illarum Partes ac Præparationes quæ maximè sunt in Usu.

Explicatio Nominum Abbreviatorum.

- A. Litera in Margine polita indicat plantas illas quibus | Præp. prefigitur, aut Angliæ Indigenas elle, ant in Agris pallim cultas.
- . B. Caspari Baubini Pinax.
- Ger. Gerardi Historia emaculata d' Tho. Johnsono.
- J. B. Joannis Bauhini Plantarum Historia Universalia.

Raij H. Pl. Indicat Historiam Plantarum Universalem ab

Eruditissimo Viro, & Botanico Peritissimo D. Joanne

Raio nuper Editam.

Explicat Partes istius Plante que in Arte Pharma-

centica quotidie u'urpantur.

Raij Cat. Angl. Denotat Catalogum Plantarum Anglia &

Insularum adjacentium

Ossicinis nostris frequentissime exhibentur.

- Offic. Exbibet Nomina Plantarum, quibm vulgò vocantur in Officins nostru.
- urk. Joannis Parkinsoni Theatrum Botanicum.

A. Lígusfrum Offic. vulgare. Park. Raij Cat. Angl. & H. Pl. 1603. Usu Fol. Floret tota ferè æstate. In Sylvis & Sepibus. Primprint or Printet. Sarifraga pratensis Offic. Seseis pratense nostras Ag. quæ ad flatus Infantium præffat. Hæc & altera species Park. Raij Cat. Angl. & H. Pl. 45 3. Ulu Tota Planta. Præp.

A. Sedum majus seu Sempetvívum Offic. majus vulgare C. B. Raij Cat. Angl. & H. Pl. 687. Víu Fol. In Muris & Techis provenic. Common Poulleek.

Rhabacharum Offic. Icha verum Antiquorum

1702. Usu Fol. Lign. Corr. & Sem. Lingua Avu dichum. Prap. Aq. Sal. In Autumno Semen perficitur. E specie huius Arboris Memna colligitur. The All-Tree.

A. Frantmus Offic. vulgaris Park. Raij Cat. Angl. & H. Pl.

lape promiseud usurpatur. Mendow Sarifranc

Stramonium offic. Solanum fætidum pomo fpinoso oblongo C. B. Raij H. Pl. 745. Usu Fol. & Sem. Præp. Ung. In Hortis colitur. The Chosn-Apple.

A. fracticulum Offic. vulg. Germanicum C. B. Raij Cat. Angl. & H. Pl. 457. Ulu Rad. quæ inter quinque ac etiam inter duas Radices Aperientes locum obtinet, & Sem. quod inter quatuor calida majoraCarminativa annumeratur. Præp. Aq. Syr. & Ol. Semen in Autumno perficit. Community Fennel.

Aftlepias & Aincetaritum Offic. fl. albo C. B. Rajj H. Pl. 1090. Ulu Rad. In Hortis Curiolorum nutritur. Swallow-wort. Alphodelus Luteus & Valla Regia Offic. Inteo & flore & radice C. B. Raij H. Pl. 1192. Ulu Rad. In Hortis Botanicis nonnunquam reperitur. Vellow Alphodel or Rithurs. Surar.

- A. Buglossum Sylvestre Offic. sylv. minus C. B. Raij, Cat. Angl. & H. Pl. 494. Ulu Rad. Interfegetes & in Arvis requietis passim occurrit. Small Casso Bu-
- A. Istitannica Antiquorum Cleta Muningij;in Tractatu de verâ Antiquorum Herbâ Britannicâ. Lapathum maximum aquaticum five Hydrolapathum J. B. Raij Cat. Angl. & H. Pl. 171. Ulu tota Planta sed præcipue Radices. In palustribus ac stagnantibus locis. The Cette Istitannica of the Anticents.
- A. Cpanus Phinol Offic. vulgaris Park. Raij Cat. Angl. & H. Pl. 121. Ulu Flor. inter segetes Maio & Junio Floret. The Coll. Flower or Below-Bottle.
- A. Catblata Offic. & Park. Raij Cat. Angl. & H.Pl. 571. Usi Fol. In incultis & ruderatis locis, Julio Florer. Do-thetwort.
- A. Cructata Offic hirfuta C.B. Raij Car. Angl. & H.Pl. 479. Usu Fol. Ad sepes & in Dumeuis. Bugueed or Cross-wort.

of nithemill Off. five Cucuta minor C. B. Rail H.

- hugus Kibonis Manna colligitur. The Aul-Citt.
- fera C. B. Raij Cat. Angl. & H. Pl. 1604. Ulu Cert. interior potissimum radicis. In Sylvis. The Black Aldet.
- A. flumatia Offic. purpurea Ger. Raij Cat. Angl. & H. Pl. 40ς. Ulu tota Herba. Præp. Aq. Conf. Syr. Simp. & Comp. Pil. & Sal. In fatis & aggeribus terrenis frequens eft. flumitoχp.
- A. Fungus Sambuctnus seu Aurscula Judae Officize Ger. Raij Cat. Angl. & H.Fl. 106. In Sambuci caudice frequenter nascitur. Jews-Eat.
- A. Gerantium Columbinum & Pes Columbinus Offic. vulgare Park. Raij Cat. Angl. & H. Pl. 1059. VII Fol. ad sepes & vias. Doves-foot or Doves foot Cranes-Isill.
- A. Apollus Banac Offic. Nymphea alba minima C. B. Raij Cat. Angl. & H. Pl. 1320. Ulu Fol. Hac raristime præscribuntur inter Medicos, sed aliquando substituuntur pro Umbilico Venera. In Aquis conosis crescit. The Least Collite Catater Lilly or Frog-Bit.
- A. Linum Catharticum Offic. pratenfe floculis exiguis C. B. Raij Cat. Angl. & H. Pl. 1076. Ulu Planta integra. Floret Junio ac Julio. In Pascuis Siccioribus præsertim montos. Poill-Dountain or Purging Dwarfwild-flax.
- dica. J. B. Raij Car. Angl. & H. Pl. 188. Usu Fol. & Rad. In Sylvis & aquosis locis. Orent Alitto Anterian.
- **Epanus major** Offic. maj. vulgaris. Park. Raij H. Pl. 322. Utu Fol. & Flor.qui fæpe à nonnullis usurpantur loco minoris. Seat Islew Bottle.
- A. Sophía Chitutgotum Offic. & Ger.Nafturtium Sylveftre tenuisfimè divisum. C. B. Raij Cat. Angl. & H. Pl. 812. Usu Fol. & Sem. In ruderatis & incultis locis.

- Rhabarbatum Offic. Kha verum hntiquorum Gr. Raij H. Pl. 170. Ulu Rad. Præp. Syr. Troch. Pil. & Extr. Rhubarb.
- Lapathum Patientia Offic. hortense sol. oblongo five secundum Diescondis C.B.Raij HP1. 171. Usu Rad. in Hortis nutritur. Sathen Patience or Ponks-Rhuharh.
- A. Alfola Battía Offic. purpurea fl. simplici odoro C.B. Raij Car. Angl. & H. Pl. 1049. Usi Sem. Fol. quæ sint è numero Herbarum quinque Emollientium, & Flor. qui inter quatuor Cordiales socum obtinent. Marrio & Aprili floret in aggeribus Sepium. The Association.
- Armeniaca Bala Offic. Malus Armeniaca major Park. Raij H. Pl. 1514. Ulu Frust. & Nuclea. Præp. Ol. per express. Chrysomelinum dictum. The Apricock Cree.
- A. Martubium Rigrum Offic. nigrum fætidum, Ballote Dioscoridis C.B. Raij Cat. Angl. & H. Pl. 571. Usu Fol. in ruderatis & ad sepes. Stinking Hoteljound.
- A. Afinca pervients Offic. Clematis Daphnoides minor C. B. Raij Car. Angl. & H. Pl. 1091. Ulu Fol. Ad sepes & in aggeribus fosfarum. Periwinskie.
- Antihora feu Antíthora Offic. Antithora flore luteo Aconiti J. B. Raij H. Pl. 705. Ulu Rad. In Alpibus Helvericis, in Sabandia &c. crescit. The Wilfolesonic Helmet flower.
- A. Enaplatium longifolium humile ramosum capitulis nigris Raij H. Pl. 295. & Cat. Angl. Usu Herba tota. loco Pedis Cati seu Gnapbalij Offic. usurpatur. In Scrobibus siccis & locis depressionibus reperitur. Blackifeabede Cudweed.
- Raij H. Pl. 431. Ulu Fol. & Sem. Pray. Aq. Maio floret. Elent fweet Gathen Chethil or fweet Citely.
- A. Petfoliata Offic. vulgaris Ger. Raij Cat. Angl. & H. Pl. 471. Ulu tota Planta. Inter Segetes. Chotott.

- Epithpmum Offic. sive Cuscuta minor C. B. Raij H. Pl. 1094. Usu Integra Planta. Præp. Syr. Variis Herbis adnascitur. Epithpmum or Dodder of Cime.
- A. Etica Offic. vulg. humilis sempervirens st. purpureo & albo J. B. Raij Cat. Angl. & H. Pl. 1713. Usi Fol. & Flor. In Ericetis. Common Death.
- A. CIIMUS Offic. vulgatissimus fol. lato scabro Ger. Raij Cat. Angl. & H. Pl. 142?. Usu Rad. Fol. Cert. In Sepibus & Sylvis crescit. The Elm.
- A. Almaría & Begina Pratí Offic. Barba Capri floribus Compactis C. B. Raij Cat. Angl. & H. Pl. 623. Usu Fol. Præp. Aq. Junio & Julio floret, in Pratis Humidis frequenter occurrit. Deadous-Sueet.
- A. Clttica Offic. urens maxima C. B. Raij Cat. Angl. & H. Pl. 160. Usu Herba tota. Prep. Aq. Æstate floret, ad sepes & in locis ruderatis. Common Nettse.
- Cinata & Scolynnus Offic. Sativus Spinosus & non Spinosus J. B. Raij H. Pl. 299. Usu Fel. Caules & Capitum Squamma, sed frequentius in Culium quam in Officinus unsurgatur. Attichouse.
- Rola Daniascena Ger. & pallida Offic. Raij H. Pl. 1468. Usu Flor. Præp. Ag. Spr. Conf. Mel. Spr. &c. The Dannask Rose.
- Rosa Rubra Offic. & Ger. Raij H. Pl. 1468. Usu Flor. (qui inter quatuor Cordiales Flores recensentur) & Anthera. Præp. Aq. Cóns. Syr. Mel. Ol. Ung. &c. The Red Rose.
- Sumach & Rhots Obsoiorum Offic. Rhus folio Ulmi C. B. Raij H. Pl. 1590. Usu Fol.& Sem. In Hispania & Gallia crescit. Common Sumnath.
- Cerebinthus Augaris Offic. & C.B. Raij H. Pl. 1577. N.B. Terebinthinam veram, Officinis esse ignotam; Officinalemque esse vel resinam Lariceam, vel Abiernam. The Curpentine-Aree.





Hortus Siccus Pharmaceuticus:

SIVE,

Collectio Euriofissina Plantarum

Officinalium;

Tam Exoticarum quam Indigenarum, Alphabetico Ordine diligenter digelta.

11 11

Synonymis Necessariis, Locis Natalibus & Tempore Vigendi.

Accesserunt etiam illarum Partes ac Preparationes quæ maximè sunt in Usu.

Explicatio Nominum Abbreviatorum.

- A. Litera in Margine posita indicat plantas illas quibus prafigitur, aut Anglia Indigenas este, aut in Agris passim autin.
- C. B. Castari Baubini Pinax.
- Ger: Gerardi Historia emaculata à Tho. Johnsono.
- J. B. Foannis Banbini Plantarum Historia Universalus
- Office Exibet Nomina Plantatum, quibus vulgò vocantur
- Raij Cat. Angl. Denotat Catalogum Plantarum Anglia & Infularum adjacentium.

 Raij H. Pl. Indicat Historiam Plantarum Universalem ab Erudstissimo Vro & Botanico Peritissimo D. Joanne Raio

PIRP. Engreat Pr. containes fen Medicamenta que in Offici-

- A. Pinpinella Saliguistio Offic. vulg. seu minor. Park. Raij Cat. Angl. et H. Pl. 401. Usi Fol. et Rad. Przy. 29. in montolis pratis et pascuis. Swall Biunet.
- Diver Indicum five Capficum Offic. Raij H. P. 675. Ulu Schque, sed cause ulurpande. Outmy Denner.
- Piffachia Offic. et Ger. Raij. H. Pl. 1682. Usu Nuclea cum Corr. ejus. In Persia, Syria, Italia, &c. nascitur. 76e Pissachia or Fistick Prite Cree.
- A. Patrinica Office vulg, fol. longo ferrato fl. albo. 7.8.
 Rauj Car. Angl. & H. Pl. 244. Ulu Rad. & Fel. funio Floret. In Prairs udis & Pilafribus. Successor of 251:
- A. Poutegium Regale Offic. latifolium C. B. Raij Car. Angl. & H. Pl. 153. Ulu 161. & Fior. Place, Aq. Spl. & Offic. Still. Junio & Julio Florer. In Locis Pelutribus & aquefis gaudee. Puddung Etals or Jernny Ropal.
- Pulmonaría maculola Ofic. & Got. Raij H. Pl. 488. Ula Fol. cum Fl. r. Prap. 19. Aprili Fl. rec. Jerufalem Comílips, Saige of Jerufalem or Spotted Lung.
- Pyrus Offic. Sativus C. B. Raij H. Pl. 1450. Ulu Finet.
 Apilii & Maio Floret, & por totam lei è altatem Finetus
 maturat. Ibe Bent-Cree.
- Salbia Hout, major Offic major vulg, Perk. Raij H. Pl. 509. Ulu Ed. & Flor. Fræn, ex Flor. Cenf. ex Fol. A. & Syr. Junio & Julio fleret. Common Sange.
- A. Salvia Sylvestris seu Scordonia Oste. Secretais seu Sendatis seu Sendatis Feu Sendana Fol. Salvia J. B. Raij H.Pl. 570. & C... Angl. Ulu Fol. In Sylvis & domeris crescit. Telos.
- A. Saunbutcus Offic fructu in umbella nigro C. B. Raij Cat. Angl. & H. Pl. 1609. Uti Fol. & eyus Gemma, Flor. Cott. & Bacca. Præp. ex Fol. Ung. ex Flor. Aq. et Ol. ex Bacc. Aq. Svr. Svmp. & Comm. vul. Rod. & 1729. Gr.

Satistaga Alba Offic. & Gor. Rail Cat. Angl. & 11

Ulu Explicat Partes istim Plantae que in Arte Pharmacentica

. Probee Norman Plantarum, quibus oulge vocantur Officins nofths. Roamie Baubins Planiavum Historia Universalu

Raij H. Pl. Indicat Historium Plantarum Universitiem ab truper Lidium. 100 & Bounce Verifiem D. Foature Raio

Foamus Parkinsoni Theatrum Botanicum

Dollitum Offic. feu Pulicaris Herba. Ger. Raij H. Pl. 881. Uli Sem. guod in Autumno perficitur. flentwort.

- A: Plantage Angussisolia (20 Ausninguenervia Oster Ray Car. Angl. & II. P., & S. U. J. W. in Pajeus frequene. Ribwort or Eliver Plantain.
- Polygonatum & Signillum Solomonig Offic. La-lat vulg. C. B. Raj Cat. Angl. & H. Pl. 664. Uiu Rad. Pass. In Sylvis quibuldam in Occidentalibus Angliæ gramme Solomong Scal.
- Polygonum & Centinodia Offic & F. B. Raij Cat. Logi. & H. Pl. 184. Oli West. Prep. Ag. in Aros & jux-ta Semina pallim. Anot grass.
- A. Polypodium Offic & Go. Raij Cat. Angl. & H. Pl. 13. Ulu Rad. Prxp. Ol. Stil. In Muris andquis, Truncis arbanm & ad eatum radices. Baippoup.
- 1. Poputius Offic. nigra Ger. Raij Car. Angl. & H. Pl. 1417. Ulu Gemma. Prap. Ol. & Ung. ad Ripus fluviorum Kin Sewels. The Black Bunlar
- Dottum Offic. commune capitatum C. B. Raij H. Pl. 1126. Ulu Rad. & Fol. Piep. Ag. in Hortes colitur.
- Daztulaca Offic. Horrensis Latifolia F. B. Raij H. Pl. 1019. Ulu Fol. & Sem. quod inter quatuor Sem. frigid. minora locum obtinet, Præp. Aq. Syr. & Leboc. Sarven
- Datalplis & Primula Veris Offic. odorata fl. luteo fim-plici. J. B. Raij Cat. Angl. & H.Pl. 1081. Usi Fol. ac Flor. Præp. Ag. Syr. & Conf. Aprili floret in Pafens ubique. Daigles or Coullips.

Explicat Partes istim Planta qua in Arte Pharmacentica quotidie u/urpaniur.

A. Prunella, Offic, Vulgaris Park. Raij Cat. Angl. & H. Pl. vet. Uli Fol. ac Elor. Præp. Aq. Floret Jumo in Pascuis. Self-heal.

Detfict offer mas Perfer F. B. Ruj H. Pl. 1515. Utu Flor. Nucl. & Frust. Præp. Syr. March & Aprils here, Frust. Frust. Schrift here, Frust. Septembri perfectur. The 10cm of 1.

- A. Perstearia seu Dydropiper Offic vols acris su minor. Park. Raij Car. Angl. & H. 17. 182. Usa Fol. Judio & Lagufo floretad Revulos & Bolio. Lake-Weed or Ar-
- A. Petulites Office vulg. rubons roturdiore fol. C.B. Raij Car. Angl. & H. Pl. 2 (9. Ufa Rad. Floret initio Fores & in aquodis ac Palufribm croftic. Definitut wort or Buffer Dur
- 19htt feu Claseriana Offic. Valeriana major, odorata radice F. B. Raij H. Fl. 388. Ulu El. & Rad. in Hens Great Garven Malerian or Sett-wall. crefcir,
- A. Phyllitis Lingua Cerviina Offic. C.B. Raji Cari. Angl. & H.Pl. 135. Ulu Fd. que inter quinque Herbus Capillares enumerantur. In Umbrofis, Uliginifi & Saxofis.
- A. Pilotella, Auticula Butis. Offe. minor vulg. repens Park. Raij Car. Angl. & H. Pl. 242. Ulu Fol. čum Flor. Prap. Aq. & Syr. Comp. Junio & Jalio interdam etiam Maio floret in Pascus arenosas. Doute-Cat.
- Angl. & H. Pl. 878. Ulu Fel. Rad. & Sem. Prep. Ag. & Sir. Julio spicas fort. Ad vias ubique allay by gan A. Plantago Latifolia Offic Vulgaris, Park. Raij Cat. o: Plantain.

Actos Angl. & H. Pl. 1600. Uli Til & cius Genmue, Flor. In sepibus & locis ruderatis.

Sakistiaisa Alba Offic. & Ger. Raij Cat. Angl. & H. Pl. 1084. Ulu I ol. cum Fier. & Rad. granuless circa finem Aprilis & Initio Mari floret, in Pratis & Pafeuis ficeoribus dreftit. Callite Sauffrage.

Ulu Fol. P. &p. Tint. & Ap. Comp. Junio ac Julio Floret, A. Scottlett Offer & Ger. Raij Car. Angl. & H. Pl. 576. in Locis Hamidis. Water Sermander or Scordinn.

Scotzonera Vispanica Ofic. seu Viperaria Ger. Raij H. Pl. 248. Ulu Rad in Hortis nothris Jule mente Florer. Clipers Orald or Servizonera.

- Angl. & H. Pl. 754. Un El & Ral C. B. Raj C. L. gufto in Dumetis & Sepibus. Stown with with or file. wort.
- Setale Offer velgativs Park. Raif Cat. Ingl. & II. F. 1241. Uli Grana. Julio materiant, & in Agent forestern. Bir.
- A. Settetia & Etypetan Officianing ville. C. B. Rail Cat. Angl. & H. P. 29. c. Ulu Hins integra. Vero He. rere incipit & per totam aftarem durat. In Arris, Hortis, & Arois abique occurrit. Erraturaless.
- C. B. Raij Cel. Angl. & H. Pl. 522. Ulu tota Herba. Præp. Aq. 8pt. & Ol. 841. in monichs fed tailus. Lemon
- A. Scrpillum O.E. Vilg. minus Park. Raij Cat. Angl. & H. Pl. 521. Ulu Fol. cum Flor. Prvp. Aq. Mayo et funo Floret. In Montesis et Pajenis Sieciorièmis. Botifist of Chome.
- A. Soldanella, et Istallica matina Ofic. vulg. scu. Volubilis marina Park. Raiz Cor. 1200/20 & H. Pl. -26. Ulu El. in Lattoribus Maris Arca. sc. Sed Isund-weed, Sea Cole-wort or Scottiff Scutty-strafe.



Erra A							
						•	
	1						
	,						
	*						
		*					
							4
,							
	· 0.72701.9+						
	19						
			~v.				

- Solanum Offs. Hortenfe feu Vulgare J. B. Raij Cat. 1820. 8. 11. P. 672. Ulu Fol. Prap. Ag. & Ung. Augusto Septembri mentibus floret, in finiciis & locis ruderatis.
- H. Pl. 222. Usi Fel. in hortis, vineis & Cepibus. Saneljus Luevis Offic. volg. Park. Kaij Car. Angl. Anorth Some Chiffle or Hares Lettice
- Ataechas Arabica Offic. vulgaris Park Raij H. Pl. with a do by or french Mavender
- Arbor tendir Liquorem pinguem inftar Balfami Styrar Lightion Offic. Aceris folio Raij H.Pl. 1681. Chaffum, graveslens & vocatur Liquid Stopat.
- Stillet Offic Latifolium perpetud virens C. B. Raij H. P., 1292. Ulu Cortex cua varijs usibus inservit In Hi
 - A. Cinnacettiini Offic. vulg. Inteum C. B. Raij Cat. Angl. & H. Pl. 265. Ulu Herba Cum Flor. & Sem. Prep. Aq. & Syr. Floret Junio ac Julio; In montibus Borealibus Angliæ. Gania, Infirmia &c. crefeit. The Corte Cute
- A. Camarifens Offic. Narbonensis Ger. Raij H. Pl. 1704. Usi Fel. et Cart. è Radice detracta. Prap. Ay. Maio &
 - Wila Offic. fam. fol. majore C. B. Raij Cat. Angl. & In Areis et Compitis, The Line, Line or 11. Pl. 1694. Uh Her. qui Maio & Junio producuntur.
- Aparine Offic. & Ger. Raij Cat. Angl. & H. Pl. 484. Ulu Foi, vel Planta integra. Prap. Aq. ad Sepes & unter Cleavers or Goofe Oralis Segeres.
- Ageratum & Eupatozium Heliucs Offe. Ballamita fremina teu Ageraum Ger. Raij H. Pl. 264. Ulu Fol. & Iller. Prap. Syr. Pil. & Troch. Baublin or Baublin

- A. Pallis Splutsfiris Offic. sen Agrestis F.B. Raij Car. Angl. & H.Pl. 1444. Usu Frust. è succo ejus Attifice extrahitur. Flores Aprile et Octobri Fractus maturos profert. In Sylvis et Sepibus. The Colifornity or Crali
- Bajotana Latifolia seu major Anglica Park. Raij H. Pl. 539. Usu Fol. Pro Olere usurpatur, urxle à Nothracibus dicitur 190t Partoran
- Detus Offic. Communis Italica C. B. Raij H. Pl. 1502. Ulu Fel. et Bacc. Prap. Syr. & Ol. In Regionibus calidis crescit. The 99 pittle.
- Micotiana major Latifolia C. B. Raij H. Pl. 713. Fumi Nicotiana in Tubo accensa ore attracti et redditi per totam Europam Usus est creberrinus. Ex Virginia, in Circinia Co magna Copia nobis Transportatur.
- Carr. & Feb. In Italia, Gallin &C. provenit. The Ditte Dinig Offic. Sativa C. B. Raij H. Pl. 1398. Usu Nucrs
- Scatiola Offic major vulg. Ger. Raij Car. Angl. & II.Pl. 174. Ulu Fos. Przep. Valent. Scab. Aq. Syr. Simp. et Comp. Junio & Julio floret in Pafcuis, nec minus inter fegetes
- Camarindus Offic. & Gov. Raif H. P. 1748. Un Pul. pa acida nigra quæ in siliqua conditur. In Arabia & utra que India crefeit. 7 be Camarina Crec.
- Clerbascum & Capitus Barbatus Offic. mas Litifolium luteum C. B. Raij Cat. Angl. & H Pl. 1094. Ulu Fol. & Flor. Prup. Tapfimel. & Tapfivalinis. Lungwort, High-taper or Hullein.
- Cithynnalus Offic. Sylvations toto anno folia retinens. F. B. Raij Cat. Angl. & H. Pl. 863. Ulu Fol. In Sylvis & Dumetis crescit. Calood Spurice.
- H. Pl. 821. Ulu Sem. inter Segetes reperitur. Bithti-A. Chlashi Offic. Vaccaria solio Park. Raij Car. Angl. &

- Cethelia Offic. communis ceruleo flore C. B. Raii Cat. Augl. & H. Pl. 535. Ulu Herba integra. Præp. Ag. & Ol. Inful. Æltate floret. in Areis & compitis. Elethallt.
 - Labbanum Offic. Ciffus Ledon folis Laurinis C. B. Raii H. Pl. 1003. E Foliis cujus exudat Liquor quidam Labdanum sen Ladanum dietus. Modus colligendi tradit Dioscorides. Labbanum
- Cletonica Office mus vulgaris supina. Park. Raji Car. Angl. & H. Pl. 851. Ush Fel. cum Flor. Prap. An. fishate floret. In Arenosis & Sabulosis descentur. Park. Speedwel or Fluellin.
- Angl. & H. Fl. 278. Usit Ed. cum For. Prap. Ag. Augu'ro Flore. In montosts Sylvis & Ericetis. Galusti 1983.
- Hyemen durant. Semper virens oft & Frutex reculians H. Pl. 1933. Ulu Fel. Initio veris floret Scheeke per tecans Clifetim Ger, vulg. Baccis albis C. B. Rai Cat. Arri. ? variis Arboribus innafcens. Biffe tot.
- Lachymz ex amoutatis palmitibus extillantes; fic etUanm, Tartains, Uva Paffa, Agrefta. &c. ex quibus varie Pracon-Cittis Cimifera C.B.Raij HP/1612. Ulu Flac etism ratings funt. In Vincia & Horits diligenter colunium In Luiume Findiss maturatu : The Stape or Gine
- Criffetti Offic Spica mutica Ger. Raij Cat. Angl. & H. Pl. 1226. Uli Frum.mta. Seritor in Agris. Willtat.
- A. Cemanerum, Uth Fol. Maio & Junio Floree, In cala-Crifolium paludoliun Ger-Fibrinum Tederamantam Aribus & aquosis. Buck-brang or March Crefoil.
 - Angl. & H. Pl. 259. Ulu Fel. Flor. & Rad. Prxp. Aq. & Cumiago & Karara Offic. vulg. C. B. Raij Core Syr. Martio foret, antequam emittit Folia, ideoque a nonruni & loca rigua provenire amar. Foals Foot or Colts nullis Filius ante Parrem dicitur. Circa Scatturigines aqua-
- A. Maletiana Stacca Ger. & Park. Raij Cat. Angl. &

ricum Offic. Rail H. Pl. 167. Fungus est albicans truncis hujus Arboris adnafcens. Præp. Extract. Troch. & Pil. The Larch-Tree Mulbrame or Agarick

Hoc fape usurpatur pro vero illo Officinarum, (quod Gnaphalium montanum seu Pes Cati Park. elt) Inficci-Sigaphalium minus seu Herba Impia Park. Raij Cae. oribus, flerilioribus & ad vias passim. Detb Inpions or Ingl. & Hift. Plant. 293. Ulu Planta integra. Præp. Aq

Doivenn Offic diffichum Park. Raij Cat. Angl. & H.Pl. 1243. Ulu Grans & ejus Farinz, in Agris seritur. ComDyostepannis niget Offic. & Ger. Raig Cat. Angl. & H. ". "II. Usu Fel. Rad. & Sem. quod usurpatur loco Alsi. in Finteris et Locis ruderatis, Junio Floret. Denbane.

Tietebra minor feu l'ormicularis Osse. Sedum parvum acre il. lurco. J. B. Raij Cat. Angl. & H. Pl. 1041. Ulu Id. Cum For. In muris, 10 dis et ficcioribus, Junio Iloret. Addit-Pepper or Stone-crop Laurenia Offic. Semper-virens fl. luteolo F. B. Raij Car. Angl. & H.Pl. 1587. Ulu Fel. et Bace. InSylvis et Sepibus, Pebruario et Martii initio Floret, Irudus Æstatis fine et adventante Autumno maturitatem confequuntur A. Lichen Arboum fea Hulfins Journanatius Office er nigri sint. Dwarte or Spurise Lawrel

Lotus Ardana et Trifolium Ddaratum Offici Louis Hortenfis odora C. B. Raij H. Pl. 950. Ufu Herba wim Flow Junio er Julio Hortis Floret. Sucet Ctefoil.

& Park. Raij Car. Angl. & H. Pl. 119. Arboribus præfer-

tim Quoreni adnascitur. Cree Lungwott or

Cat. Angl. & H. Pl. 599. Ulu Fol. Flor. Sem. Præp. Ag. Cerl. Syr. Junio ac Julio Floret. In locis ruderatis & ad ad ieres. Common Ballows

Foenum Statcum Offic. Sativum, Park. Rail H. Pl. 954. Ulu Sem quod in Autumno perficitur. If peijugter is

Cornentilla Offic. & Ger. Raij Cat. Angl. & H. Pl.

AND THE PARTY OF T

Craincantha Offic. vera. Park. Raij H. Pl. 933. Usu 617. Ulu Fol. & Rad. Prap. Aq. In Pascuis montosis, & Gummi quod Tragacantha dicitur, & exudat radice incifa vel (ut alij volunt) sponte sua. In Sicilia &c. provenit scioribus. Septfail or Cormentil The True Goats-Chozn Sabiina Offic. vulg. Park. Raij II. Pl. 1415. Ulu Fol. fera est (ut Antiqui Authores volunt) sed verè Conifera. ut primum observavit, & mihi ostendit D. Sam. Dale Vir ta instar Thee apud Ditiores magni Estimantur: & cti-Salvia Absinthites Gor. Raij H. Pl. zrr. Usu E.L. ux cum aqua cocta & parum facchari edulcorata & epo-Prap. Ag. Ol. Stil. & Infus. Hac nec sterilis neque Baccium à nonnullis dicitur viribus, Thee præstare. in Cognitione stirpium apprime gnarus.

Malabathrum et Folium Indum Oke & F.B. surpantur sunt Cinamomi Sylvestris species. Malabatbrum Antiquarum quid sit inter Botanicos nondum convenit. Rail H. Fl. 1563. Ulu Fol. que in nostris Officinis nunc u-Camalapatica or the Indian Leaf

qui Septembri perficitur. In Italia & Germania frequens est. The Sanih of This America Spyllis Offic, fativa C.B. Raij H.Pl. 1456. Ulu Frusk. The Soul or True Servife Tree

Spatula factida & Fpris Offe. Nyris seu spatula sertida. C.B. Raij Cat. Angl. & H. Pl. 1190. Usu Rad. In sepibus & Dumetis. Stinking Sladdon or Sladwon,

Spinachia Offic Lapathum Hortense seu Spinachia femine spinolo. C. BRaij H. Pl. 162. Usu Fol. que inter Plant. 18 Culmarius celebrantur. Colitur in Hortis. Spillacile.

aliarum Plantarum frequenter adnascitur. Ex Succo ejus Dypociffig Offic. Ciffus mas vulg. Park. Raij H.Ph. 1007. do ut Orobinole (cujus species est) eradicibus Genista & e radicibus hujus Fruticis Hypseifix excrescit eodem mo in Sole exiccato Hypsciffus praparatur, quod nostris Anridors (ape ingredicur. Depocities

Dlea Offe, fativa C. B. Raij H. Pl. 1541. Uft Fulf. in Boransphyllis falsd venditur, loco alianum Specierum Othcinalium Valerianarum. The Steek Clafetiall, and by some Lacobs Laddet

tantur & Æstimantur, ac etiam præbent Oleum duorum generum. 1. Quod exprimitur a Frustibus maturis elt 2. Quod ex Olrois immaturis deducitur Ol. Omphacinum Muria conditi antequam plane maturuerint, à variis est-Ol. Olivarum commine. Dyl Dilve, or Sallet Dyl dicitur. The Dlive Crec.


& inter dum ac etiam inter quinque Radices Aperientes locum obtinet. Prap. 14. Floret Junio ac Julio. Satuti vulgo. C. B. Raij H. Pl. 488. Ulu Sem. Fel. & Rad. quæ **Pettolelinum Offe.A**pium Hortenfe feu Petrofelinum

Ufu Fol. Her. Rad. Prap. Ag. Syr. Conf. Eur. Floret Maio tum majus vulgare. Park. Raij Car. Angl. & H. Pl. 505. Confolida major, seu Spunphytum Offic. Symphyac Junio. In Aquofis, ad Rivos & Foffs. Enula Campana, five Descritum offic. & Park Raij Cat. Angl. & H. Pl. 272. Usu Rad. Prap. Constr. An 01. & Ung. In Pratis & Pafeuis humidioribus, Fanto & Julio floret. Elecampane.

. Equificant & Cauda Equina Offic. arvenfe Longi-oribus fetis. C. B. Raji C. A. Angl. & H. Pl. 123. Un Fl. In Arvis. Dozle-tail

Pl. 815. Usu Herba tota. Prap. S/r. Ad Sepes & muros A. Ctyfinum Offer velgare C.B. Raij Car. Angl. & H. Funio & Falso flores, **Deduce Bustari** Fieug & Carica Offic, vulgaris Park. Raij H. Pl. 1431. Ulu Fust.qui post biennium maturantur. In Hortis nostris sape nutritur. The figu-Tree

FILE mas Offic mas non ramofa pinnulis latis densis minutin dentatis. Ger. Raij Car. Angl. & H. Pl. 141. Ulu Rad. In aggeribus sepium provertim umbrosis. Bale



-				
			•	
•				
		•		

- Euphraffa Offic & C. B Raij H. Pl. 771. Uiu Horba cum Flor. Ping. Aq. & Comf. Post mediam ferè aftarem in Authenaum usque Floret. In pascuis sterilioribus. Cochritait.
- 1. fata Offic. major Hortenfis Ger. Raij Cat. Angl. & H. Pl. 909. Ulu Flor. Sem. & ejus Farins. Præp. Sal. & Ag. ex Fl. r. & Silig; In Hortis Evitur & Agris, Aprilj & Maso Florer Julio & Lugufo fructum fort. Oat Det Detug.
- A. Filippindula Offic. vulg. Park Raij Cat. Angl. & H. P., C23. Ulu Rad. Junio & Julio Floret, in pratis & Pul. cuis montosis. Dop-Wilt.
- A. Delifotus Offe. Germaniæ C.B. Raij H.Pl. 951. & Cat. Angl. Uiu Fel. & Flor. Præp. Empl S. & Comp. Junio & Julio floret, ad sepes & in arvis. Common Belitot.
- A Baffurtium Aquaticum Offic. aquat. vulg. Park. Raij Cat. Angl. & H. Pl. St. 6. Ufu Her ba tota. Præp. Aq. & Spr. Julio & Augutho Mensibus florets, in Aquis & ad Rivules frequens. Utater Crestes.
- A. Stille Formitta Offic. ramofa major pinnulis obtufis non dentatis. C. B. Raij Car. Angl. & H. Pl. 149. Utin Rad. Ubique nimis frequens. Semiale feeth, or Community Arabed.
- A. Frayaria Offic vulgaris C.B. Raij Cat. Angl. & H.Pl. 600. Ulu Icl. & Fruch. Prap. Aq. Syr. & Tink. In Sylvis & Horris, Janio Fruchus fert. Common Strawberry.
- Calega & Ruta Capratía. Offic. vulgaris Park. Raij H. Pl. 911. Víu Herba cum Flor. Præp. Aq. & Conf. Junio & Julio in Hortis floret. Goats Rue.
- A. Callium Office Inteum C. B. Raij Cat. Angl. & H. Pl. 482. Ulu Herba tota. In Pascuis siccioribus passim. Cheese Rening, or Pellow Ladics ISO-straw.
- A. Genista Offic. vulg. & Scoparia Park. Raij H. Pl. 1723 & Cat. Angl. Usu Fol. Flor. & Sem. Præp. Aq. Conf. & Sel. Florer eirea finem Aprilis & initio Min. In Pascuis menolis trequences occurrie. Estiman Istoonn.

- Spicotiana Petum & Tabacum Offic. Hyofcyamm Intens Ger. Tabacco Anglicum Park. Raij fl. Pl. -15. Ul. Fol. & ejus Sucens. Prap. Aq. Ung. & Ol. English Cobacco, and by some Pellow Den-batte.
- Migella, Melanthium & a quibusdam Gith Offic. Nigella st. minore simplici candido C. R. Raij H. Pl. 10-11. Usi Sem. que valde odorata sunt. Migella or festivell felosor.
- A. Duniphata & Nenuphar Offic. major alba. C. B. Raij Car. Angl. & H. Pl. 1319. Ulu Flor. & Rad. Præp. Syr. Junio & Julio floret, in Lacubus et Fluviis. The Clatter-Lilly.
- EDELITA Office vulgi odore Citri F. B. Raji H. Pl. 572. Ulu Herba tota, Præp. Aq. & Spr. Syr. acConf. Junio Julio & Augusto Floret. Dautin.
- Agela Offic. vulg. C. B. Raij H. Pl. 644. Ulu Sem. quæquatuor Seminibus frigidis Majoribus referuntur. The Agelan.
- Dentha Offic. anguilifol, spicat. C.B. Raij H. Pl. 532. Usa Herba cum Edr. Præp. Aq. Spt. Syr. Conf. & Ol. Infuf. & Stil. Julio Floret. Satuten or Specat Hint.
- A. Detruitalis, Bonn Hemicus & Teta Bona Offic. Birtum Bonn Hemicus dictum Raij H.Pl. 195. Ulu Fel. Præp. Syr. & Mel. Aprili & Maio Floret, ad vias & in Oleraceis. All group, or English Detruity.
- A. Betum Athamanticum Office fol. Anethic. B. Raij H. Pl. 432. & Cat. Angl. Ulu Rad. Maio & Junio Floret. In Prais & Pafeur nonnunquam occurrit. Betwo or Spignell.
- 16 A. Diffetolium Offic terrefiris rulg. Ger. Raij Cat. Anzl. & H. Pl. 348. Utu Fol. Junio Floret in Pascus ubique: Partow or Hillott.
- Bolus Offic. nigra vulg. Park. Raij H. Fl. 1429. Ulu Fridt. qui in Angufto maturescit, Prap. Syr. et Diamoron.

- A. Passinaca sativa tenuisolia Ger. Raij Cat. Angl. & H. Pl. 465. Usu Sem. & Rad. gux Culinarix sunt. In Agris sericur. The Continon Catrot.
- A. Detecpiet Anglorum Lobellj Ger. Raij Cat. Angl. & H.
 Pl. 209. Uli Herba integra. Inter Segetes & in foloreflibili fleriliore palitim. Joanfly-piett or Joanfly breakextens
- A. Querring Offic. vulg. Gen. Rai, Can. Angl. & H.Pl. 7605.
 Unit 191. Glandes conumque Calves. Prep. Ar. In. Sylvas & Sepibus ubique vulgacitinna. The Date.
- A. Quinquelesstants: Pentaphyllun oss. vulg. Ger. Raj Car. Angl. & H. Pl. Stat. Uh. R. Let Rad per totam ashatem Flo-et, ad semitas pashin. Five-senbed Crass
- Raphanustr enfis of. val., P. R., H. P. 804. Un Ral. & Sem. The Batter Radiff.
- A. Raphanus Sylvestris for Rustianus Ofic.531volt. Isu Armoracia in ites. J. B. Ray II. Pl. Sec. & Cor. Angl. Ulu Rad. Pres. Aq. S. & Comp. Spc. & Fyr. Comp. Locis humidis. Hotic-Radish.
- A. Kapa Offic. fativa rotunda C. B. Raij Car. Angl. & H Pl. 800. Ulu Rad. P. xp. 4q. & Syr. In Agris & thoris pathin feriur. The CULINGA.
- A. Bhanning Catharticus et Spins Cethina Offi. folintius leu Spins Infebria vulg. Park. Raif Cat. sugl. et II-P. 1625. Ula Bacca que coliquitur in fine Septembris aut initio Offobra. Prep. Succ. et Syr. In Sylvis & Sopibus Crefcit. The Common Patryling Chour or Ishakilan.
- Rosa Ag. Mais & Jun'o Hortis luxuriar. The Callite Rose.
- Ruhia Cincocum offic & Ger. Raij H. Pl. 480. Un Rad.In Horris colitur. Baddet.

The contract of the trace of th

- aregosis frequenter occurrit. Common Droun. Recaing, or Pellow Ladies Bed from. 4 12. CILL HETON
- Pl. 1057. & Cat. Angl. Ulu Fol. Præp. Ag. odirata. In Anglia fed 1210. Egustopy, or Busiked Cranes. tum fol. ad Myrrhidem accedente majus I. B. Raij H. A. Erranium Holthatum Offic. Cicutæ fol. Moscha-

Definium & Kalisteum Offic. Vulg. majus Park. Raij H. Pl. 541. Usu Fol. ac Sem. Floret Junio & Julio, in Hortis Creldic. Bassell.

- A. Cohiegiafilli Offic. sive Lingua Serpentina, Park. Raij Cat. Angl. & H. Pl. 126. Usu Planta integra. In Pritis humidis aprili & Maio luxuiint. Apperid Colliffie.
- Digannum Offic. Anglicum Ger. Raij Cat. Angl. & H. Pl. 539. Usu Fol. cum Flor. Præp. Ol. (iill. Floret Junio ac Julio in dumetis & ad Sepes. Caill Hanjozam.

Pocenta und Offic. & Park. Raij H. Pl. 693. Usu Rad. Flor. & Sem. Maio Florer. The Hale Hoonly.

Polium Hentanum offe. Rail H. Pl. 121. Usu Fol. & Flor. In Italia & Gallia crelcit. Polley Douite Mattuttum Hottenfe Offs. & Go. Raij H. Pl. 825. Um Fel. (quæ in Acetarijs po supe afurpantur.) ac Sem. Præp. Aq. Floret Maio & Junio. Edvott Crest.

Naffuttium Indicum Park. Raij H. Pl. -8-. Ostu Flores qui grati funt saports & a quibusdam ad Viola reducuntur. Prap. Cenf. in Horts nutritur. The Judian Crelg.

- Kummulatia Offic. vulg. Park. Raij Cat. Angl. &. H. Pl. 1099. Ulu Fel. Prwp. Aq. à Maio mente in multam æstatem Floret. In Folfis & Pratis humidis. Bottep. wort or Herbstwo-pelice.
- A. Bepeta Offic. major vulg. Park. Meniba Felma feu Cattaria, Ger. Raij Cat. Angl. & H. Pl. 548. Usu Herba reta. Junio & Julio stora, ad sepes invenitur. Bepp or

The Windshipmen

Re H. Pl. 345. Ulu Foi. Junio Fiolice III From English

Hepl. 113. Ulu Planta tota. In Aridis Locis. Ä

Angl. & Eup or Chalice Bols. Drylapathum Offic Lapathum Fol. acuto plano. C. B. Raij Cal. Anzl. & H.Pl. 1-2. Ulu Rad. & Sem. In incultis pastim. Syarp pointed Dock.

Poconia Kaemina Offic. & Ger. Raij H. Pl. 644.Usu Rad. Fler. & Sem. quod Julio ad maturitatem pervenit. Prap. Ag. S. & Cemp. Syr. Cenf. The Kemale Poconp.

Danaver album Offic. Hort. sem. albo. C. B. Raij Hist. Plant. 853. Ulu Capita cum Sem. Præp. Syr. in Horis colicur. The Collitte Sarven Poppy,

- Jonpavet Rhoens Ger. seu Erraticum. Osic. Raij Cat. Angl. & H. Pl. 855. Ulu Flor. Prep. Aq.S. & Comp. Syr. & Conf. floret Junio inter Segetes pattim. Red Poppy or Curr-Rafe
- Parietatia Offic. & Diosc. C. B Raij Cat. Angl. & H. Pi. 789. Ulu Fol. quæ locum tenent inter Herbe quinque Emollientes, Pray, Aq. In Maris & locis Ruderatis. [3011]. Ä
- A. Baranpehía Chilgaría Ger. Raij Cat. Angl. & H. Pl. 189. Usa Herba tota Marrio Flores. In Mario, Il oribus & Aronofe. Cammon Collittom-Orales.
- A. Durannehia Rutacea Folio Gen. Rail Car. Lugl. & H. P. Ulu Hoba integra florer April & Meio in Mura Icors & Siccioribus. Ruis Leav's Clyitlaw Graff.
- Ballittatu Hortensis Osse. Latifol. sativa Ger. Ray Cat. Long. St. H. Pl. 410. Ulu Rad. & Sem. in Agris & Hortis coltuir. Carden Parlinip.
- 19819 Offe Cucurbita major rotunda flore luteo fol. afpero C. S. Kail H. Pl. 396. Ulu Sem. quæ lunt e numero quatuor frigedierum majerum. The Pinnpion, called by the Vulgar, The Delon

H. Pl. 515. Ulu Fol. Sem. & Flor. qui Authos vocantur. Prap. Aq. Spr. Conf. et Mel. Aprili et Maio ac deinceps Flo-Rounarinus Offic. Hortenfis angultiore Fol. C.B. Raij ret. Roscinary.

Buth Tingeruppoffice on that It as

Bole.

Ros Solis Offic. fol. retundo Ger. Raij Cat. Angl. et H. Pl. 1105. Usa Herba tota. Prap. Aq. In palustribus. Ros Solis or Sun-dew. Ribes Offic vulg. fructu rubro Ger. Raij H. Pl. 1485. Usu Fruct. Przp. Rob et Syr. Aprili mense aut serus proveris Conflitutiones floret. Fruct. Fruct. matturantur. Red Currang.

- perficient. Prap. Syr. & Diaman. In Sopibus & Dumetis frequencer occurrit. The Brannlife or Blatk-Wetty Angl. & H. Pl. 1539. Uhi Fel. & Frack. quod in Autumno Rubug vulg. Offic. major frudu nigro J. B. Raij Cat. K
- Kubus Jdatus Offe. frinofes Fuelu releto. CB. Raij Cat. Angl. & H. Fl. 1645. Ulu Frust. P. 22. 44. & Syr. In. Monthus. Wall & & Septentrionvillus Anglia Greleit. The Dindberry, Framboife of Rasistry.
- H. 1. oct. Un R. Linwisk and a Aperientes launtener in Supplies & Duncies Miles 199111 of Littlifers. Ruffers & Benfers Offs. & Ger. Feel Cat. Angl. & bjoom.

fall Ung. In He is your more about a set of State or State or State or Rate. H. P.L. 874. Via I. L. & S.m. Prant Leg. C of. O. Sal. & Su-Ruta Datterffs, Caralgo I. Fort, letital. C.B. Raij

Salvia Dort, white of any a Andelea J. R. Raij H Pl. 415. Undeld High Sign Signs of Sagreet

- A. Sappliatis Office valg. Perk. Reif Cen. Angl. & H. Pl. 999. Uli Fel. Floret Junes, Julio & a. in Archeris & ad Ripas fluvictum provenit. Soppolitati.
- A. Pifum Offe. Raij H. Pl. 893 & Car. Angl. Ulu Sem. m Agris feritur. Deale,

					- *
19					
			+1		



Colassium Ashim Office Rais H. Pl. 62. In Mari re-

Corallium Kuthuuti Offic. & Ger. Raij H. Pl. 60. est Frutex sub aqua Marina generatus. Presp. Syr. & Sal. Red Corall. Cojiandrum Offic, vulg. Park. Raij H. Pl. 470 Usin Sem. Hzc Planta dum viget halitum gravem prabet, reficeata autem suavem acquirat odorem. Cojiander.

- A. Copplied Office five New Avellana fylveftris Park. Raij Cat. Angl. & H. Pl. 1279. Usin Frust. & Lign. in Sylvis & Sepibus. The Datell Mut.
- A. Ctíthmum feu Feniculum marinum Offic. minus C. B. Raij Cat. Angl. & H. Pl. 457. Ufu Herba tota. Foliin Muria condita in acetarijs, &c. Szpe ufurpatur. In clivis & Rupibus maritimis frequenter occurrit. Rock Sammire.
- A. Crottis Offic. verus fativus Autumnalis Park. Raij Cat. Angl. & F. Pl. 1176. Usa stamina Florum. Prep. Sp. Syr. Ext. Spec. & Empl. Floret Septembre. In Essexia & Comitatu Cantabrigae permultum colitur. Sastitum.

Cucunis Hortenfis Offic. Sativus vulg. C. B. Raij H. Pl. 645. Uju Fruct. cellinis 82 Sem. Officinis quainter quator frigidiora majora enumerantur. Satoen Colu cumber.

Cultunis Spidefitis feu Afrins Offic. & I. B. Raij H. Pl. 647. Excels Succi Fructus hujus Planta dicient Efiterium. Fructus fert in Autumno. Califoto de la Cultura feut.

Cupteffus Offic. & C. B. Raij H. Pl. 1406. Use Fruct. feu Conos. in Hortis crefcit. The Cpptls: Ctt.

A. Cuscusta Offic. seu Cassusha Ger. major C. B. Ra Cat. Angl. & H. Pl. 1903. Usu Herba tota cum sloribus Herbis adnascitur passim; Junio, Julio, & Angusto sloret.

Deveta Cettefftis Offic. vulg: Park. Raij Cat. Angl. & H. Pl. 567. Ulu Herba integra. Præp. Aq. Conf. & Syr. Menfe Aprili Floret, ad Sepes. Gill 130 by the Ground. Cuntifoof, Ale-hoof, and Sepes. Gill 130 by the Ground.

collebous Albus Offe. & Ger. Raij H. Pf. 168. Usurad. Przep. Vin. Califte Cellebore.

pelleborus Ilívet Offic, niger verus Ger. Raij H. Pl. 697. Ulu Rad. Præp. Ext. Syr. & Oxymel. in Hyeniem Floret. The True black Pellebore.

1902minum Sativum Offic. vulgare feu Sclarea. Park. Raij H. Pl. 543. Ulu Fol. & Flor. Præp. Ag. & Sp. Junio & Julio Floret. In Hortis feritur. Satven

A. Dozminum Spluestic & Deculus Christi Oste. Sylv. Lavendulæ Flore. C. B. Raij Car. Angl. & H. Pl. 848. Usu Planta integra, in glareosis frequens. English

A. Opperferni Office vulgaris C. B. Raij Cat. Angl. & H. Pl. 1018. Ufu Fol. ac Flores. Prap. Aq. Frust. Ol. Simp. & Comp. Floret Falso in Dumeris & ad Sepes. St. John's 771 int.

Dyffeptig Offic. vulg. spicatus angultifolius I. B. Raij H. Pl. 516. Ulu Fol. nc Flor. Præp. Aq. Syr. & Ol. Srill. Floret Julio& Augusto, Crescitque in Hortis. Satuett

A. Depatica & Lichen rulg. Offic. & Park. Ray Cat. Angl. & H. Pl. 124. Ula Fol. In alignofis & umbrofis ad rivulorum ripas. Common Liver-wort.

A. Jacobea Offic. vulgaris laciniata. C.B. Raij Cat. Angl. & H. Pl. 284. Usi Fel. In Pascuis & ad vias passim provenit. Seggeum or Ragwort.

Jasminum Offic. vulgatius fl. albo C. B. Raij H. Pl. 1599. Usu Flores. Præp. Aq. Balf. & Elsen. seu Ol. per Infus. per integram serè æstatem Flores. Wisite Jasmine or Jestime.

Lanium Rubium Offic. vulg. folio subrotundo slore rubro, Park. Raij Cat. Angl. & H. Pl. 559. Usi Fol. ac Flor. Primo vere & deinceps per totam æstatem Floret. Small Dead Mettie, or Red Atchangel.

Labendula Offic. major feu vulgaris Park. Raij H. Pl. 512. Ulu Fel. & Fleres. Præp. Conf. Aq. & Sp. Junio ac Julio menfibus floret, in Hortis. Labendet.

Laurus Offic. vulgaris C. B. Raij H.Pl. 1668. Ulu Fel. & Bace, que maturantus feb finem Autumni. Præp. Elegt. Empl. Ung. & Ol. Stil. The Billy Cute.

A. Leng Paluffig Offic. fou aquatica vulg. Pårk. Raij Cat. Angl. & H. Pl. 117 Ulu. tota Herba. In aquis flagnantibus. Duchs-theat.

Levisticum Offic. vulg. ver. Liguilicum vulg. folio. Apij I. B. Raij H. Pl. 427 Ulb. F. I. Rad. & Sim. Pl. R. Aq. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. Aq. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. S. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. D. & Ol. Sril. Ælta* Ulb. F. Sim. Pl. R. D. & Ol. Sril. Ælta* Ulb. F. D. & Ol. Sril. Ælta* Ulb. F. D. & Ol. Sril. Ælta* Ulb. F. D. & Ol. Sril. Ælta* Ulb. Ella* Ulb. F. D. & Ol. Sril. Ælta* Ulb. F. D. & Ol. Sril. Ælta* Ulb. F. D. & Ol. Sril. Ælta* Ulb. Ella* Ulb. Ella*

Lillium Album of the constant of the constant

A. Lifting Conduction of the Pek. Raij Car. Angl. & H. Pl. 667. The property of the Conf. Major mentle floret, in the property of the Calify, or Soat-Lifty.

Lincula mala Offic Raif R. P. 18486. Ulu Fract. com. Cont. & Sem. Pract. Ag. Syc. Conf. C. S. e. Fract. ex. Higherit & Leilia ad nos deferences. The Leimmon.

A. Linun fabrum Offic & Gen Ray Car Angel & H. Pl. 10-2. Un Sem. Prap. O', per Engrish in Agels fabrum.

Raij Cat. Angl. & H. Pl. 503. Ulu Sem. In Agrorum marginibus ficcioribus. Flor. Maio & Junio & comill or Occuminell.

A. Lujula & Acctofella Offic. Trifolium acetofum vulg. Park. Raij Car. Angl. & H. Pl. 1098. Usu Fel. Floret A-

Us. Kad. In Curiosorum Horris colitur. Sow Bread Cat. Angl. & H. Fl. 1905. Off Helio, de Augusto Horer. Merris adrafeitur passinn; Junio, Julio, & Augusto Horer. Epciament five Arthanita office nail ext.

Fruct. & Sem. Prap. Succ. Syr. Vin. Rob: Miva, &c. Fructicat in Octobri. The Quilite-Cree. Cydonia fen Cotomea malas Raij H. Pl. 1452. Uju

- Tynoglossunt Offic. majus vulg. Ger. Raij Cat. Angl. & H. Pl. 489.Us. Fol. & Rad. Præp. Pill. ad vias ubique Junio Floret. Hounds Countie
 - Cynostatos & Cynotthodos Offic. Rols fylv. vulg. flore odorato incarnato. C. B. Raij Cat. Angl. & H. Pl. 14-0. Usu Fruck. & Spong. Prap. Cons. Autumno fru-ticat in Sepibus. The Dog Role, Poepp, or Celifa
- Cyperus Longus odoratus Offic. & Park. Raij Cat. Amgl. & H. Pl. 1294. Ulu Rad. In palustribus. The Sweet Cyperus, or Einfillh Salangal
- vulg. Park. Raij Cat. Angl. & H. Pl. 1058. Uiu Planta integra. Ad lepes ubique. Heth. Robert.
- Park. Raij Cat. Angl. & H. Pl. 910. Usu Rad. cum Succo Ejustdem. Prap. Syr. Comp. Floret Julio & Semen sert Septembris, varijs locis apud nos colitur. Liquorsth.
 - A. In agris & Hortis nimis frequens, Common Dougs Orals, Quick Orals, or Couch Grals. Oramen Canimun Offic. arvense seu Gramen Diosc. C. B. Raij Cat. Angl. & H. Pl. 1255. Usu Rad. Prap.

Exoflutaria sive Alva Crispa, Offic. Raij H. Pl. 1484 Ufu Frust. Floret Aprili, fructum fert Maio, quem Junio & Julio perficit. The Gogberty

Transta vel Punica Osc.& Ger.RaijH.Pl.1462.Usi Poma cum Cort. Prap. Syr. The Pomentanate-Tree.

yevera Offic. corymbola nigra. Park, Raij Cat. Angl. & H. Pl. 1505. Usu Fol. Bacc. Lig. & Gummi. Floret Autumno, Baccas fert hyeme, Arboribus passim nascitur. Climing or Watten Ave

Thirty our interest Præp. Ag. Balf. & Ellent. feu Ol. per liftle ver interest. Per darmie and Imperatoria offic. five Afrantia vulg. Park. Rail H. ffertiout, and by some erroncoully Bellitoly of Pl. 436. Usu Rad. Prap. Aq. in Hortis nutritur. Ba-

Just Nottens Offic. purpurea feu vulg. Park. Rail H. Pl. 1180. Usu Rad. Præp. Ag. & Succ. floret Maio & Junio, in Horis frequenter. The Posttyle flower De

- A Jugians Offic, sive Regia vulg. C. B. Raij Cat. Angl. & H. Pl. 1376. Ulu Nuces virides dum tenera sunt & Fructum ante initium Septembris apud nos non perficit. Cort, interior Arboris. Præp. Aq. Simp. & Comp. & Mel.
- A. Juniperus Offic. vulgaris baccis parvis purpureis. I. B. Raij Cat. Angl. & H. Pl. 1411. Ulu Bacc. Lign. & & Ol. still. Fruttus maturescit Anno secundo, Autumno & Gummi quod Sanivataca dicitur. Prap. Ag. Sp. Conf. tunc cæruleo tingitur colore. In Ericetis crescit.

220. Usu Fol. & Sem. quæ annumerantur inter Semina Lactura Offic. fativa non capitata J. B. Raij H. Pl. quatuor Figida qua minora dicuntur. Prap. Aq. OUL den Lettice

- major I. B. Raij Cat. Angl. & H. Pl. 392. Folia Verno tempore, cum Aceto Sale & Olco jucunde commedum-A. Lactica Aguina Ger. Valeriana campestris inodora tur cum alijs acetarijs Herbis. Inter segetes. Lambs
- A. Leng Offic. vulgaris C. B. Raij Cat. Angl. & H. P. 904. Usu Sem. Sentur in Agris. Lentills, and by some
- A. Lamiunt Alluum Offic. vulg. album seu Archangelica slore albo Park. Raij Cat. Angl. & H. Pl. 559. Ulu Flor. Prap. Conf. & Aq. Mense Aprili & Maie floret, ad Sc. pes. Earlite Archangel, or Dead Bettie

Etuca Offic. Latifolia alba fativa Diosc. C. B. Raij E. Pl. 806. Ulu Sem. Rockett.

prili ac principio

Scommen.

Lupinus Offic. vulg. semine & flore albo sativus J. B. Raij H. Pl. 906. Usu Sem. Æstate floret, & Autumno Sem. perficit. Lupines.

Lupulus Salictarius Offic. & Raij Cat. Angl. & Hift. Plant. 156. Usu Flor. Augusto in sepibus passim abundat. Lentiscus vulgaris C. B. Raij H. Pl. 1579. Usu Lig-num & Gummi quod è caudicibus hujus Arboris exudat & Baffich vocatur. The Baffich-Cree Dajorana & Samplucum Offic. vulg. æftiva. Park. Ruij H. Pl., 538. Usu Fol. ac Flor. Prxp. Aq. Ol. Stil. & Comf. Floret Julio & Augusto. Sweet Partolam.

Mandraydra Offic. fructu rotundo C. B. Raij H. Pl. 668. Usu Fol. & Rad. in Hortis Botanicis colitur. The Pale

A. Pasifium Atount Offic. Marrabium album valg. C.B. Raij Cat. Angl. & H. Pl. 555. Usin Ed. Præp. Aq. & Syr Floret Fallo, ad vias publicas & in locis ruderatis. Callite Hogelfound.

lens. C.B. Raii H. Pl. 520. Usin Herba tota, in Augusto store october Bastich or Bastich Cine, and by some Harum Offic Sam Juchus fen Maram Mastichen reco-

neutories in caonter imponitur. Angrian Ballich Ulu Fel. & 18 18 regratifimo & vegeto funt odore, & fler-Syarung Syriacum vel Creticum Park. Rail H. Pl. 538.

- Ang S. F.P. 355. Ulu Herba cum Flor. Prap. Aq. Junio & Matticotia & Parthenium Offic Vulg. Park.. Raij Cat. JOSE WELLE & Hound Sepes & locis ruderaris. If Chirefth.
- Erringuntt Offic. marinum Matth. C. B. Raij Cats ,84. Ust Rad fecat. aut Condit. In Livo. sign of the 184. On Rad. secar. auchonic in a sign, we seem to the safeture Seasoup our Crysty.



The second second				
yetti i				
ž.				
		•		
			,	
				•
3.5				
No.		6		
				/

- LEURIOIUIII Hortenfe Offic. angulfifolium majas. C. B. Ray H Pl. 401. Util 181 R.id. & Flor. que funt e nume o quatuoi Florum andialium. Presp. exclein cum Borragine. In ho. tis no. 11 is no. 11 i
- A. Diffilled for Confolida media Offic vulgaris Fl. cerumleo. P. N. Raij Car. Angl. & H.P. 575 Ulu Herba integra. Majo mense Florer. In sylvis & Praiis hum dis. Diffic.
- A. Diuth Palluis Offic. major vulg. Park. Raij Car. Angl & H.P. S. Un Phyastim occurrit totaque unate viget. Syppheneus Putte.
- A. Dupittialmum Offenam longe diffat a Buphthalmo Doje, & Math. reliis major Ger. Raij Cat. Angl. & H. Pl. 75°. Usa Fol. & Flex. In pratis & pascuis Majo & Junio Floret. The Stort Dasie, or Dr. Epc.
- A. ISHTHE Offer arborescens C. B. Raij Cat. Angl. & H. Pl. 1653. Usu Flor. & Ligna Præp. Ol. Stil. In varijs locis Auglia. The ISON CICE.
- A. Calamintha Offic. altera odore Pulegij folijs maculofis Park. Raij H. Pl. 5'9. & Cat. Anzl. Ul'i Herba tota Prep. Aq. Spec. Diacalaminth. Simp. & Comp. Svivi. & ful.5 Autumnum Floret. In aggeribus fepium & folo Iteriliore & arenofo provenit. Hae pro vulgari in Officinis mafris femper ufurpatur & forfan non immeritò. Itota
- A. Calanilitha vulgaris fl. magno I. B. Roij H Pl. 569. & Cat. Angleviget Altate ad aggeres follarum & lerium. Common Calamint.
- A. Calannis Aromaticus Offic. Acorus verus feu Calamus Aromaticus Offic. C. B. Raij Cat...Angl. & H. Pl. 1613.
 Usa Rad. Locis humidis gaudet. Plantam hanc non esse Verum Calamum Aromaticum seu odoratum Antiquerum Botanici eruditiores confentiunt, sed Acorum verum Dioscoridir. The Sweet suessilling Islay, or Calannus.

Calendula Offic. Caltha Vulgaris C. B. Raij H. Pl. 237. Ulu Fol. & Flor. Prap. Ag. Acet. & Conf., ver totaun fene Miller in Horis. Earthern Barringolo.

- Cathuning & Cinita Offic. Hortenlis Park Raij H 21.9 ... Ulu Florenzap. Syr. & Conf. in Hortis luxuriat ac in Julio & potter Floren The Clove July flower.
- A. Catum Office vulgare Park. Raij Cat. Angl. & H. Pl. 446. Use Somewood unum est e quatuor calidis majoribus Ca minatisis. Parp. Aq. Spr. Ol. Stil & Confect. In palufiribus & pratis humidis apud nos invenitus. Catamaics.
- A. Catyonly lata Offic. vulg. fl. parvo luteo F. B. Raij Cat. Angl & H. Pl. 606. Ulu Rad. & Fel. In Maio & Junio floret ad fèpes & in dumetis. Deth' Bellutt, or Allena
- A. Castested Office Gen. Raij Cat. Angl. & H. Pl. 1582. Ulu Frace. Coun. Cort. ejus. In montibus & acclivibus. The Ch. E-stite Test.
- Cataputia Pajot vulg. seu Ricinus Offic. & Park. Raj H.P. Callia Nuclei Fructús, The Steater Sputige, or Palmachtillí.
- Cataputia Pinoz Office fou Lathyris major Hortenfis. Park. Ray H. Pu. S. 6. In Hortis frequenter provenit, Dof. Baccar, ad no. 10. Fel. ad no. 6. The Satuen Spurge.
- Ceveus Conifera Libani. Folijs Laricis C. B. Raij H. Pl. 1407. Arborem hanc ene Cedrum Libani in facris Literis tantopere celebratum cujus thus erat ad Temblum Solomonis edificandum minimè dubium est. The Cevat of Libanus.
- A. Centaurium Pinus Offic. A. purpureo F. B. Raij H. Pl. 1.92. & Cat. Angl. Ulu Fol. ac Flor. Præp. Aq. Ext. & Sal. Jolio mense floret in pascuis siccionibus frequenter Smail Centaury.
- Dactylus Ossc. seu Palma Dactylifera major vulgaris, Johnst. Dendrolog. Raij Hist. Plant. 1352. Usu Fruck. & Ossc. Præde Biaphæn. Fern. The Palm, or Date-Eree.
- A. Daucus Offic. seu Pastinaca sylv. tenuisolia Diosc. C. B. Raij H. Pl. 465. & Cat. Angl. Usu Sem. quæ locatur and garagement. The Pastinia Garage and the Catter of the

- Ceba Offic. vulg. C. B. Raij H. Pl. 1116. Usu Fol. & Rad. In Hortis colitur. Floretque Junio & Julio. Duyons.
- A. Cetalus fructu nigro Offic. major ac Sylvefris frusctu fubdulci nigro colore inficiente. C. B. Raij H. Pl. 1538. & Cat. Angl. Ulu Fruct. cum Nucleis & Gummy, Pixp. Aq. Sp. Vin. & Syr. The Diack Chetiy.
- Cheretolium Offic. Cerefolium vulgare sativum. Ger. Raij H. Pl. 430. Um Fel. & Sem. Præy. Aq. Maio Flores. Sweet Chevil, or Sis.
- Chameding & Triffago Offic. Valg. Park. Raij H. Pl. 527 Ulu Fol. cum Floribus. In Hertis Junio & Julio Florer. Sermander.
- A. Chamaemetium vulg.Offic odoratifimum repens FI. fimpl. I. B. Raij Cat. Angl. & H. Pl. 354. Ulu Fel.& Fler, Prap. Aq. Ol. Still. & Injul. Syr. & Sal. In passuis uliginosis Julio Floret. Common Chamoniil.
- **Chamaenneium Romanum** FI. pleno Offic. repens odoratuhmum perenne FI. multiplici. I. B. Raij H. Pl. 353. Uli Flor. Przp. exdem cum vulgari.In Hortis Luxuriat. Douthle filotuted Chamoniil.
- A. Chamachitys & Iva Aribritica Offic. Intea vulg. fol. trifido C. B. Kaij Cat. Angl. & H Pl. 57., Ulu Herba tota. Pix., Pil. In agris reltilibus Junio & Julio Floret. Oronna-3nic.
- A. Chelidonium majus Offic. & Ger. Raij Cat. Angl. & H. Pl. 8;8. Uti Fel. cum Fierib. & Rad. Præp. Aq. ab Aprili mensc in multam æstatem Floret. in ruderatis & Sepibus umbrosis. The Steattet Celandine.
- A. Clyclidonium minus offic. Schrephularia minor feu Chelidonium minus vulgò dictum I. B., Raij Car. Angl. & H. Pl. 579. Ulu tota Herba cum Radicibus. Floret Martio & Aprili in Pratis humidis, & ad Sepes. Hoc longè differt (in specie Methodica) a majore. Hoc Rammoule affinis, illud Papaveri Siliquos congener est. The Lester Celandine, or Poste-wort.

lere Attatem floret in Hortis. Oat vent Bat pgold,

Cannabis Offic. Sativa C. B. Raij H. Pl. 158. & Cat. Angl. Ulu. Sem. in Agris serieur. Denip.

Pl. 140. Usu Fol. qux inter quinque Herbas Capillares locum tenet.Pixp. Syr.Officinis nosfris Adianthi veri seu Caoulgare Offic. Trichomanes mas. Ger. Raij Cat. Angl. & H. villi Veneris succedaneum oft. English Black Bat Capillus Cleneris & Adianthum Migrum

rupus quotannis per totam Galliam alialque nationes pra-Cavillus Cleneris Verus Officinis Menspelis, unde Syparatur, & in magni Bitimatione habetur. Adianibum Fol. Coriandri C. B. Raij H.Pl. 147. The True Buton-

line, &c. luxuriat, & Gemme Florum Aceto conditæ ab illis Capparís Offe. Spinola fructu minore fol. rotrinlo. Troch. & Ol. Comp. In locis arenofis & ruderatis Italics, Gal-C. B. Rail H. Pl. 1629. Ulu Cort. præcipuè Radicum. Pian locis nobis transportantur, & in frequenti usu tunt.

Germanicum. C.B.Raij Cat. Angl. & II.Pl. 1450. Ulu F.l. & Flor. Incipit florere Maio & per totam feie Estatem Caprifolium Offic. Perichmenum non perfoliatum uxuriat, in sepibus frequenter. Doncy-suchse or

Usu Fel. cum Capitibus & Sem. Prxp. Aq. Syr. & 303. Ulu Fol. cum Capitivus & sem. 11212. 117-31. Sal. sub Antumnum Semen perficit. In Hortis colitur. Carding Beneditus Offic. & Ger. Raij H. Pl.

Park. Raij Cat. Angl. & H. Pl. 312. Ulu Fol. Sem & nonnunquam Rad. ad Agrorum margines & in aggeribus Fossamm frequencer occurrit. The Bilkie or Ladies Carduns Barine Offic. albis maculis notatus vulg.

Carthamus Cuicus Offic. & Ger. Raij H.Pl. 302. Usu Kor. & Sem. Dos. ad 3vi. Prap. Syr. Elect. & Spee Floret Julio & Augusto. Hx Floret nonnullis venduntur pro Cross & ab illis Crosm Hiffam. dicitur, Baffath Saffton.

que occurrit. Willa Carrot. Deng Leonis & Taraxacum Offic. latiore fol. C. B.

Raij Cat. Angl. & H. Pl. 244. Usu tota Herba, per totam affatem floret in Pascuis ubique. Danpelion Dictaminus Creticus Offic. & C.B. Raij H. Pl. 37. Ulu Folia. Dittany of Erete

Dictamus Albus offe. Fraxinella Ger. Raij H. Pl. 698. Ulu Kad. Frakinell, or Bastard Dittany.

Angl. & H. Fl. 767. Ulu Fol. & Flor. Junio & Julio floret in Arenolis. Por Gloves. Digitalis Offic. purpurea fol. afpero C. B. Raij Catal.

Dozanicum Offe. Rad. Scorpij C. B. Raij H. Pl. 274. Ulu Rad. In montibus prope Genevam copiose provenie. Leopards-bane Draco Octba Ger. Raij H. Pl. 373. Usus ejus in acetarijs irequentifimum est. Catranol

1211. Ulu Herba tota cum Rad. Prap. Aq. Dragong. Dracontium Oste major vulgaris Park. Raij H.

Dulcamara & Solanum Lignolum Offic. Amara lunio & Julio Floret, in sepibus præsertim juxta aquas. Dulcis Ger. Raij Cat. Angl. & H. Pl. 672. Ulu Fol. & Rad. Witter fiveet or Coloody Right-flade

A. Culling Officin. seu Sambucus Humilis Ger. Kouniaurn Diose. Usu totin Fruten. Floret Junio & Jelio ad vias inque Agrorum limitibus & cometerijs. Dante wort, Calaste Ethium Officia. vulg. Ger. Raij Cat. Angl. & H. Pl. 498. Ulu. Fol. & Flor. Æstate stotet in Arvis Bettioribus & muris frequenter occurrit. Alivers Bunfals

Elatine Offic. seu Veronica fann. Fueblij. Ger. Raig Car. Angl. & H. Pl. 759. Ust Fol. Serids post mediam afta-tem floret in arvis. Bounte-seuve, or Female

In Hilpania, Italia, Gallia, &c. passim in agris seritur. Floret Junio & Julio mensibus atque subinde in siliquis Cicet Offic. Sativum Ger. Raij H. Pl. 917. Ulu Sem. lemen producit. Chich-Peale, or Cicers

attinis, illud *Papaveri Siliqusf*i congener **elt. The Lettee** Œchmune, or Dite-maje.

Cichotium Offic. Sylvestre seu Officin. C. B. Raij Cat. Angl. & H. Pl. 255. Ulu Fol. Flor. Rad. & Sem. quæ inter 4. frigida minora enumerantur. Prap. Cons. & Syr. ad agrorum margines sponte provenit. Æstate & post solsti tium Floret. Succory

Raij H. Pl 1392. Coccum fert Maio mense quod duobus ubbus infervit, nimirum Medicina (ex quo celebris Confectio Cheemies Offic. Hex aculeata cocciglandifera. C. B. Alkermes nomen habuit) & Land tingendis. The Scat.

45 t Ulu Fol. & nonnunquam Rad. Junio & Julio Floret in aggeribus terrenis & ad folias. Demilock. Cícuta Officiulgaris major Park Raij Cat. Angl. & H. Pl.

Cituis Offic. Malus Citria feu Medica Raij H.Pl. 1654. Ulu Fruchus Prap. ex Cort. Ol. still. Elect. Syr. & Condit. The Citron Cree Cochicaria Britanica seu Barina Offic. Fol. sinuato C. B. Raij Cat. Angl. & H. Pl. 823. Usu & Præp. eædem cum Hortense. In salsis maritimis plurimis in Locis. Sca Scurbygalg

A. Colocinthis Offic. fructu rotundo minor. C. B. Raij H. Pl. 642. Usu Medulla arida Fructuum. Dos. ad 3j. Colocinthis Offic. fructu rotundo minor. C. Prav. Trock. Coloquintiva Lochsteatia Hottensis Offic. fol. subrotundo C. B. Raij H. Pl. 882. & Cat. Angl. Usu Herba tota. Plæp. Aq. Sp. Cons. Syr. & Sal. Floret Aprili & Maio in Hortis. Sardeir Seurvy-Grafs.

Offic. & Park. Rail Cat. Angl. & H.Pl.65. Ulu tota Plants Cotallina Offic. Muscus marinus seu Corallina alba Accrescit Rupibus aqua marina opertis juxta litus ubique

					* -	18
					*	2 * 4
						-
						t
			*			
			*			
				,		
						4
						*
	•					
						54
				,		, , , , ,

	•		
A			
<u>:</u>			
	~		
•			
· {		·	

Gieg Park. Taxifolijs Raij Hist. Plant. Pag. 1;94. In Us lunt Folia & ejus Resma quæ duplex ab hac Arbore elicitur. Prima, Liquida Osficinis Terebinthinam Veneram (at nostratibus Argentoratensem, i.e. Strajburg) falso vocant. Secunda, Sieca quæ Tbus zamulatur. Lignum Deal vocatur Præp. Ol. Still. Hæc non est Abies, qua vulgo utuntur in nostris Osficinis, attamen multo præstantior est. The Beth. Leaved. or Leaven.

Abites Officiarum. Picea Park. Raij Hig. Plant. 1396. Ulu Felta. Hac Arber copiofam Refimam ultione præbet; Imprima, cum tenuioris Confiltentiæ fit; Pix Liquida feu Catt vocatur; fed cum decoctione dephlegmatur, craffelcitque, Pix Sicca, vel Pittly, nobis dioitur. The Compiler, or Pittly, craffelcitque, pix Sicca, vel Pittly, nobis dioitur. The Compiler.

Guintanum mas Officia, mas angultifolium majus, C. B. Raij H. Pl. 371. Ulu Fol. Præp. Aqua & Conferva. In Hortis colitur. Southeth-Clood.

Abrotantin frmina, Chamxeypariffus Offic. Abrotantin tem. vulg. Park. Raif H. Pl. 359. Ulu Fol. & Sem. Praxp. Ag. & Confero. Lavender Cotton.

- A. Abfinthium Vulgare Offic. & Park. Raij H. Pl. 166. & Catal. Anglia. Ulu Fl. ac Semen. Præp. Aq. Spr. Vinum, Syr. Treeb. Ol. Stil. & Sal. Crefcit in Ruderatis & Incultis locis frequenter. Common Clom-Clood.
- Marinum Anglicum Park. Raij H. Pl. 370. & Cat. Angl. Uf. 180. ac eius Summitates. Præp. eædem cum vulgari. In Maritiniis Eingliff Sea (Cloum-tellom), and commonly (but taility) called Konnun (Cloum-twood).

Charthium Romanum verum. Tennifolium Ponticum Galeni Ger. Raij H. Pl. 367. The True Roman Angl. Ulu Fol. Rad. & Sem. Præp. Syr. & Conserv. In practis & pascuis & pascuis. Quinnon Sopret.

Stretofa rotundifolia Hortensis C. B. Raij H. Pl. 180. Stree usurpatur soc vulgaris, & Prap. exdem, cui praffat; acidiore sapore prædita. Roman, or strenth

Acous Adulterinus Offic. Itis paluftris lutea, Ger. Raij

Allull Offic. & Ger. Expostor Diofe, Raij H. Pl. 1125.

- A. Allitte Offic. minor fou media. Ger. Raij H. Pl. 1030. & Car. Angl. Ulu Herba Integra. Common Chick-Cleed.
- A. Althea Offic, vulgaris Park. Raij H. Pl. 602. & Cat. Angl. Ulu Syr. Rad. & Fol. Ex Quinque Herb. emolient. Prw. Sem. & Ung. In paluftribus marinis. Butifigental orangement.

Animi vulgare Offic. & Ger. Raij H. Pl. 455. Ulu Sem. Common Bilhops-Caleed. Ampadalus Offic. Dulcis & Amara, I. B. Raij H. Pl. 1519. Ulu Nuclea. Prxp. Ol per Expressionem. Primo Vere

Floret. The Almand Trec.

A. Amagallis mas Offic. Flore phæniceo. Park. Raij H.

Pl. 1023. & Cat. Angl. Ulu. Fol. & Flor. in Areis & Hortis. Agale (2011). Pl. 1023.

A. Anagallis Aquatica, sive Becabunga. Offic. Ger. Ray H. Pl. 852. & Cat. Angl. Usin Fol. vel planta integra. In rivulis & ad Fluviorum margines exit. Common 13500k-lime.

Aucthum Offic. Hortenfe feu vulgare. I. B. Raij H. Pl. 415, Ulu Fol. Flor. & Sem. Præp. Ol. Stil. & per Inful. In Hortis nutritur. Auct, or Dill.

Angelica Offic. Sativa. Ger. Raij H. Pl. 454. Ulu Fol. Rad. & Sem. quod in Julio & Augusto perficitur. Præp. Ol. Stil. Garben Angelica.

Anistum Offic. & Ger. Raij H. Pl. 450. Usu Scm. qui inter quatuor Calida majora Carminativa locum possidet. Præp. Aq. Sp. & Ol. Sril. Annist.

- A. Atyentina feu Potentilla Offie. & I. B. Raij H. Pl. 617. & Cat. Angl. Usu Planta integra. Circa semitas & loca aquosa frequens. (Ilistanta integra. Circa seluct. Illesta.
- A. Angl. Usu Rad. ac-potissimum ejus Cortex. Prap. 4q. In

Attiplet Offic. olida Gev. a quibufdam Valvaria dicitur. Raij H. Pl. 198. & Cat. Angl. Ufu Fol. & Sem. juxta parietes & in locis ruderatis crelcit. Attinkting Drach.

A. Auent Offic. Vulg. five alba. C. B. Raij H. Pl. 1253. &c Cat. Angl. Ulis Calinare potius off quam Officinale. In Agris scrittur. Datts.

A. Autuntia Malus Offic. Raij H. Pl. 1658. Usu Fel-Flor. Sem. Cert. ac Medulla Fructuum. Præp. ex Flor. Ag. & Ung. e Cort. Syr. Condit. & Ol. Stil. Flores & Fructus toto Anno profert. The Dianing Cute. Lialauffum Offic. Balaufium Fl. pleno majus C. B. Raij H. Pl. 1643. Hac Arbor Flores (vulgo Lialauffi-1169 vocantur) folum fert, & illi tantum iu Use lunt. The greater wild, or bouble bioffomed Johnegranates of rec

Balfanita Mas Offic. & Ger. Coffus hortensis sex. Mentha Corimbistera I. B. Raij H. Pl. 363. Usu Herba 10-ta. Prwp. Aq. odorata. Costinato.

- A. Wattballa Major Offic. & Ger. Personata, sive Lappa major, aut Bardana. C. B. Raij H. Pl. 332. & Cat. Ang. Usu R. A. Sem. & Fol. In incultis passim. Street Idlic. Dock, or Clot-burn.
- A. DSellis fylv. minor C. B. Confolida & Symphisum minimum. Offic. Raij H. Pl. 3+9. & Cat. Angl. 'Viu Tota Planta. Vere primo & deinceps per totam æftatem Floret. In Pratis & pascuis passim provenit. Common
- A. Dethetis Offic. & Park. Raij H. Pl. 1605. & Cat. Angl. Ulu Cortex interior Ligni, & Frust. ex quo Prap. Conf. Spr. Rob. & Troeb. In Sepibus sylvolis & ripis Fluviorum. Frust. in Autumno perficitur. Bathetise a quibuldam Piptetidues.

Letta Alba Offic. & Ger. Raij H. Pl. 204. Ulu Rad. Fol. & Sem. Præp. Succ. una elt e quinque Herbæ Emollientes. The Ullite Beet.

Ista Rubta Offic & Ger. Raij H. Pl. 204. Un Rad.
Fol. & Sem. Rev Istet.

Isetonica Ger. Cylvestris seu vulgaris Offic. Raij H.

- H. Pl. 1186. & Cat. Angl. Ulu Rad. Ad Ripas Fluviorum. Pellow Water flag, or flower-de-luce.
- A. Abianthum Albam Offic. Ruta muraria, seu Salvia Vita, Park. Raij II. Pl. 146. & Cat. Angl. Usu Fol. quainter quinque Herbas Capillares locum occupant. In muris antiquis, & pomtibus Lapideis nascitur. (Ulliste Buirben-Pair, Chall-Rue, or Cent-Chot.
- A. Abianthum Aureum Offe. Polytricum Aureum majus, Park. Raij II. Pl. In Pharmacopeia nominatur, fed in Usuraro venit. In Sylvis udis ac paludosis crescit. Solvis udis ac paludos crescit. Solvis or Selvis natur.

Agung Castus & Atter, Officia. Vitex solijs angusioribus Cannabis modo dispositis, I. B. Raij H. Pl. 1695. Usi Sem. Leu potius Finctus, Prxp. Troch. The Chaster

- A. Agrifolium feu Aquifolium, Park. Raij H. Pl. 1622. & Car. Angl. In Ufu raro occurrit, attamen Fol. & Bacca ad dolorem Colicum & tormina intestinorum mire concucunt, & ex Corticibus hujus Arboris Vifeus paratur. The Dolly: Ette.
- A. Agrimonia Ger. & Eupatorium Gracorum Offic. Raij H. Pl. 400. & Cat. Angl. Ulu Fel. Prap. Aq. Sal & Syr. Æltate Floret in Palcuis & ad sepes. Agrimonp.

Alten Agyptiaca, Abel Moseb sen Mosch Arabum. Semina valde odorata sunt, & cum Odorarijs frequenter in Isto. Beneik Septil.

A. Alchimilla Offic. & Ger. Raij H. Pl. 209. & Cat. Angl. Ulu kol. In pascuis & pratis montosis. Ladics: Gauttle. Alkekeingi & Halicacabumi, Offic. Solanum Vesicarium, C. B. Raij H. Pl. 681 Usu Fuct. qui Autumno maturant. Prap. Troch. Calinter-Cherries.

1. Alliatia Ger. Raij H. Pl. 792. & Cat. Angl. Usu Fol. Vere storet, in sepidus & aggeridus fossarum. Jack by the yedge, or Sawce alone.

Reffessionionionis nequence occurric. Cumoen, Reffessionis

- A. Apium Paluftre seu Ospeinarum C. B. Raij H. Pl. 447, & Cat. Angl. Usu Fol. & Sem. quæ inter 4. Calida minora locum tenet. & Rad. quæ ad Aperientes refertur Præp. Aq. & Ung. in Paludosis & Foshis præsettim maritimis,
- A. Aquilegia Offic. Flore fimplici. I. B. Raij H. Pl. 706. Ulu Fol. & Sem. in Sylvis montosis invenitur. Colum-

Atistolochia longa & rounda, Offic & Ger. (nam in Folijs parum disserunt) Usu Rad. Long and Round. Routed Bittly-wort.

- A. Arthemisia offic. Mater Heibarum. Ger. Raij H. Pl. 372. & Cat. Angl. Usu Herba tota. Præp. Ag. Syr. & Sal. Ad Sepes & agrorum margines. Duspussurt.
- A. Atum Offic. Vulgare Ger. Raif H. Pl. 1208. & Cat. Angl. Ufu Rad. & Fol. Præp. Ag. Pulv. comp. & Facula. Primo vere exit in Sepibus & Umbrofis. Cuckow-Point, Clarke Robin, and Atom.

Afatum Ger. Vulgare Offic. & Park. Raij H. Pl. 207. Usu Fol. Dos. ad No. 9. infuta & expressa. Rad. Dos. ad 3j. Afatabateta.

Asparagus Sativus Offic. & Ger. Raif H. Pl. 683. Usu Rad. que e quinque Rad. Aperient. Turiores seu Germina in cibis magni æstimantur verno Tempore. Spartagus, or Specialic, corruptly Spattow-Erasis.

A. asperusa sive Asperala odorata Ossic. & Park. Raij H. Pl. 483. & Cat. Angl. Usu tota Herba. Majo mense Florer in Sylvis & dumetis præsertim montosis. Casodoogoos.

A. Applementant, five Scolopendra, Ceterach, Offic. C. B. Raij H. Pl. 139. & Cat. Angl. Ulu Fok (qux inter quinque Herbas Capillares enumerantur) Prxp. Syr. In Saxorum crepidinibus ing, venultis muris provenit. Ceterach, Spilenwolf, or Diffwall.

Atripler Sativa. Offic. & Ger. Raij H. Pl. 191. Usu Fol. & nonnunguam Sem. Satten Diach.

- A. Betonica Ger. fylvestris seu vulgaris Osse. Rais H. Pl. 550. Usu Fol. & Flor. Præp ex Fol. Ag. Syr. Simp. & Comp. Empl. & Sal. ex Flor. Comf. In Sylvis & Sepibus Junio & Julio Floree. (UN000 Betonly.
- A. Betonifa Pauli & Verenica mas Offic. Veronica pratomlis Serpyllifolia. C. B. Raij Cat. Angl. & H. Pl. 848. Ulu Fol. Majo & Junio Floret. In pratis hunidis & Sylvis crescit. Little smooth fluellin, or Paulis Detains.
- A. Istula Offic. & Ger. Raij Cat. Angl. & H. Pl. 1410 Ulu Fol. Cort. & Liquor ab illa extillante. In Sylvis humidis & Ericetis colligitur. The Istich Ctt.
- A. Dissistat offic. major vulgaris Park. Raij Cat. Angl. & H Pl. 180. Ulu Rad. In pratis humidis. Stake-weed, or Dissistant.

Bail H. Pl. 1064. Ulu Sem. & Lanugo que Gossipium le B. Osficinis vocatur. The Cotton Shtill.

Jövrtang Offic. Hortensis Ger. Raij H. Pl. 493. Usu Rad. Fol. & Flores (qui inter quatuor Cordiales Famosos recensentur) Præp. ex Fol. Aq. & Syr. ex Flor. Syr. & Conf. In Junio Julio & Augusto Florer. **Jövrtange**.

Nottys & Anthrofia Offic. Vulg. Park. Raij H. Pl. 196. Ulu Fel. Prap. Syr. In hortis colitur. Dak of Jerusasem.

Branca Eletina Offic. Acanthus fativus. Ger. Raij H. Pl. 1326. Ulu Fol. Brank-Eletine, or Bears-Breech.

Astallica Sativa. Offic. capitata alba Ger. Raij H. Pl. 794. Uli Fol. & Sem. Ontven Cabbance.

A. Dityonfa alta Offic. afpera feu alba baccis rubris.
C. B. Raij Cat. Angl. & H. Pl. 659. Ufu Rad. Dof. ad 3j.
P. Ep. Aq. Comp. & Facal. In Sepibus crefcit. Collite
Dityony.

A. Detyania Pligta Offic. lavis feu nigra racemofa. C. B. Raij Cat. Angl. & H. Pl. 650. Ulu Rad. In Sepibus & Dumetis nalcitur. Dinth Bryony.

•			1
			1
			_
			÷
			- 2
			, X
			,
		Y X	
			* مدر

Fucus five Alga marina latifolia vulgatissima Raij Sycops. 2. H. Pl. 70 & Cat. Angl. Quercus marina Ger. Fig. 1567. Park. Fig. 1293. It grows to Stones and Rocks on the Sea-Coasts plentifully. The Post common broad-leaved Sca-Wrack.

Fucus five Alga latifolia major dentata Raij Cat. Angl & H. Pl. 71 Synops. 2. It grows in deep Seas. 232000-10011: et Sea-Callacti.

Fucus angustifolius vesiculis longis Siliquarum æmulis Raij H. Pl. 73 Fasc. 6 & Syn. 4. It's found on the Seashoar in many Places. Mr. Ray's Marrow leaved Whack, with Pod-like Bladders.

Alga Ger. fig. 1569. Fucus marina feu Alga marina graminea Park. Fig. 1292. Raif H. Pl. 75. Fasc. 75. & Syn. 5. Very common on the Sea-shoar. Sints-White.

Alga marina platiceros porosa. I B. Fucus tela lineam Sericeámve textura sua æmulans Raij Syn. 4. H. Pl. 71. & Cat. Angl. 11. It's often cast on the Sea Shoar. Silkett: Chiach.

Lichen marinus Ger. Fig. 1566. Fucus marinus Lactuca marina dictus Park. Fig. 1293. Raij Cat. Angl. & H. Pl. 77. Synops. 6. It grows to Rocks and Stones in the Sea. Dyffer-Green.

Corallina Muscosa seu Muscus marinus tenui capillo Spermophoros L. Pl. Phyt. Tab. 168. Fig. 3. I first observed this (A. D. 1691.) on the Sea-shoar in Essex. I fine leaved Coralline Bass.

Conferva geniculata minima nostras L. Pl. Phyt. Tab. 84. Fig. 9. Observed by Mr. Doody, in the Ponds and Ditches, about the Fort at Sheerness plentifully. Geniculated Conferva, or Crow-Silk.

Muscus arboreus supinus marginibus pilosis Raij Cat. Angl. & H. Pl. 116. Syn. 14. It's found on many Trees, but especially the Ash, flourishing in May, Mr. Ray's Black Spangle headed Grey Hoss, with sine curied Leaves.

Muscus crustaceus bracteolatus slavus. crusta aut Lichenis modo Arboribus Petrisve adnascens slavus Raij Cat. Angl. & H. Pl. 116. Syn. 14. On old Trees and Stone-Walls frequently. Pellow Spangled crustaceous Poss.

Muscus terrestri vulgari similis lanuginosus Raij Syn. 18. Alpinus ramosior erectus slagellis brevioribus lanuginosus L. Pl. Phyt. Tab. 47. Iig. 5. It's very common on the Mountains in Wales. Mr. Lloy d's Dairy Earth Doss.

Muscus montanus gracilis ramosus viticulis longioribus glabris, L. Pl. Phyt. Tab. 47. Fig. 4. It's found in Mountainous shady places. Dr. Plukenet's stender-branched Mountain Moss.

Muscus palustris gracilis summo ramosus, parvus Stellaris C. B. & L. Pl. Pkyt. Tab. 47. Fig. 6. In Boggy Places, Small upgight Bogg-Poss, with starry Cops.

Muscus Filicinus Ger. Fig. 1561. Park. Fig. 1310. Major C. B. pin. 360. Raij Cat. Angl. & H. Pl. 112. Syn. 13. Its very common in shady Woods. Ferm Moss.

Muscus trichodes medius capitulis sphæricis Raij in App. Syn. 243. Its found early in the Spring on Dry Ditch Banks in Heathy Ground. Mr. Doody's Golds-socks, with Ronno. heads.

Muscus trichoides foliis capillaceis capitulis minoribus Raij Syn. in App. 243. In moist Hearby Grounds. Mr. Doody's fine-leaved Goldslocks, with small Deads.

Muscus trichoides capitulis oblongis restexis Raij Syn. App. 243. It grows frequently on the Walls about London. Mr. Doody's small Goldslocks, with Red Stalks, and Green pendulous Deads.

Muscus trichoides parvus soliis Musci vulgaris capitulis longis acutis Raij Syn. 243. Its very common on Walls and old Houtes. Mr. Doody's common Gold locks, with small crooked beads.

Muscus trichoides hirsutior capitulis longis restexis, pediculis brevibus insidentibus Raii Syn. 243. On Brick Walls frequently. Mr. Doody's Pairy Goldslocks, withpendulous peads, and very short soot stalks.

Muscus trichoides minor capitulis longissimis Raij Syn. in App. 243. On shady Banks. Mr. Doody's small Goldstocks, with very tong and slender heads.

Muscus trichoides minus foliis ad caulem convolutis, capitulis subrotundis reflexis Raii Syn. in App. 244. On moist shady Walls. Mr. Doody's Goldslocks, with Leaves growing like a Bulbous Root.

Adianthum aureum medium foliis tenuissimis, capitulis erectis acutis Raij Syn. 237. Its found in Heaths and Boggs. Mr. Bobart's fille leaved Boggs py Death. Dols.

Muscus pennatus capitulis Adianthi Raij Syn. 236. In Boggs and moist places. Mr. Bobari's Pennated Goldiscks.

Muscus Saxatilis Fæniculaceus Anglicus, I sirst observed it A. D. 1692. on the highest Rocks in CharleyForrest, in Leicester-shire. Rock-Dast.

Muscus trichoides lanuginosus Alpinus. I sound this about Midsummer, A. D. 1692. on some Rocks in Leicester shire. Small Dairy Hountain Goldslocks.

Muscus Lichenoides arboreus repens foliis rotundis D. Doody Musci cauliculis compressis altera species Ejust. Rais Syn. 244. Its frequently found early in the Spring, growing to Trees in shady Places. Mr. Doody's creeping black-headed Liverwort Hose, with Round-Leaves.

Fungus holosericus arboreus inserne lamellatus Raij Syn. 244. Its sound on Trees with the other. Mr. Doody's Small Lamessated velvet Husproom.





						35
						1
					4	. :
					÷	
						3
						į.
						(1)
•						-1
						* -
				- 4		
			53	00/5		
						,
		•				
	,					
						and the same of th
					٠	



